

MRA template for prokaryotic or eukaryotic genome sequences

Article title

Draft genome sequence *Chryseobacterium treverense* isolated from human clinical sources

Authors

Analisa Coppens^a, Isabella Fregoso^a, Rachel Haakma^a, Kimberly Habon^a, Grace Rodrigues^a, Tricia A. Van Laar^a

All contributed equally, organized by last name

Affiliations

a. Department of Biological Sciences, California State University, Stanislaus

Running title

Chryseobacterium treverense Genome Assembly

Corresponding author's email address

Address correspondence to Tricia A. Van Laar, tvanlaar@csustan.edu.

Abstract

We report the draft genome sequence of *Chryseobacterium treverense* is isolated from human clinical sources. The genome is 2,379,279 base pairs and contains genes enhancing bacterium survival and evasion of the host's immune system, therefore amplifying broad antibiotic resistance.

Announcement

The Gram-negative *Chryseobacterium treverense* is isolated from Trier City, Germany (49.7501° N, 6.6372° E) ([5](#), [6](#)). It is a yellow-tinged, aerobic, non-spore forming, and non-motile bacterium in soil, plants, human clinical specimens, and food. It potentially causes urinary tract infections and pneumonia ([1](#), [2](#)). Genomic sequencing provided taxonomic revisions, accurately classifying it in the genus *Chryseobacterium* ([3](#)). Data from JGI allowed scientists to genomically sequence *C. treverense*, enabling researchers to investigate its impact on eukaryotes ([4](#)). *C. treverense* was aerobically grown on Columbia agar, supplemented with 5% sheep blood, and preserved in fetal calf serum by lyophilization ([1](#)).

The draft genome of *C. treverense* was generated at DOE JGI using Illumina technology. Illumina HiSeq-2500 1TB platform sequenced a 300 bp insert standard shotgun with the 150-bp paired-end format. Illumina Std PE generated 10,312,988 reads totaling 1,546.9 Mbp. The raw data was filtered using BBduk. Reads with more than one "N," quality scores averaging less than 8, contigs less than 1 kbp, or reads shorter than 51 bp were discarded. Using BMAP, the remaining reads were aligned to masked reference genomes of humans, cats, and dogs, discarding those exceeding 95%. The filtered, normalized reads were assembled using SPAdes v3.6.2 ([7](#)). The final draft, based on 1,500.0 Mbp of Illumina data with coverage of 625.3X, contained 10 contigs and 10 scaffolds, totaling 2.379 Mbp. The final draft was annotated according to the standard JGI Microbial Genome Annotation Pipeline. The genome was 17% contaminated with a complete genome of 99.97% ([8](#)). DSMZ will provide details on growth and DNA isolation.

The genome had 2,379,279 base pairs. Scaffold N50 is at 1,189,639 base pairs. The average fold coverage was 4.3 with a GC content of 39.46%. There were 2305 genes and 2193 coding sequences with 5 rRNAs and 37 tRNAs (Table 1). The probability of *C. treverense* being a human pathogen is 0.227 with zero matched pathogenic families and 0.68% input of proteome coverage; it is predicted to be non-pathogenic ([9](#)). Virulence factors include genes enhancing the probability of bacterium survival and protecting from the host's immune system (VFDB 2022). Using CARD, 95 loose hits and one strict hit were detected in *C. treverense* ([10](#)). The bacteria exhibits broad antibiotic resistance, particularly to fluoroquinolones, glycopeptides, oxazolidinones, and tetracyclines. The results of CRISPRCasFinder displayed 10 analyzed sequences and 3 CRISPR sequences with evidence ([11](#)). Comparing our results to JGI, no CRISPR sequences were listed. antiSMASH (v 7.0). Analysis revealed type I PKS cluster, indicating possible terpene production.

Feature	Findings
Length (bp)	2,379,279
Status	complete
No. of contigs	10
GC content %	39.46
No. of genes	2305
No. of coding sequences	2193
No. of rRNAs	5
No. of tRNAs	37

Table 1: Genome Statistics

Figure 1: This phylogenetic tree of *Chryseobacterium treverense* was constructed using the Type Strain Genome Server (TYGS) and Genome BLAST Distance Phylogeny (GBDP) analysis.

Data availability statement

This Whole Genome Shotgun project is deposited in GenBank under accession no. [FORQ01000001](#). The version described is the first version, [FORQ01000001.1](#).

Acknowledgments

The work (proposal DOI[:xxx]) conducted by the U.S. Department of Energy Joint Genome Institute (<https://ror.org/04xm1d33>), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under contract no. [DE-AC02-05CH11231](#).

References

1. [Yassin AF, Hupfer H, Siering C, Busse H-J. 2010. *Chryseobacterium treverense* sp. nov., isolated from a human clinical source. *Int J Syst Evol Microbiol* 60:1993–1998.](#)
2. [Siddaramappa S, Narijala A, Viswanathan V, Maliye C, Lakshminarayanan R. 2019. Phylogenetic insights into the diversity of *Chryseobacterium* species. *Access Microbiol* 1:e000019.](#)
3. [Info - *Chryseobacterium treverense* DSM 22251. \[https://genome.jgi.doe.gov/portal/ChrtreDSM22251_FD/ChrtreDSM22251_FD.info.html\]\(https://genome.jgi.doe.gov/portal/ChrtreDSM22251_FD/ChrtreDSM22251_FD.info.html\). Retrieved 20 February 2025.](#)
4. [Info - *Chryseobacterium treverense* DSM 22251. <https://genome.jgi.doe.gov/portal/ChrtreDSM22251/ChrtreDSM22251.info.html>. Retrieved 4 March 2025.](#)
5. [Prakash O, Verma M, Sharma P, Kumar M, Kumari K, Singh A, Kumari H, Jit S, Gupta SK, Khanna M, Lal R. 2007. Polyphasic approach of bacterial classification - An overview of recent advances. *Indian J Microbiol* 47:98–108.](#)
6. [Polyphasic approach of bacterial classification - An overview of recent advances - PubMed. <https://pubmed.ncbi.nlm.nih.gov/23100651/>. Retrieved 24 February 2025.](#)

7. [Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen I-MA, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard operating procedure of the DOE-JGI Microbial Genome Annotation Pipeline \(MGAP v.4\). Stand Genomic Sci 10:86.](#)
8. [Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2023. CheckM2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. Nat Methods 20:1203–1212.](#)
9. [Cosentino S, Larsen MV, Aarestrup FM, Lund O. 2013. PathogenFinder - Distinguishing Friend from Foe Using Bacterial Whole Genome Sequence Data. PLOS ONE 8:e77302.](#)
10. [Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA, Edalatmand A, Petkau A, Syed SA, Tsang KK, Baker SJC, Dave M, McCarthy MC, Mukiri KM, Nasir JA, Golbon B, Imtiaz H, Jiang X, Kaur K, Kwong M, Liang ZC, Niu KC, Shan P, Yang JYJ, Gray KL, Hoad GR, Jia B, Bhando T, Carfrae LA, Farha MA, French S, Gordzevich R, Rachwalski K, Tu MM, Bordeleau E, Dooley D, Griffiths E, Zubyk HL, Brown ED, Maguire F, Beiko RG, Hsiao WWL, Brinkman FSL, Van Domselaar G, McArthur AG. 2023. CARD 2023: expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. Nucleic Acids Res 51:D690–D699.](#)
11. [Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCasFinder, an update of CRISPRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins. Nucleic Acids Res 46:W246–W251.](#)
12. [Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw BR, Metcalf WW, Helfrich EJN, van Wezel GP, Medema MH, Weber T. 2023. antiSMASH 7.0: new and improved predictions for detection, regulation, chemical structures and visualisation. Nucleic Acids Res 51:W46–W50.](#)