

**Copyright
by
Ryan Patrick Hannigan
2023**

The Dissertation Committee for Ryan Patrick Hannigan certifies that
this is the approved version of the following dissertation:

Stranger Things at the LHC

Committee:

Christina Markert, Supervisor

Peter Onyisi

Richard Hazeltine

Chandrajit Bajaj

Stranger Things at the LHC

by

Ryan Patrick Hannigan, B.S.

Dissertation

Presented to the Faculty of the Graduate School of
The University of Texas at Austin
in Partial Fulfillment
of the Requirements
for the degree of
Doctor of Philosophy

The University of Texas at Austin

May, 2023

To Jaynee, who unquestionably is the reason why this document exists.

Stranger Things at the LHC

Publication No. _____

Ryan Patrick Hannigan, Ph.D.

The University of Texas at Austin, 2023

Supervisor: Christina Markert

Among the most mysterious of the four fundamental forces is the strong nuclear force. Responsible for both the binding together of nucleons within an atom, as well as the

Contents

List of Figures	viii
List of Tables	ix
Chapter One: Introduction	1
1.1 What is fundamental?	1
Chapter Two: Experimental Apparatus	5
2.1 The LHC	5
2.2 The ALICE Detector	6
Chapter Three: Λ Production In and Out-of Jets: Analysis Overview	11
3.1 Objective	11
Chapter Four: Analysis Details	12
Chapter Five: Systematic Uncertainties and Cross-checks	13
Chapter Six: Results and Discussion	14
Chapter Seven: Summary and Conclusions	15

List of Figures

1.1	A diagram depicting the particles we currently believe are fundamental within the so-called “Standard Model” of particle physics.	2
1.2	A coffee cup with the Standard Model Lagrangian density printed on its side. Please ignore the “+ h.c.” term following the $i\bar{\psi}\not{D}\psi$, it is the result of a small lapse in judgement from the mug makers.	3
2.1	A diagram depicting the LHC with its various main detectors shown underground. Illustration by Phillippe Mouche, from BBC News.	6
2.2	A 3-D schematic of the ALICE detector, with labels for all of the sub-detectors. Note the humans-for-scale in the bottom left of the diagram. .	7

List of Tables

Chapter One: Introduction

The purpose of this chapter is to transform a scientifically literate reader who may be vaguely familiar with the idea of a “quark” into one who can understand the motivation and techniques behind the analysis presented in the rest of this thesis. If you are an extremely educated physicist who regularly performs lattice quantum chromodynamics calculations in their head, I would also encourage you to read this chapter it its entirety as it may contain egregious errors that need to be corrected.

1.1 What is fundamental?

The answer to the question “What are the fundamental building blocks of our universe?” has changed drastically over the course of human history. The idea that all matter is composed of smaller, uncuttable pieces has been around since 5th century BCE when Greek philosophers Democritus and Leucippus first introduced the concept of an atom. While this idea was initially ignored in favor of more theological descriptions of our universe, over the span of a millennium or two the atom became the de facto indivisible building block of nature. However, everything changed around the turn of the 20th century when scientists like Rutherford and Chadwick determined that the supposedly indivisible atom was composed of even smaller particles, eventually named protons and neutrons. The notion that protons and neutrons were unbreakable was relatively short lived, as not even half a century later the deep inelastic scattering experiments performed by Kendall, Friedman and Taylor revealed that protons (and subsequently neutrons) were actually composed of even smaller parti-

Standard Model of Elementary Particles




Figure 1.1: A diagram depicting the particles we currently believe are fundamental within the so-called “Standard Model” of particle physics.

cles, dubbed “partons”. This discovery was one of the largest contributing factors to the creation of the so-called Standard Model of particle physics, a theory which describes all of the fundamental particles and the way in which they interact with each other. A diagram of those fundamental particles can be seen in Figure 1.1. It should be noted that all of the particles labeled as quarks and leptons – collectively as “fermions” – have corresponding anti-particles with opposite electric charge. The equation that describes all of these particles and their interactions, often incorrectly¹ referred to as the “Standard Model Lagrangian”, can be compactified into a relatively palatable form that can easily fit on a coffee cup like the one shown in Figure 1.2.

While this equation may appear brief¹, it can be used to completely describe three

¹It is “incorrect” because this is technically a Lagrangian density (i.e. Lagrangian per unit volume), but as it is usually integrated over all space the distinction is mostly irrelevant.

¹Here “brief” is in the eye of the beholder, but certainly its brevity is misleading as even in the




Figure 1.2: A coffee cup with the Standard Model Lagrangian density printed on its side. Please ignore the “+ h.c.” term following the $i\bar{\psi}\not{D}\psi$, it is the result of a small lapse in judgement from the mug makers.

of the four fundamental forces of nature:

1. The Electromagnetic Force, which is responsible for the electrons pushing against each other to keep you from falling through your chair,
2. The Weak Nuclear Force, which is responsible for initiating the nuclear fusion reactions that fuel our sun, and
3. The Strong Nuclear Force, which is responsible for holding quarks and gluons together in bound states known as hadrons, like the protons and neutrons that make up everyday matter.

The only fundamental force missing from this list is the Gravitational Force, which is described by a completely separate set of equations²

first line the $F_{\mu\nu}$ refers to three completely different gauge field tensors...

²Specifically, the Einstein Field Equations, $G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}$, but this is the thesis of a particle physicist so gravity is taboo.

Each of the three forces that are described within the Standard Model are mediated by different gauge bosons. For example, the electromagnetic force is mediated by the boson known as the photon, the weak nuclear force is mediated by the W and Z bosons, and the strong nuclear force is mediated by bosons known as gluons. In this thesis we will be primarily focusing on the Strong Nuclear Force, which acts solely on particles with color charge – an intrinsic property of quarks and gluons. The “color” charges are red, green, and blue with antio

Even though each of the electromagnetic, weak and strong forces can be described using the Standard Model Lagrangian, the way in which they appear within the equation is not easy to determine. For example, the electromagnetic force actually corresponds to line 1

Chapter Two: Experimental Apparatus

As this thesis is focused on the physics of heavy-ion collisions, it stands to reason that the data analyzed in this thesis was gathered using the only detector along the LHC dedicated to studying such collisions: the ALICE detector. In this chapter, a brief synopsis of the LHC will be provided, followed by a much more detailed overview of the ALICE detector and its corresponding sub-detectors.

2.1 The LHC

Located along the Swiss-French border near Geneva, Switzerland, the Large Hadron Collider (LHC) is the largest particle accelerator on the planet. At a circumference of 27 kilometers, its tunnels lie almost 200 meters beneath the surface of the earth. Inside the tunnels are two high-energy particle beams pointing in opposite directions, with the beam pipes being kept inside of an ultra-high vacuum. The particles inside the beam are guided by a multitude of superconducting magnets: 393 quadrupole magnets keep the beam focused, while 1232 dipole magnets bend the particles along the circular path. The beams are designed to collide at four intersection points along the LHC, each with a corresponding detector surrounding the collision points: (1) ALICE, which specializes in heavy-ion collisions; (2) ATLAS, which specializes in studying high- p_T particles produced in pp collisions, (3) CMS, which TODO and (4) LHCb, which is designed to study CP violations through measurements of B mesons at forward rapidity. A diagram of the LHC with these four intersection points can be seen in Figure X.




Figure 2.1: A diagram depicting the LHC with its various main detectors shown underground. Illustration by Phillippe Mouche, from BBC News.

Currently, the highest center of mass energies achieved for each of the main collision systems are $\sqrt{s} = 13$ TeV for pp, $\sqrt{s} = 7$ TeV for p-Pb and $\sqrt{s} = 5.02$ TeV for Pb-Pb. The LHC underwent a long shutdown from XXXX to YYYY, in order to upgrade the beam luminosity and COM energies. The projected final COM energies for each collision system will be ...

2.2 The ALICE Detector

The detector used by the ALICE collaboration, unsurprisingly known as the ALICE detector, has the primary focus of investigating the physical properties of the strongly interacting quark-gluon plasma created during heavy-ion collisions. Building the detector was a massive effort, requiring the help from over 1000 people from 105 institutes in 30 different countries. The detector itself is also massive, weighing in




Figure 2.2: A 3-D schematic of the ALICE detector, with labels for all of the sub-detectors. Note the humans-for-scale in the bottom left of the diagram.

at around 10,000 tons and spanning 26 meters in length with a 16-meter height and width. It is composed of 18 sub-detector systems, all of which work together to help reconstruct the event. A diagram of the detector with its corresponding sub-detector systems can be seen in Figure 2.2, which contains a banana for scale. As the primary focus of the ALICE detector is to study heavy-ion collisions, all of its components must work together to reconstruct very high multiplicity events.

2.2.1 The Inner Tracking System

The Inner Tracking System (ITS) is the inner-most component of the ALICE detector, lying closest to the beam pipe. It is composed of six cylindrical layers of silicon detectors that are coaxial with the beam pipe and cover the pseudorapidity range $|\eta| \leq 0.9$. The distance from the beam line varies from 3.9 cm for the first layer to 43 cm for the sixth layer. The ITS uses different types of silicon detectors for each layer:

Silicon Pixel Detectors (SPD) for the first and second layers, Silicon Drift Detectors (SDD) for the third and fourth layers, and double sided Silicon Strip Detectors (SSD) for the fifth and sixth layers. Because of its proximity to the interaction point, the ITS is invaluable for reconstructing both primary and secondary vertices and enhancing the tracking capabilities of the ALICE detector near the interaction point. Moreover, the ITS can also track particles that are not detected or missed by the external barrel detector due to acceptance limitations and momentum cutoff.

2.2.1.1 ITS Upgrade

The LHC underwent a fairly substantial upgrade to the beam luminosity from X to Y. In order to utilize all of the The writer of this thesis was the main reason why the ITS Upgrade actually finished, as he is the smartest person of all time and actually hand built the entire thing.

2.2.2 The V0 Detector

V0 is made of two arrays of scintillator counters set on both sides of the ALICE interaction point, and called V0-A and V0-C. The V0-C counter is located upstream of the dimuon arm absorber and cover the spectrometer acceptance while the V0-A counter will be located at around 3.5 m away from the collision vertex, on the other side. It is used to estimate the centrality of the collision by summing up the energy deposited in the two disks of V0. This observable scales directly with the number of primary particles generated in the collision and therefore to the centrality. V0 is also used as reference in Van Der Meer scans that give the size and shape of colliding beams and therefore the luminosity delivered to the experiment.

2.2.3 The Time Projection Chamber

The largest component of the ALICE detector is known as the Time Projection Chamber (TPC). The TPC is a gas-filled volume with The ALICE Time Projection Chamber (TPC) is a large volume filled with a gas as detection medium and is the main particle tracking device in ALICE.[19][20] Charged particles crossing the gas of the TPC ionize the gas atoms along their path, liberating electrons that drift towards the end plates of the detector. The characteristics of the ionization process caused by fast charged particles passing through a medium can be used for particle identification. The velocity dependence of the ionization strength is connected to the well-known Bethe-Bloch formula, which describes the average energy loss of charged particles through inelastic Coulomb collisions with the atomic electrons of the medium. Multiwire proportional counters or solid-state counters are often used as detection medium, because they provide signals with pulse heights proportional to the ionization strength. An avalanche effect in the vicinity of the anode wires strung in the readout chambers, gives the necessary signal amplification. The positive ions created in the avalanche induce a positive current signal on the pad plane. The readout is performed by the 557 568 pads that form the cathode plane of the multi-wire proportional chambers (MWPC) located at the end plates. This gives the radial distance to the beam and the azimuth. The last coordinate, z along the beam direction, is given by the drift time. Since energy-loss fluctuations can be considerable, in general many pulse-height measurements are performed along the particle track in order to optimize the resolution of the ionization measurement. Almost all of the TPC's volume is sensitive to the traversing charged particles, but it features a minimum material budget. The straightforward pattern recognition (continuous tracks) make TPCs the perfect choice for high-multiplicity environments, such as in

heavy-ion collisions, where thousands of particles have to be tracked simultaneously. Inside the ALICE TPC, the ionization strength of all tracks is sampled up to 159 times, resulting in a resolution of the ionization measurement as good as 5

2.2.4 The Electromagnetic Calorimeter

The EMCal is a lead-scintillator sampling calorimeter comprising almost 13,000 individual towers that are grouped into ten super-modules. The towers are read out by wavelength-shifting optical fibers in a shashlik geometry coupled to an avalanche photodiode. The complete EMCal will contain 100,000 individual scintillator tiles and 185 kilometers of optical fiber, weighing in total about 100 tons. The EMCal covers almost the full length of the ALICE Time Projection Chamber and central detector, and a third of its azimuth placed back-to-back with the ALICE Photon Spectrometer – a smaller, highly granular lead-tungstate calorimeter. The super-modules are inserted into an independent support frame situated within the ALICE magnet, between the time-of-flight counters and the magnet coil. The support frame itself is a complex structure: it weighs 20 tons and must support five times its own weight, with a maximum deflection between being empty and being fully loaded of only a couple of centimeters. Installation of the eight-ton super-modules requires a system of rails with a sophisticated insertion device to bridge across to the support structure. The Electro-Magnetic Calorimeter (EM-Cal) will add greatly to the high momentum particle measurement capabilities of ALICE.[26] It will extend ALICE’s reach to study jets and other hard processes.

Chapter Three: Λ Production In and Out-of Jets: Analysis Overview

This chapter will provide a high-level description of the main analysis topic of this dissertation: using the Λ baryon to investigate the origins of strangeness enhancement in p-Pb collisions. First, the goal of this analysis will be clearly stated, along with the steps needed to achieve this goal. Then, each step will be described with more detail, including justification. Finally, the analysis will be summarized, and the next chapter will describe the analysis in more detail.

3.1 Objective

While the phrase “origins of strangeness enhancement” is concise, it lacks the specificity required to bring the analysis to fruition. In order to make

Chapter Four: Analysis Details

Chapter Five: Systematic Uncertainties and Cross-checks

Chapter Six: Results and Discussion

Chapter Seven: Summary and Conclusions