

# Soundness for Linear Logic regarding Phase Semantics

Ryo Haruyama

Nagoya Univ.

January 12, 2021

One kind of soundness for Linear Logic as to Phase Semantics is dealt with.

- 1 What is Linear Logic?
- 2 What is Phase Semantics?
- 3 What is soundness?
- 4 Proof of soundness
- 5 Future work

# What is Linear Logic?

Features:

- Decomposition of Classical Logic
- Realizing constructivity keeping Duality (symmetry) intact
- Resource sensitive (i.e. each hypothesis can be used exactly at once)
- Suitable for expressing parallel computing

# Decomposition of Classical Logic

Familiar connectives “ $\wedge$ ” and “ $\vee$ ” break into weaker four connectives:

- “ $\wedge$ ” into “ $\otimes$ ” and “ $\&$ ”
- “ $\vee$ ” into “ $\wp$ ” and “ $\oplus$ ”

Another viewpoint:

- *Multiplicative* are “ $\otimes$ ” and “ $\wp$ ”
- *Additive* are “ $\&$ ” and “ $\oplus$ ”

System containing only Multiplicative and Additive is called *MALL*

# Resource sensitiveness

“ $\multimap$ ” is linear version of “ $\Rightarrow$ ”

Usual logic :

$$\frac{X \quad X \Rightarrow Y}{Y \text{ (but } X \text{ still holds.)}}$$

Linear Logic :

$$\frac{X \quad X \multimap Y}{Y \text{ (and then } X \text{ disappears!)}}$$

$\otimes$  : Tensor, suitable for expressing parallel computing

$$\frac{f : A \multimap B \quad g : C \multimap D}{f \otimes g : A \otimes B \multimap C \otimes D}$$

Meaning : Programs which do NOT share the same resource can be executed at the same time.

## & : Cartesian product

$$\frac{f : X \multimap Y \quad g : X \multimap Z}{f \& g : X \multimap Y \& Z}$$

and

$$Y \& Z \not\vdash Y \otimes Z$$

Meaning : Programs which share the same resource CANNOT be executed at the same time unless copying the resource, while we CAN CHOOSE either  $Y$  or  $Z$ .

In fact, copying and deleting of resources are explicit.

$\oplus$  : Additive Sum, dual of “&”

$$Y \oplus Z \not\vdash Y$$

nor


$$Y \oplus Z \not\vdash Z$$

Meaning : Either  $Y$  or  $Z$  holds, but we CANNOT choose neither  $Y$  or  $Z$ .



# Off topic: Variants of Linear Logic

- Multiplicative are “ $\otimes$ ” and “ $\wp$ ”
- Additive are “ $\&$ ” and “ $\oplus$ ”
- *Exponential* are “ $!$ ” and “ $?$ ”



# What is Phase Semantics?

First off, what is *semantics*?

- To provide rigorous definitions that abstract away from implementation details
- To provide mathematical tools for proving properties of programs

(Amadio, Curien)

Phase Semantics is a kind of semantics, which is based on the idea of *Tarskian style*:

- “A” means A which is truth value (true or false)
- “ $A \wedge B$ ” means “A” and “B”

and so on.

This seems obvious, however, there is another semantics which is not the case: *Coherent Semantics*, which is BHK style inconsistent semantics.

*Phase Space* is Phase Semantics for MALL.

# Other semantics

- Coherent semantics
- Categorical semantics
- Geometry of interaction
- Game semantics  
and so on.

# What is *soundness*?

Formulae derived using specific rules are semantically valid, which is minimum requirement for semantics in general (Systems which yields lie are completely useless).

# Syntax of MALL

## Definition 1

Formula of MALL.

$$\begin{aligned}
 A ::= & p \mid p^\perp \\
 & \mid A \otimes A \mid A \oplus A \\
 & \mid A \& A \mid A \wp A \\
 & \mid \mathbf{1} \mid \mathbf{0} \mid \top \mid \perp
 \end{aligned}$$

# Inference rules of MALL I

Inference rules of MALL.

$$\begin{array}{c}
 \frac{}{\vdash A, A^\perp} \text{ (identity)} \qquad \frac{\vdash \Gamma, A \quad \vdash A^\perp, \Delta}{\vdash \Gamma, \Delta} \text{ (cut)} \\
 \\
 \frac{\vdash \Gamma, A, B, \Delta}{\vdash \Gamma, B, A, \Delta} \text{ (exchange)} \\
 \\
 \frac{}{\vdash \mathbf{1}} \text{ (one)} \qquad \frac{\vdash \Gamma}{\vdash \Gamma, \perp} \text{ (false)}
 \end{array}$$

## Inference rules of MALL II

$$\frac{\vdash \Gamma, A \quad \vdash \Delta, B}{\vdash \Gamma, A \otimes B, \Delta} \text{ (times)} \qquad \frac{\vdash \Gamma, A, B}{\vdash \Gamma, A \wp B} \text{ (par)}$$

$$\frac{}{\vdash \Gamma, \top} \text{ (true)} \qquad \text{ (no rule for zero)}$$

$$\frac{\vdash \Gamma, A \quad \vdash \Gamma, B}{\vdash \Gamma, A \& B} \text{ (with)} \qquad \frac{\vdash \Gamma, A}{\vdash \Gamma, A \oplus B} \text{ (left plus)}$$

$$\frac{\vdash \Gamma, B}{\vdash \Gamma, A \oplus B} \text{ (right plus)}$$

# Phase Space

We give semantics for MALL.

## Definition 2

Phase Space  $:= (M, \perp)$

where  $M$  is commutative monoid and  $\perp \subseteq M$  is defined.

## Definition 3

Commutative monoid  $M$  holds

- commutativity:  $pq = qp$
- associativity:  $(pq)r = p(qr)$
- identity:  $1p = p1 = p$

for all  $p, q, r \in M$ .



## Definition 4

$X \multimap Y$  is defined as

$$\begin{aligned} m \in X \multimap Y \\ :\Leftrightarrow \forall x (x \in X \Rightarrow mx \in Y) \end{aligned}$$

## Definition 5

orthogonal

$$X^\perp := X \multimap \perp$$

## Definition 6

$X$  is *fact* iff

$$X = X^{\perp\perp}$$

or equivalently,  $X$  is of the form  $Y^\perp$ .

## Definition 7

For convention,

$$xy \in X.Y :\Leftrightarrow x \in X \wedge y \in Y$$

Let  $X$  and  $Y$  be fact. Connectives are interpreted in this way (more precisely, we are defining interpretation function from *formula* to  $\wp(M)$ ):

$$X \otimes Y := (X.Y)^{\perp\perp}$$

$$X \wp Y := (X^{\perp}.Y^{\perp})^{\perp}$$

$$X \multimap Y = (X.Y^{\perp})^{\perp}$$

$$X \& Y := X \cap Y$$

$$X \oplus Y := (X \cup Y)^{\perp\perp}$$

$$\mathbf{1} := \{1\}^{\perp\perp}$$

$$\mathbf{0} := \emptyset^{\perp\perp}$$

$$\top := M$$

# Validity

## Definition 8

Sequent

$$\vdash \Gamma, A$$

is interpreted as subset of  $M$

$$\Gamma \vDash A$$

## Definition 9

$\underline{X}$  (as formula) is valid iff  $1 \in \underline{X}$  ( $\underline{X} \subseteq M$ )

# Proof of soundness I

## Theorem 10

*Sequent which are provable in MALL are all valid in Phase Space*  
*i. e.*

$$\vdash \underline{X} \Rightarrow 1 \in \underline{X}$$

where  $\underline{X}$  are all fact.

By straightforward induction on inference rules.

$$\textcircled{1} \frac{}{\vdash A, A^\perp} \text{ (identity)}$$

$$A \wp A^\perp = A \multimap A \ni 1$$

( $\because$  definition of identity 1)

# Proof of soundness II

$$\textcircled{2} \quad \overline{\vdash \Gamma, \top} \text{ (true)}$$

$$\Gamma \not\approx \top = \mathbf{0} \multimap \Gamma$$

Since  $\mathbf{0}$  is the smallest fact,  $\mathbf{0} \subseteq \Gamma$ . This implies  $\forall z, z \in \mathbf{0} \Rightarrow z \in \Gamma$ .

Hence  $\forall z, z \in \mathbf{0} \Rightarrow 1z \in \Gamma$ .

$\therefore$  By definition of “ $\multimap$ ”,  $1 \in \mathbf{0} \multimap \Gamma = \Gamma \not\approx \top$

$$\textcircled{3} \quad \overline{\vdash \mathbf{1}} \text{ (one)}$$

Obviously,

$$1 \in \{1\} \subseteq \{1\}^{\perp\perp} = \mathbf{1}$$

# Proof of soundness III

$$④ \frac{\vdash \Gamma, A \quad \vdash A^\perp, \Delta}{\vdash \Gamma, \Delta} \text{ (cut)}$$

We have to show that if  $1 \in \Gamma \wp A$  and  $1 \in A^\perp \wp \Delta$  then  $1 \in \Gamma \wp \Delta$ .  
In fact this is equivalent to

$$\begin{aligned} 1 \in \Gamma^\perp \multimap A, 1 \in A \multimap \Delta \\ \Rightarrow 1 \in \Gamma^\perp \multimap \Delta \end{aligned}$$

This is easily followed by

$$\begin{aligned} \Gamma^\perp \subseteq A, A \subseteq \Delta \\ \Rightarrow \Gamma^\perp \subseteq \Delta \end{aligned}$$

$$⑤ \frac{\vdash \Gamma, A, B, \Delta}{\vdash \Gamma, B, A, \Delta} \text{ (exchange)}$$

$\therefore$  “ $\wedge$ ” is commutative.

# Proof of soundness IV

$$\textcircled{6} \quad \frac{\vdash \Gamma, A \quad \vdash \Gamma, B}{\vdash \Gamma, A \& B} \text{ (with)}$$

We have to show that if  $1 \in \Gamma \mathcal{J} A$  and  $1 \in \Gamma \mathcal{J} B$  then  $1 \in \Gamma \mathcal{J} (A \& B)$ . Here we use distributivity of  $\mathcal{J}$  over  $\&$ :

$$(\Gamma \mathcal{J} A) \& (\Gamma \mathcal{J} B) = \Gamma \mathcal{J} (A \& B)$$

By definition of  $\&$  and hypothesis, left hand side contains 1, and so does right hand side.

$$\textcircled{7} \quad \frac{\vdash \Gamma, A}{\vdash \Gamma, A \oplus B} \text{ (left plus)} \quad \frac{\vdash \Gamma, B}{\vdash \Gamma, A \oplus B} \text{ (right plus)}$$

Similarly, we use half-distributivity of  $\mathcal{J}$  over  $\oplus$ :

$$(\Gamma \mathcal{J} A) \oplus (\Gamma \mathcal{J} B) \subseteq \Gamma \mathcal{J} (A \oplus B)$$

# Proof of soundness $\forall$

$$\textcircled{8} \frac{\vdash \Gamma}{\vdash \Gamma, \perp} (\text{false})$$

$$\Gamma \wp \perp = \mathbf{1} \multimap \Gamma$$

By definition,  $\mathbf{1}$  is the smallest fact contains  $1$  i. e.  $\mathbf{1} \subseteq \Gamma$  so that  $\forall e, e \in \mathbf{1} \Rightarrow e = 1e \in \Gamma$  hence  $1 \in \mathbf{1} \multimap \Gamma$ .

$$\textcircled{9} \frac{\vdash \Gamma, A \quad \vdash \Delta, B}{\vdash \Gamma, A \otimes B, \Delta} (\text{times})$$

We have to show that if  $1 \in \Gamma \wp A$  and  $1 \in \Delta \wp B$  then  $1 \in \Gamma \wp \Delta \wp (A \otimes B)$ . Hypotheses can be transformed into



# Proof of soundness VI

$1 \in \Gamma^\perp \multimap A$  and  $1 \in \Delta^\perp \multimap B$  respectively. Therefore, we have  
 $\Gamma^\perp \subseteq A$  and  $\Delta^\perp \subseteq B$

$$\Rightarrow \Gamma^\perp . \Delta^\perp \subseteq A.B$$

$$\Rightarrow (\Gamma^\perp . \Delta^\perp)^{\perp\perp} \subseteq (A.B)^{\perp\perp}$$

$$\Rightarrow \Gamma^\perp \otimes \Delta^\perp \subseteq A \otimes B$$

$$\Rightarrow \Gamma^\perp \otimes \Delta^\perp \multimap A \otimes B \ni 1$$

$$\Rightarrow \Gamma \wp \Delta \wp (A \otimes B) \ni 1$$

$$\textcircled{10} \frac{\vdash \Gamma, A, B}{\vdash \Gamma, A \wp B} (par)$$

This is tautology. □

# Future work

- Categorical semantics: symmetrical monoidal (closed) category
- Application to functional programming: Combinatorial linear logic, categorical and linear machine
- Game semantics