

## ORIGINAL ARTICLE

# Antiphospholipid syndrome: antibodies to Domain 1 of $\beta$ 2-glycoprotein 1 correctly classify patients at risk

V. PENGO,\* A. RUFFATTI,† M. TONELLO,† S. CUFFARO,† A. BANZATO,\* E. BISON,\* G. DENAS\* and S. PADAYATTIL JOSE\*

\*Clinical Cardiology, Thrombosis Centre, Department of Cardiac Thoracic and Vascular Sciences, University of Padua; and †Rheumatology Unit, Department of Medicine, University of Padua, Padua, Italy

**To cite this article:** V. Pengo, A. Ruffatti, M. Tonello, S. Cuffaro, A. Banzato, E. Bison, G. Denas, S. Padayattil Jose. Antiphospholipid syndrome: Antibodies to Domain 1 of  $\beta$ 2-glycoprotein 1 correctly classify patients at risk. *J Thromb Haemost* 2015; **13**: 782–7.

**Summary.** *Background:* Determination of lupus anticoagulant (LA), anticardiolipin (aCL) and  $\beta$ 2-Glycoprotein 1 (a $\beta$ 2GP1) antibodies is mandatory to classify patients with antiphospholipid syndrome (APS) into risk categories. *Objectives:* To measure relevant antibodies, considered to be those of the IgG isotype directed towards  $\beta$ 2GP1 and particularly those directed to Domain 1 (Dm1) of the molecule. *Patients/methods:* In this cross-sectional study we measured IgG a $\beta$ 2GP1-Dm1 by a chemiluminescent immunoassay in a group of individuals initially positive for IgG a $\beta$ 2GP1 and classified as triple (LAC+, IgG aCL+, IgG a $\beta$ 2GP1+,  $n = 32$ ), double (LAC–, IgG aCL+, IgG a $\beta$ 2GP1+,  $n = 23$ ) or single positive (LA–, IgG aCL–, IgG a $\beta$ 2GP1+,  $n = 10$ ). *Results and conclusion:* Geometric mean and standard deviation expressed as chemiluminescent units (CU) in triple, double and single positive groups were  $273.0 \pm 6.2$ ,  $18.2 \pm 9.6$  and  $4.4 \pm 2.2$ , respectively. The geometric mean obtained in 40 healthy subjects was  $2.0 \pm 2.0$ . Mean CU values were significantly different among groups and with respect to values found in 40 healthy subjects ( $P < 0.0001$ ). Positive values of IgG a $\beta$ 2GP1-Dm1 (above 14.2 CU) were found in 45 individuals while 20 individuals ( $20/65 = 30.8\%$ ) positive for IgG a $\beta$ 2GP1 were negative for IgG a $\beta$ 2GP1-Dm1. There was a significant association between positive IgG a $\beta$ 2GP1-Dm1 and thromboembolic events ( $P = 0.001$ ). Positive and negative values of IgG a $\beta$ 2GP1-Dm1 were consistently confirmed after 12 weeks, with only three low positive values being negative after 12 weeks. In conclusion, IgG a $\beta$ 2GP1-Dm1 seems a robust and reproducible test that in association with the classic tests may be useful in clinical practice

in identifying individuals at high risk of developing thromboembolic events.

**Keywords:** antibodies; antiphospholipid syndrome; epitopes; glycoproteins; thrombosis.

## Introduction

Antiphospholipid syndrome (APS) is characterized by the presence of antiphospholipid (aPL) antibodies in patients with thrombosis or pregnancy morbidity [1]. To categorize APS patients, three tests exploring the presence of aPL antibodies (lupus anticoagulant (LA), anticardiolipin (aCL) and anti- $\beta$ 2-Glycoprotein 1 (a $\beta$ 2GP1) antibodies) are performed. In this way, APS patients are classified into category I when more than one test is positive, and IIa, IIb and IIc according to the presence of only one positive test [1]. When all three tests are positive (triple positivity), the association with thrombosis is very strong; it gets weaker in double-positive (with LA negative) and single-positive patients [2]. Antibodies determining positivity in all the three tests are those directed against  $\beta$ 2GP1, a plasma protein without a defined function [3, 4]. Indeed, affinity-purified IgG a $\beta$ 2GP1 spiked into normal plasma made all three tests positive [3]. Moreover, a subgroup of IgG a $\beta$ 2GP1, those directed to Domain1 (Dm1) in the NH2 terminal region of the molecule, are able to interfere with the coagulation process and are associated with thrombosis [5]. The aim of this cross-sectional study was to measure both basal and after 12 weeks values of IgG a $\beta$ 2GP1-Dm1 in individuals initially positive for IgG a $\beta$ 2GP1.

## Material and methods

### Study population

Consecutive individuals referred to a rheumatology outpatient clinic dedicated to APS patients and initially

Correspondence: Vittorio Pengo, Clinical Cardiology, Thrombosis Centre, University of Padua, Via Giustiniani 2, 35128 Padova, Italy. Tel./fax: +39 49 8215658. E-mail: vittorio.pengo@unipd.it

Received 15 October 2014

Manuscript handled by: M. Levi

Final decision: F. R. Rosendaal, 24 January 2015

positive for aPL antibodies were evaluated. The selection criteria for individuals were not based on the diagnosis of the patients, but rather on the presence of APS-related symptoms (thrombosis and pregnancy morbidity) and aPL antibody positivity. Because IgG a $\beta$ 2GPI-Dm1 is a subgroup of IgG directed to  $\beta$ 2GPI, the study focused on the group of individuals initially positive for IgG a $\beta$ 2GPI. They were classified into three groups: triple positive (LA+, IgG aCL+, IgG a $\beta$ 2GPI+), double positive (LA-, IgG aCL+, IgG a $\beta$ 2GPI+) and single positive (LA-, IgG aCL-, IgG a $\beta$ 2GPI+). Two of us (VP and AR) consulted the available hospital records and diagnosed previous thromboembolic events only in the presence of objective criteria and if confirmed by positive compression ultrasonography or venography for deep vein thrombosis, and spiral tomography, ventilation-perfusion lung scan or pulmonary angiography for pulmonary embolism. Positive tomographic scanning, magnetic resonance imaging or angiography confirmed the diagnosis of arterial thromboembolism. The presence of pregnancy morbidity according to Miyakis [1] was assessed by individual interview. Hospital records were also consulted for the presence of associated autoimmune diseases (systemic lupus erythematosus [SLE] or other connective tissue disease). No subject was lost to follow-up after 12 weeks. We restricted this study to individuals positive for IgG a $\beta$ 2GPI but we have also tested IgG a $\beta$ 2GPI-Dm1 in individuals negative for IgG a $\beta$ 2GPI but positive for one or more tests exploring the presence of aPL.

#### *Determination of aPL antibodies*

Venous blood was collected in 3.8% sodium citrate (9 : 1) and centrifuged twice at 2000  $\times g$  for 15 min at 4 °C and obtained plasma was stored at -80 °C until used. Patients' plasmas were screened for the presence of aPL antibodies using home-made IgG and IgM aCL ELISA as previously described [6] and the recommendations proposed by a taskforce during the 13th International Congress on Antiphospholipid Antibodies (Galveston, TX, USA) [7]. LA was detected using dRVVT and SCT (Instrumentation Laboratory, Milan, Italy), according to the ISTH guidelines [8]. The presence of IgG a $\beta$ 2GPI antibodies was confirmed by a chemiluminescent immunoassay (QUANTA flash  $\beta$ 2GPI IgG; Inova Diagnostics, San Diego, CA, USA). The cut-off value of tests exploring the presence of aPL antibodies was calculated by means of the 99th percentile of results obtained in 40 healthy subjects. The cut-off value for both dRVVT and SCT in mixing studies was 1.2; the cut-off values of IgG and IgM aCL and IgG and IgM a $\beta$ 2GPI were 10 GPL and 8 MPL and 13 and 7 home units, respectively. The IgG a $\beta$ 2GPI cut-off value in the chemiluminescent immunoassay was 19.4 CU.

#### *Determination of IgG a $\beta$ 2GPI-Dm1 antibodies*

Autoantibodies to  $\beta$ 2GPI-Dm1 were measured by a chemiluminescent immunoassay (QUANTA Flash Domain 1 IgG; Inova Diagnostics) using the BIO-FLASH® technology (Inova Diagnostics). The plasma sample was diluted 1 : 10 by the instrument in a disposable plastic cuvette. The technician performing the assay was blinded to patient/individual classification categories. Small amounts of the diluted patient plasma and the assay buffer were all combined into a second cuvette containing purified recombinant  $\beta$ 2GPI-Dm1 coated onto paramagnetic beads, mixed and incubated at 37 °C. Isoluminol-conjugated anti-human IgG antibody was then added to the cuvette and incubated at 37 °C. The light coming from the luminescent reaction produced by isoluminol conjugate was measured as Relative Light Units (RLU) by the BIO-FLASH® optical system. The RLU are proportional to the amount of bound isoluminol conjugate, which in turn is proportional to the amount of a $\beta$ 2GPI-Dm1 antibodies bound to the  $\beta$ 2GPI-Dm1 on the beads. Relative light units are converted to chemiluminescent units (CU) when the assay is assigned to a lot-specific master curve, and this curve is calibrated with QUANTA Flash calibrators to obtain the final CU values for results. The approximate cut-off value for IgG a $\beta$ 2GPI-Dm1, calculated by means of the 99th percentile of results obtained in 40 healthy subjects, was 14.2 CU. Intra-assay and inter-assay coefficient of variation are 6.3% and 2.0%, respectively.

#### *Statistics*

Descriptive statistics are reported as appropriate: categorical data are expressed as frequencies (percentage); continuous data are reported as mean (standard deviation). Due to non-Gaussian distribution, data of IgG a $\beta$ 2GPI-Dm1 (CU) were transformed into log10. In this way the distribution became Gaussian and the difference between groups was tested using one-way analysis of variance (ANOVA). The chi-squared test for independence was used to evaluate the association of thrombosis and pregnancy loss with positive values of IgG a $\beta$ 2GPI-Dm1. Linear (Pearson) correlation was performed between IgG a $\beta$ 2GPI-Dm1 (CU) initial data and data obtained after 12 weeks. Spearman correlation was used to compare Domain1 and the whole molecule titres. GRAPH-PAD INSTAT 3 (GraphPad Software, San Diego, CA, USA) was used for the statistical analysis (www.graphpad.com).

#### *Results*

During the period 2009–2013, 1120 individuals were tested for the presence of aPL antibodies (LAC, aCL IgG and IgM, a $\beta$ 2GPI IgG and IgM). Reasons for testing

were vascular thrombosis, previous pregnancy morbidity, SLE or other autoimmune diseases, familial aPL, oral contraceptive or hormone replacement therapy, accidental finding of prolonged activated partial thromboplastin time or false-positive test results for syphilis. The final study cohort was selected based on aPL antibody positivity and anti  $\beta$ 2GPI antibody positivity. Among the 106 individuals positive for one or more test, 65 were positive for the presence of IgG a $\beta$ 2GPI. These individuals were grouped according to their aPL profile into triple positive (LAC+, IgG aCL+, IgG a $\beta$ 2GPI+,  $n = 32$ ), double positive (LAC-, IgG aCL+, IgG a $\beta$ 2GPI+,  $n = 23$ ) and single positive (LAC-, IgG aCL-, IgG a $\beta$ 2GPI+,  $n = 10$ ). DRVVT and SCT were both positive in triple-positive patients/carriers, and both were negative in the other two groups. Table 1 shows individuals' characteristics. Of note, only young females represented the single-positive group, while autoimmune disease was lower in the triple-positive group. The rate of clinical events related to APS (thromboembolism and pregnancy loss) was also different among groups; the rate of thromboembolic events was higher in triple-positive individuals, while the rate of pregnancy loss was more frequent in double-positive individuals. Figure 1 shows the values expressed as chemiluminescent units (CU) of IgG a $\beta$ 2GPI-Dm1 in the three groups. Geometric mean and standard deviation expressed as CU in triple, double and single-positive groups were  $273.0 \pm 6.2$ ,  $18.2 \pm 9.6$  and  $4.4 \pm 2.2$ , respectively. The geometric mean obtained in 40 healthy subjects was  $2.0 \pm 2.0$ . One-way analysis of variance (ANOVA) showed a statistically significant variation among columns in the mean values of IgG a $\beta$ 2GPI-Dm1 antibodies ( $P < 0.0001$ ).


The approximate cut-off value set at the 99th percentile obtained in 40 control subjects negative for the presence of IgG a $\beta$ 2GPI was 14.2 CU. Positive values of IgG a $\beta$ 2GPI-Dm1 (above 14.2 CU) were found in 31 out of 32 triple-positive (97%), 13 out of 23 double-positive (43%) and in 1 out of 10 single-positive (10%) individuals. When comparing the mean values of IgG a $\beta$ 2GPI-Dm1 between the groups, we found a significantly higher value in triple-positive individuals ( $P < 0.0001$ ).

There is a significant correlation between the two tests (Spearman  $r = 0.76$ ; 95% CI, 0.6–0.85;  $P < 0.0001$ ) (Fig. 2). However, there were 20 individuals out of 65

(20/65 = 30.8%) positive for IgG a $\beta$ 2GPI and negative for IgG a $\beta$ 2GPI-Dm1, all but one of them in the double and single-positive groups.

Thromboembolic events were present in 29 of 45 individuals (64%) with positive values of IgG a $\beta$ 2GPI-Dm1 (above 14.2 CU) and in four of 20 (20%) individuals with negative values (Fig. 3).


Overall, there was a very significant association between positive IgG a $\beta$ 2GPI-Dm1 and thromboembolic events ( $P = 0.001$ ), with a sensitivity of 88% and a specificity of 50%. On the other hand, pregnancy loss was present in 16 out of 39 women (41%) with positive values of IgG a $\beta$ 2GPI-Dm1 and in three out of 19 women (16%) with negative values ( $P = 0.07$ ). All the individuals were tested after 12 weeks. As shown in Fig. 4, there was a consistent confirmation after 12 weeks of data obtained at initial screening (correlation coefficient [ $r$ ] was 0.99; 95% CI, 0.98–0.99 [ $P < 0.0001$ ]). Overall, of the 45 positive values at initial testing, only three low positive values of IgG a $\beta$ 2GPI-Dm1 (two in double-positive and one in single-positive groups) were negative after 12 weeks. On the other hand, of 20 negative values at initial testing not one was positive after 12 weeks. The correlation coefficient between basal and 12-week values of 45 initially




**Fig. 1.** Variations of geometric mean of IgG a $\beta$ 2GPI-Dm1 values expressed as chemiluminescent units (CU) among groups are significantly greater than expected by chance ( $P < 0.0001$ ). All the individuals in the three groups are positive for IgG a $\beta$ 2GPI.

**Table 1** Characteristics of individuals positive for IgG a $\beta$ 2-glycoprotein 1 according to the aPL antibody profile

	LA+, IgG aCL+, IgG a $\beta$ 2GPI+ $n = 32$	LA-, IgG aCL+, IgG a $\beta$ 2GPI+ $n = 23$	LA-, IgG aCL-, IgG a $\beta$ 2GPI+ $n = 10$
Age, years (mean $\pm$ SD)	41 $\pm$ 13	45 $\pm$ 12	32 $\pm$ 17
Male, no. (%)	6 (19)	2 (9)	0 (0)
Autoimmune disorders, no. (%)	2 (6)	7 (30)	3 (30)
Venous or arterial thrombosis, no. (%)	24 (75)	8 (35)	1 (10)
Pregnancy loss, no. (%)	7 (22)	10 (43)	0 (0)
Carriers, no. (%)	5 (16)	8 (35)	9 (90)



**Fig. 2.** Correlation between IgG aβ2GPI and IgG aβ2GPI-Dm1 ( $P < 0.0001$ ). Dots inside the rectangle denote individuals positive for IgG aβ2GPI and negative for IgG aβ2GPI-Dm1.




**Fig. 3.** Rate of thromboembolic events in individuals with values positive or negative for IgG anti-β2GPI-Domain1.

positive IgG aβ2GPI-Dm1 values was extremely significant (Pearson  $r = 0.94$ ,  $P < 0.0001$ ), as was that of 20 initially negative IgG aβ2GPI-Dm1 values (Pearson  $r = 0.89$ ,  $P < 0.0001$ ).


Of the 106 initially positive individuals, 41 were negative for IgG aβ2GPI. In this group, 16 were triple positive ( $n = 2$ ) and double positive ( $n = 14$ ) for IgM aβ2GPI and 25 were positive for only one test (8 IgM aβ2GPI, 8 IgG aCL, 7 IgM aCL, 2 LA). As shown in Fig. 5, no one was positive for IgG anti Dm1.

## Discussion

In this study, we assessed individuals testing positive at initial screening for IgG aβ2GPI alone or in combination with the other two tests required to classify APS patients. Of these, 69% were positive for antibodies directed against β2GPI-Dm1, a rate somewhat higher (55%) than previously reported [9]. Our results show that IgG aβ2GPI-Dm1 values are significantly higher in individuals with a triple-positive aPL profile, and that there is an association between the concentration of these antibodies



**Fig. 4.** Linear correlation between data of IgG anti-β2GPI-Domain1 (CU log10) obtained at initial testing and after 12 weeks.



**Fig. 5.** Values of IgG aβ2GPI-Dm1 expressed as relative light units (CU) in triple IgM isotype positive subjects ( $n = 2$ ), double IgM isotype positive subjects ( $n = 14$ ) and 25 individuals positive for only one test.

and thrombosis risk categories. All the triple-positive individuals, except one, had values well above the approximate cut-off calculated in 40 healthy subjects. Thus, both high titre IgG aβ2GPI-Dm1 and triple positivity identify patients with previous thromboembolic events at high risk of recurrence [10] and carriers at high risk of developing a first thromboembolic event [11]. At variance with other risk categories, triple-positive individuals have positive LA and this is a peculiar feature of the subgroup of IgG aβ2GPI antibodies that are directed against Dm1 [5,12,13]. However, IgG aβ2GPI-Dm1 are also positive in some individuals without LA: one explanation might be that in this case, a low titre of antibodies is present, such that it is insufficient to determine LA activity; another reason might be that the detected subset of IgG aβ2GPI-Dm1 is not directed against the Gly40-Arg43 epitope. The main epitope of Dm1 possesses in fact fine specificity, and it is discontinuous in nature, involving arginine 39-arginine 43, aspartic



acid 8-aspartic acid 9, and possibly the interlinking region between DI and DII [14–16]. Conversely, despite positivity in IgG a $\beta$ 2GPI, many individuals in double and all but one in single-positive groups had negative values of IgG a $\beta$ 2GPI-Dm1. In these individuals, IgG a $\beta$ 2GPI may be directed to other Domains of the molecule. For example, anti-Domain 4/5 antibodies have been detected in non-thrombotic conditions, such as atherosclerosis [17], leprosy [18] and in children with atopic dermatitis [19] or those born to mothers with systemic autoimmune diseases [20]. The presence of antibodies preferentially directed against Domain 4/5 is not associated with thromboembolic events [20]. Interestingly enough, lower levels of IgG a $\beta$ 2GPI-Dm1, as in the double-positive group, might be relevant in pregnancy morbidity. Indeed, we have previously shown when analyzing data from retrospective studies, that low positive aCL and double positivity are more strongly associated with pregnancy morbidity than with thromboembolic events [2,21].

According to present guidelines, confirmation of aPL positivity is mandatory to exclude the presence of transient antibodies [1]. This is particularly true for individuals with single test positivity as confirmation is poor [22]. Results from the present study show that IgG a $\beta$ 2GPI-Dm1 are not transient and confirmation values are very close to those initially detected. Positivity was confirmed after 12 weeks with a very high rate of concordance. As pointed out in a recent comprehensive review paper [23], we feel that this chemiluminescent immunoassay using the BIO-FLASH<sup>®</sup> technology is more sensitive and reproducible than the existing ELISA assays. The package insert for this CE marked assay contains ample testing of QUANTA Flash Domain 1 on a variety of disease controls, including IBD, infectious osteoarthritis, RA and systemic sclerosis, which shows a very high specificity (> 99%). However, prospective studies need to make clear their clinical utility. In conclusion, in addition to the other aPL tests, the finding of high levels of IgG a $\beta$ 2GPI-Dm1 antibodies identifies individuals at high risk of thrombosis, and thus might become an additional useful tool in clinical and therapeutic decision-making in aPL-positive individuals and APS patients.

## Addendum

V. Pengo and A. Ruffatti designed the research, analyzed and interpreted data and wrote the manuscript; M. Tonello and S. Cuffaro, collected samples and data; E. Bison, A. Banzato, G. Denas and S. P. Jose performed research and G. Denas performed statistical analysis.

## Disclosure of Conflict of Interests

The authors state that they have no conflict of interests.

## References

- 1 Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, Derksen RHWM, de Groot PG, Koike T, Meroni PL, Reber G, Shoenfeld Y, Tincani A, Vlachoyiannopoulos PG, Kritis SA. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006; **4**: 295–306.
- 2 Pengo V, Biasiolo A, Pegoraro C, Cucchini U, Noventa F, Iliceto S. Antibody profiles for the diagnosis of antiphospholipid syndrome. *Thromb Haemost* 2005; **93**: 147–52.
- 3 Pengo V, Biasiolo A, Rampazzo P, Brocco T. dRVVT is more sensitive than KCT or TTI for detecting Lupus Anticoagulant activity of anti-b2-Glycoprotein-I autoantibodies. *Thromb Haemost* 1999; **81**: 256–8.
- 4 Banzato A, Pozzi N, Frasson R, De Filippis V, Ruffatti A, Bison E, Padayattil Jose S, Denas G, Pengo V. Antibodies to domain I of  $\beta$ 2Glycoprotein I are in close relation to patients risk categories in Antiphospholipid Syndrome (APS). *Thromb Res* 2011; **128**: 583–6.
- 5 de Laat B, Derksen RH, Urbanus RT, de Groot PG. IgG antibodies that recognize epitope Gly40-Arg43 in domain I of beta 2-glycoprotein I cause LAC, and their presence correlates strongly with thrombosis. *Blood* 2005; **105**: 1540–5.
- 6 Pengo V, Biasiolo A, Fior MG. Autoimmune antiphospholipid antibodies are directed against a cryptic epitope expressed when b2-glycoprotein-I is bound to a suitable surface. *Thromb Haemost* 1995; **73**: 29–34.
- 7 Pierangeli SS, de Groot PG, Dlott J, Favaloro R, Harris EN, Lakos G, Ortel T, Meroni PL, Otomo K, Pengo V, Tincani A, Wong R, Roubey R. 'Criteria' aPL tests: report of a task force and preconference workshop at the 13th International Congress on Antiphospholipid Antibodies, Galveston, Texas, April 2010. *Lupus* 2011; **20**: 182–90.
- 8 Pengo V, Tripodi A, Reber G, Rand JH, Ortel TL, Galli M, de Groot PG; Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the International Society on Thrombosis and Haemostasis. Update of the guidelines for lupus anticoagulant detection. Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the International Society on Thrombosis and Haemostasis. *J Thromb Haemost* 2009; **7**: 1737–40.
- 9 de Laat B, Pengo V, Pabinger I, Musial J, Voskuyl AE, Bultink IE, Ruffatti A, Rozman B, Kveder T, de Moerloose P, Boehlen F, Rand J, Ulcova-Gallova Z, Mertens K, de Groot PG. The association between circulating antibodies against domain I of beta2-glycoprotein I and thrombosis: an international multicenter study. *J Thromb Haemost* 2009; **7**: 1767–73.
- 10 Pengo V, Ruffatti A, Legnani C, Gresele P, Barcellona D, Erba N, Testa S, Marongiu F, Bison E, Denas G, Banzato A, Padayattil Jose S, Iliceto S. Clinical course of high-risk patients diagnosed with antiphospholipid syndrome. *J Thromb Haemost* 2010; **8**: 237–42.
- 11 Pengo V, Ruffatti A, Legnani C, Testa S, Fierro T, Marongiu F, De Micheli V, Gresele P, Tonello M, Ghirarduzzi A, Bison E, Denas G, Banzato A, Padayattil Jose S, Iliceto S. Incidence of a first thromboembolic event in asymptomatic carriers of high risk antiphospholipid antibody profile: a multicenter prospective study. *Blood* 2011; **118**: 4714–8.
- 12 de Laat HB, Derksen RH, Urbanus RT, Roest M, de Groot PG. Beta2-glycoproteinI-dependent lupus anticoagulant highly correlates with thrombosis in the antiphospholipid syndrome. *Blood* 2004; **104**: 3598–602.
- 13 Zuily S, De Laat B, Regnault V, Wahl D. Autoantibodies against domain I of beta2-glycoprotein indicate an increased risk

- for thrombosis in antiphospholipid patients. A prospective cohort study. *Arthritis Rheum* 2013; **65**: S1130.
- 14 Reddel SW, Wang YX, Sheng YH, Krillis SA. Epitope studies with anti- $\beta$ 2-glycoprotein I antibodies from autoantibody and immunized sources. *J Autoimmun* 2000; **15**: 91–6.
  - 15 Iverson GM, Reddel S, Victoris EJ, Cockerill KA, Wang YX, Marti-Renom MA, Sali A, Marquis DM, Krillis SA, Linnik MD. Use of single point mutations in domain I of  $\beta$ 2-glycoprotein I to determine fine antibody specificity of antiphospholipid autoantibodies. *J Immunol* 2002; **169**: 7097–103.
  - 16 Ioannou Y, Pericleous C, Giles I, Latchman DS, Isenberg DA, Rahman A. Binding of antiphospholipid antibodies to discontinuous epitopes on domain I of human  $\beta$ 2-glycoprotein I: mutation studies including residues R39 to R43. *Arthritis Rheum* 2007; **56**: 280–90.
  - 17 Iverson GM, von Muhlen CA, Staub HL, Lassen AJ, Binder W, Norman GL. Patients with atherosclerotic syndrome, negative in anti-cardiolipin assays, make IgA autoantibodies that preferentially target domain4 of  $\beta$ 2-GPI. *J Autoimmun* 2006; **27**: 266–71.
  - 18 Arvieux J, Renaudineau Y, Mane I, Perraut R, Krillis SA, Youinou P. Distinguishing features of anti- $\beta$ 2glycoprotein I antibodies between patients with leprosy and the antiphospholipid syndrome. *Thromb Haemost* 2002; **87**: 599–605.
  - 19 Ambrozic A, Avicin T, Ichikawa K, Kveder T, Matsuura E, Hojnik M, Atsumi T, Rozman B, Koike T. Antibeta(2)-glycoprotein I antibodies in children with atopic dermatitis. *Int Immunol* 2002; **14**: 823–30.
  - 20 Andreoli L, Nalli C, Motta M, Norman GL, Shums Z, Encabo S, Binder WL, Nuzzo M, Frassi M, Lojcono A, Avicin T, Meroni PL, Tincani A. Anti- $\beta$ 2glycoprotein I IgG antibodies from 1-year-old healthy children born to mothers with systemic autoimmune diseases preferentially target domain4/5: might it be the reason for their innocent' profile? *Ann Rheum Dis* 2011; **70**: 380–3.
  - 21 Ruffatti A, Olivieri S, Tonello M, Bortolati M, Bison E, Salvan E, Facchinetti M, Pengo V. Influence of different IgG anticardiolipin antibody cut-off values on antiphospholipid syndrome classification. *J Thromb Haemost* 2008; **6**: 1693–6.
  - 22 Pengo V, Ruffatti A, Del Ross T, Tonello M, Cuffaro S, Hoxha A, Banzato A, Bison E, Denas G, Bracco A, Padayattil Jose S. Confirmation of initial antiphospholipid antibody positivity depends on the antiphospholipid antibody profile. *J Thromb Haemost* 2013; **11**: 1–5.
  - 23 Bertolaccini ML, Amengual O, Andreoli L, Atsumi T, Chighizola CB, Forastiero R, de Groot P, Lakos G, Lambert M, Meroni PL, Ortel TL, Petri M, Rahman A, Roubey R, Sciascia S, Snyder M, Tebo AE, Tincani A, Willis R. 14th International Congress on Antiphospholipid Antibodies Task Force. Report on antiphospholipid syndrome laboratory diagnostics and trends. *Autoimmun Rev* 2014; **13**: 917–30.