

Checkpoint: Traces and level curves

Consider the functions $C(x,y) = 2x^2 + y^2$ and $S(x,y,z) = x^2 + y^2 + z^2$. (Notice that S is, like, a 4-dimensional function – three inputs and one output.)

- (a) Write equations and sketch graphs for the $y = -2$, $y = 0$, and $y = 2$ traces of C .
- (b) Write equations and sketch graphs for the $x = -2$, $x = 0$, and $x = 2$ traces of C .
- (c) What can you say about the general shape of x -traces of this function? How about y -traces? How are they similar and how are they different?

- (d) Write equations and sketch graphs for the level curves of C at 0, 2, and 4.
- (e) What can you say about the general shape of the contours of this function?
- (f) Why didn't I ask about contours at -2 or -4?

- (g) Write an equation for the contour of S at 1.
- (h) What's this equation?
(Hint: you can graph it in CalcPlot3D if you use the "implicit surface" option.)
- (i) Maybe now I can't call this a "level curve" but instead I should call it a "level surface." Why?

Checkpoint: Partial derivatives

The gas law for a fixed mass m of an ideal gas at absolute temperature T in Kelvins, pressure P in pascals, and volume V in liters is $PV = mRT$, where R is the gas constant.

- Compute each of the partial derivatives below,
- assign them the correct units,
- and say a sentence or two about what each one means. (Be careful to think about what is *constant* and what is *changing*.)

(a) $\frac{\partial P}{\partial V}$

(b) $\frac{\partial V}{\partial T}$

(c) $\frac{\partial T}{\partial P}$

(Bonus question for the physics-knowers: Why does the sign of each one make sense?)

Checkpoint: Differentials slash linear approximation slash tangent planes

Wolfram Alpha tells me that $\sqrt{(4.98)^2 - (3.03)^2} = 3.952151312892764231189305743300570249811443622895$


This is, however, way too precise to be useful. Show how to compute a reasonable estimate of this value **without a calculator** – I bet your answer will agree with the actual value in at least the first two or three decimal places.

Hints:

- Consider the function $f(x, y) = \sqrt{x^2 - y^2}$ and think about the title of this checkpoint.
- Do you remember how to multiply decimals by hand? To find $1.5 * 0.2$, you can compute $1.5 * 2$ or $15 * 2$, then move the decimal point in your result an appropriate number of places.

Checkpoint: The gradient


Consider the function $f(x, y) = x^2 + 3y^2$; here's a contour plot of the function but with the contours deliberately not labeled.



- (a) Compute $\nabla f(x, y)$.
- (b) At the point $(1, 2)$, what is the direction of greatest increase?
- (c) Draw a vector on the contour plot showing the direction of greatest increase.
- (d) Find the slope of f in this direction.

Checkpoint: Directional derivatives

The function $T(x,y) = 30 - x^2 - y^2$ measures the temperature, in degrees Celsius, at a given point (x,y) on a heated metal plate, where x and y are measured in centimeters. An ant is walking on this plate in a straight line from the point $(1,2)$ to the point $(4,5)$. Here's a contour plot with some relevant points highlighted.



- How fast is the temperature under the ant's feet changing at the beginning of its journey? Give units.
- How fast is the temperature under the ant's feet changing when it's halfway through its journey? (Hint: if you draw this out you'll see that this is at the point $(3,3)$.)
- The ant reaches $(4,5)$ and decides to walk directly back to the origin. Now how fast is the temperature under the ant's feet changing?

Checkpoint: Optimization

The temperature T of the disk $x^2 + y^2 \leq 1$ is given by $T(x, y) = 2x^2 - 3y^2 - 2x$. Find the hottest and coldest points of the disk. A contour plot of this function in these boundaries is shown below.

