

Subgroups!

Spencer Bagley

With many thanks to Matthew Macauley,
<http://www.math.clemson.edu/~macaule/>

10 Feb 2025

Goals for today:

1. Define what subgroups are

Goals for today:

1. Define what subgroups are
2. See some examples

Goals for today:

1. Define what subgroups are
2. See some examples
3. Figure out all the subgroups of all the groups of order 4

Goals for today:

1. Define what subgroups are
2. See some examples
3. Figure out all the subgroups of all the groups of order 4
4. ... of order 6

Goals for today:

1. Define what subgroups are
2. See some examples
3. Figure out all the subgroups of all the groups of order 4
4. ... of order 6
5. ... of order 8

Definition time!

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.
2. G contains the identity element.

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.
2. G contains the identity element.
3. Every element in G has an inverse element.

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.
2. G contains the identity element.
3. Every element in G has an inverse element.
4. **G is closed under the binary operation.**

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.
2. G contains the identity element.
3. Every element in G has an inverse element.
4. G is closed under the binary operation.

Trivial subgroups

Every group G has the following two boring subgroups: $G \leq G$, and $\{e\} \leq G$.

Definition time!

Here is the definition of a subgroup.

Definition

A **subgroup** of G is a subset $H \subseteq G$ that is also a group. We denote this by $H \leq G$.

Okay, but remind me what's the definition of a group?

Definition

A **group** (G, \star) is a set of elements together with a binary operation \star satisfying the following properties:

1. The operation is associative.
2. G contains the identity element.
3. Every element in G has an inverse element.
4. G is closed under the binary operation.

Trivial subgroups

Every group G has the following two boring subgroups: $G \leq G$, and $\{e\} \leq G$.

Definition

A **proper subgroup** $H < G$ is a subgroup that's not equal to the whole group.

Generating sets

We've previously looked at the **orbit** of an element:

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

But we need not restrain ourselves to generating by one element:

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

But we need not restrain ourselves to generating by one element:

Definition

Let S be a sub**set** of G .

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

But we need not restrain ourselves to generating by one element:

Definition

Let S be a sub**set** of G . A **word** in S is a finite product of finite powers of elements of S or their inverses.

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

But we need not restrain ourselves to generating by one element:

Definition

Let S be a **subset** of G . A **word** in S is a finite product of finite powers of elements of S or their inverses.

$\langle S \rangle = \{\text{words in } S\}$ is a subgroup of G , and it's called the **subgroup generated by S** .

Generating sets

We've previously looked at the **orbit** of an element:

Definition

The **orbit** of an element $g \in G$ is the **cyclic subgroup** that it generates,

$$\langle g \rangle = \{g^k \mid k \in \mathbb{Z}\},$$

and its **order** is $|g| := |\langle g \rangle|$.

In particular, if $|g| = n$ is finite, this is the set $\{g^0 = 1, g, g^2, \dots, g^{n-1}\}$.

This is a subgroup:

Cyclic subgroups are subgroups

For any element $g \in G$, $\langle g \rangle \leq G$.

But we need not restrain ourselves to generating by one element:

Definition

Let S be a **subset** of G . A **word** in S is a finite product of finite powers of elements of S or their inverses.

$\langle S \rangle = \{\text{words in } S\}$ is a subgroup of G , and it's called the **subgroup generated by S** .

And in fact every subgroup looks like this.

Example: $C_2 \leq D_3$


Writing $C_2 \leq D_3$ means *there is a copy of C_2 sitting inside of D_3 as a subgroup.*

Example: $C_2 \leq D_3$

Writing $C_2 \leq D_3$ means *there is a copy of C_2 sitting inside of D_3 as a subgroup.*

Question

How many ways can you find C_2 sitting inside of D_3 ?




Example: $C_2 \leq D_3$

Writing $C_2 \leq D_3$ means *there is a copy of C_2 sitting inside of D_3 as a subgroup*.

Question

How many ways can you find C_2 sitting inside of D_3 ?



Remark


It's more precise to express a subgroup by its generator(s).

Example: $C_2 \leq D_3$

Writing $C_2 \leq D_3$ means *there is a copy of C_2 sitting inside of D_3 as a subgroup.*

Question

How many ways can you find C_2 sitting inside of D_3 ?



Remark

It's more precise to express a subgroup by its generator(s).

$$C_2 \cong \langle f \rangle < D_3$$

$$C_2 \cong \langle rf \rangle < D_3$$

$$C_2 \cong \langle r^2f \rangle < D_3$$

Question


How about $C_3 \leq D_3$?

Example: $C_2 \leq D_3$

Writing $C_2 \leq D_3$ means *there is a copy of C_2 sitting inside of D_3 as a subgroup.*

Question

How many ways can you find C_2 sitting inside of D_3 ?



Remark

It's more precise to express a subgroup by its generator(s).

$$C_2 \cong \langle f \rangle < D_3$$

$$C_2 \cong \langle rf \rangle < D_3$$

$$C_2 \cong \langle r^2f \rangle < D_3$$



Question

How about $C_3 \leq D_3$? *There's only one!*

Groups of order 4



The two groups of order 4

Let's start by considering the subgroups of the two groups of order 4.



The two groups of order 4



Let's start by considering the subgroups of the two groups of order 4.



- Proper subgroups of V_4 : $\langle h \rangle = \{e, h\}$, $\langle v \rangle = \{e, v\}$, $\langle r \rangle = \{e, r\}$, $\langle e \rangle = \{e\}$.

The two groups of order 4



Let's start by considering the subgroups of the two groups of order 4.



- Proper subgroups of V_4 : $\langle h \rangle = \{e, h\}$, $\langle v \rangle = \{e, v\}$, $\langle r \rangle = \{e, r\}$, $\langle e \rangle = \{e\}$.
- Subgroups of C_4 : $\langle r \rangle = \{1, r, r^2, r^3\} = \langle r^3 \rangle$, $\langle r^2 \rangle = \{1, r^2\}$, $\langle 1 \rangle = \{1\}$.



The two groups of order 4

Let's start by considering the subgroups of the two groups of order 4.



- Proper subgroups of V_4 : $\langle h \rangle = \{e, h\}$, $\langle v \rangle = \{e, v\}$, $\langle r \rangle = \{e, r\}$, $\langle e \rangle = \{e\}$.
- Subgroups of C_4 : $\langle r \rangle = \{1, r, r^2, r^3\} = \langle r^3 \rangle$, $\langle r^2 \rangle = \{1, r^2\}$, $\langle 1 \rangle = \{1\}$.

It is illustrative to arrange these in a [subgroup lattice](#):



Order: 4


2

1


Groups of order 6

Subgroups of \mathbb{Z}_6

What subgroups can you find in \mathbb{Z}_6 ? I've drawn the Cayley diagram two different ways.





Hello I am secretly also the cycle graph





Subgroups of D_3

Let's figure out all the subgroups of D_3 .



Subgroups of D_3

Let's figure out all the subgroups of D_3 .





Here are the **non-trivial proper subgroups** of D_3 :

$$\langle r \rangle = \{1, r, r^2\} = \langle r^2 \rangle, \quad \langle f \rangle = \{1, f\}, \quad \langle rf \rangle = \{1, rf\}, \quad \langle r^2f \rangle = \{1, r^2f\}$$

Subgroups of D_3

Let's figure out all the subgroups of D_3 .




Here are the **non-trivial proper subgroups** of D_3 :

$$\langle r \rangle = \{1, r, r^2\} = \langle r^2 \rangle, \quad \langle f \rangle = \{1, f\}, \quad \langle rf \rangle = \{1, rf\}, \quad \langle r^2f \rangle = \{1, r^2f\}, \quad \langle 1 \rangle = \{1\}.$$

Subgroups of D_3

Let's figure out all the subgroups of D_3 .



Here are the **non-trivial proper subgroups** of D_3 :


$$\langle r \rangle = \{1, r, r^2\} = \langle r^2 \rangle, \quad \langle f \rangle = \{1, f\}, \quad \langle rf \rangle = \{1, rf\}, \quad \langle r^2f \rangle = \{1, r^2f\}, \quad \langle 1 \rangle = \{1\}.$$

Observations:

- The cycle graph helps us spot cyclic subgroups.

Subgroups of D_3

Let's figure out all the subgroups of D_3 .



Here are the **non-trivial proper subgroups** of D_3 :


$$\langle r \rangle = \{1, r, r^2\} = \langle r^2 \rangle, \quad \langle f \rangle = \{1, f\}, \quad \langle rf \rangle = \{1, rf\}, \quad \langle r^2f \rangle = \{1, r^2f\}, \quad \langle 1 \rangle = \{1\}.$$

Observations:

- The cycle graph helps us spot cyclic subgroups.
- For small groups like D_3 , the cyclic subgroups may be the **only** proper subgroups.

Subgroups of D_3

Let's figure out all the subgroups of D_3 .



Here are the **non-trivial proper subgroups** of D_3 :

$$\langle r \rangle = \{1, r, r^2\} = \langle r^2 \rangle, \quad \langle f \rangle = \{1, f\}, \quad \langle rf \rangle = \{1, rf\}, \quad \langle r^2f \rangle = \{1, r^2f\}, \quad \langle 1 \rangle = \{1\}.$$




Observations:

- The cycle graph helps us spot cyclic subgroups.
- For small groups like D_3 , the cyclic subgroups may be the **only** proper subgroups.
- There might, however, be more complicated things that are harder to clock.

Groups of order 8




Subgroups of D_4

See if you can figure out all the subgroups of D_4 .



Subgroups of D_4

See if you can figure out all the subgroups of D_4 .



What do you think is a reasonable way to, like, arrange them?

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:

■ **join**, or **sup**, or **least upper bound**

$x \vee y$

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:

■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:



■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Examples you may have seen previously are **subset lattices** and **divisor lattices**.



This seems like a good way to organize subgroups, because:

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:



■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Examples you may have seen previously are **subset lattices** and **divisor lattices**.



This seems like a good way to organize subgroups, because:

Theorem

If $H \leq G$ and $K \leq G$ are two subgroups, then $H \cap K$ is a subgroup.

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:



■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Examples you may have seen previously are **subset lattices** and **divisor lattices**.



This seems like a good way to organize subgroups, because:

Theorem

If $H \leq G$ and $K \leq G$ are two subgroups, then $H \cap K$ is a subgroup.
(Indeed, it's the largest subgroup that's contained in both H and K .)

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:



■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Examples you may have seen previously are **subset lattices** and **divisor lattices**.



This seems like a good way to organize subgroups, because:

Theorem

If $H \leq G$ and $K \leq G$ are two subgroups, then $H \cap K$ is a subgroup.
(Indeed, it's the largest subgroup that's contained in both H and K .)

Theorem

$\langle H, K \rangle$ is the smallest subgroup containing both H and K .

Lattices

A **lattice** is a **partially ordered set** such that every pair of elements x, y has a **unique**:



■ **join**, or **sup**, or **least upper bound**

$$x \vee y$$

■ **meet**, or **inf**, or **greatest lower bound**

$$x \wedge y.$$

Examples you may have seen previously are **subset lattices** and **divisor lattices**.



This seems like a good way to organize subgroups, because:


Theorem

If $H \leq G$ and $K \leq G$ are two subgroups, then $H \cap K$ is a subgroup.
(Indeed, it's the largest subgroup that's contained in both H and K .)

Theorem

$\langle H, K \rangle$ is the smallest subgroup containing both H and K .
(Note that $H \cup K$ is not in general a subgroup. Why not?)


Subgroup lattices



$H \vee K$: “smallest subgroup above both H and K ”

$H \wedge K$: “largest subgroup below both H and K ”


Subgroup lattices




$H \vee K$: “smallest subgroup above both H and K ”

$H \wedge K$: “largest subgroup below both H and K ”

Examples:





Subgroup lattices



$H \vee K$: “smallest subgroup above both H and K ”

$H \wedge K$: “largest subgroup below both H and K ”

Examples:



The subgroup lattice of D_4

The subgroups of D_4 are:

The subgroup lattice of D_4

The subgroups of D_4 are:

- The entire group D_4 , and the trivial group $\langle 1 \rangle$

The subgroup lattice of D_4

The subgroups of D_4 are:

- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2 f \rangle$, $\langle r^3 f \rangle$.

The subgroup lattice of D_4

The subgroups of D_4 are:

- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2 f \rangle$, $\langle r^3 f \rangle$.
- 1 subgroup generated by a 180° rotation, $\langle r^2 \rangle \cong C_2$

The subgroup lattice of D_4


The subgroups of D_4 are:

- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2 f \rangle$, $\langle r^3 f \rangle$.
- 1 subgroup generated by a 180° rotation, $\langle r^2 \rangle \cong C_2$
- 1 subgroup generated by a 90° rotation, $\langle r \rangle \cong C_4$

The subgroup lattice of D_4

The subgroups of D_4 are:




- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2 f \rangle$, $\langle r^3 f \rangle$.
- 1 subgroup generated by a 180° rotation, $\langle r^2 \rangle \cong C_2$
- 1 subgroup generated by a 90° rotation, $\langle r \rangle \cong C_4$
- 2 subgroups isomorphic to V_4 : $\langle r^2, f \rangle$, $\langle r^2, rf \rangle$.



The subgroup lattice of D_4

The subgroups of D_4 are:





- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2f \rangle$, $\langle r^3f \rangle$.
- 1 subgroup generated by a 180° rotation, $\langle r^2 \rangle \cong C_2$
- 1 subgroup generated by a 90° rotation, $\langle r \rangle \cong C_4$
- 2 subgroups isomorphic to V_4 : $\langle r^2, f \rangle$, $\langle r^2, rf \rangle$.




The subgroup lattice of D_4

The subgroups of D_4 are:


- The entire group D_4 , and the trivial group $\langle 1 \rangle$
- 4 subgroups generated by reflections: $\langle f \rangle$, $\langle rf \rangle$, $\langle r^2 f \rangle$, $\langle r^3 f \rangle$.
- 1 subgroup generated by a 180° rotation, $\langle r^2 \rangle \cong C_2$
- 1 subgroup generated by a 90° rotation, $\langle r \rangle \cong C_4$
- 2 subgroups isomorphic to V_4 : $\langle r^2, f \rangle$, $\langle r^2, rf \rangle$.



The subgroup lattice of D_4



The subgroup lattice of D_4



The subgroup lattice of Q_8


Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group


$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$



The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$




Every element generates a **cyclic subgroup**:

$$\langle 1 \rangle = \{1\},$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$




Every element generates a **cyclic subgroup**:

$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\},$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$




Every element generates a **cyclic subgroup**:

$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\}, \quad \langle i \rangle = \langle -i \rangle = \{\pm 1, \pm i\},$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$



Every element generates a **cyclic subgroup**:


$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\}, \quad \langle i \rangle = \langle -i \rangle = \{\pm 1, \pm i\},$$

$$\langle j \rangle = \langle -j \rangle = \{\pm 1, \pm j\},$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$



Every element generates a **cyclic subgroup**:


$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\}, \quad \langle i \rangle = \langle -i \rangle = \{\pm 1, \pm i\},$$

$$\langle j \rangle = \langle -j \rangle = \{\pm 1, \pm j\}, \quad \langle k \rangle = \langle -k \rangle = \{\pm 1, \pm k\}.$$

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$



Every element generates a **cyclic subgroup**:

$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\}, \quad \langle i \rangle = \langle -i \rangle = \{\pm 1, \pm i\},$$


$$\langle j \rangle = \langle -j \rangle = \{\pm 1, \pm j\}, \quad \langle k \rangle = \langle -k \rangle = \{\pm 1, \pm k\}.$$

Are there any other proper subgroups?

The subgroup lattice of Q_8

Let's determine all subgroups of the quaternion group

$$Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle.$$




Every element generates a **cyclic subgroup**:

$$\langle 1 \rangle = \{1\}, \quad \langle -1 \rangle = \{\pm 1\}, \quad \langle i \rangle = \langle -i \rangle = \{\pm 1, \pm i\},$$

$$\langle j \rangle = \langle -j \rangle = \{\pm 1, \pm j\}, \quad \langle k \rangle = \langle -k \rangle = \{\pm 1, \pm k\}.$$

Are there any other proper subgroups?



Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

We've seen the subgroup lattices of two groups of order 8:

Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.

Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.

Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.
- \mathbb{Z}_2^3 has seven *elements* of order 2.

Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.
- \mathbb{Z}_2^3 has seven *elements* of order 2.

Rule of thumb

Groups with elements of small order tend to have more subgroups than those with elements of large order.

Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$


We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.
- \mathbb{Z}_2^3 has seven *elements* of order 2.

Rule of thumb

Groups with elements of small order tend to have more subgroups than those with elements of large order.

The following Cayley graphs show three different subgroups of order 4 in \mathbb{Z}_2^3 .



Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$



We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.
- \mathbb{Z}_2^3 has seven *elements* of order 2.

Rule of thumb

Groups with elements of small order tend to have more subgroups than those with elements of large order.

The following Cayley graphs show three different subgroups of order 4 in \mathbb{Z}_2^3 .



Subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$




We've seen the subgroup lattices of two groups of order 8:

- D_4 has five elements of order 2, and 10 subgroups.
- Q_8 has one element of order 2, and 6 subgroups.
- \mathbb{Z}_2^3 has seven *elements* of order 2.

Rule of thumb

Groups with elements of small order tend to have more subgroups than those with elements of large order.

The following Cayley graphs show three different subgroups of order 4 in \mathbb{Z}_2^3 .



The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

But this triple-counts all such subgroups.

The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

But this triple-counts all such subgroups. In summary, the subgroups of \mathbb{Z}_2^3 are:

- The subgroups G and $\{000\}$,

The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

But this triple-counts all such subgroups. In summary, the subgroups of \mathbb{Z}_2^3 are:

- The subgroups G and $\{000\}$,
- 7 subgroups isomorphic to C_2 ,

The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

But this triple-counts all such subgroups. In summary, the subgroups of \mathbb{Z}_2^3 are:

- The subgroups G and $\{000\}$,
- 7 subgroups isomorphic to C_2 ,
- 7 subgroups isomorphic to V_4 .


The subgroup lattice of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

All 7 non-identity elements generate a subgroup isomorphic to C_2 .

All $\binom{7}{2} = 21$ pairs of non-identity elements generate a subgroup isomorphic to V_4 .

But this triple-counts all such subgroups. In summary, the subgroups of \mathbb{Z}_2^3 are:

- The subgroups G and $\{000\}$,
- 7 subgroups isomorphic to C_2 ,
- 7 subgroups isomorphic to V_4 .




The subgroup lattice of \mathbb{Z}_8

Draw the Cayley diagram of \mathbb{Z}_8 and find all its subgroups.
Arrange them in a lattice.



The subgroup lattice of \mathbb{Z}_8

Draw the Cayley diagram of \mathbb{Z}_8 and find all its subgroups.
Arrange them in a lattice.





The subgroup lattice of \mathbb{Z}_8

Draw the Cayley diagram of \mathbb{Z}_8 and find all its subgroups.
Arrange them in a lattice.



The subgroup lattice of \mathbb{Z}_8

Draw the Cayley diagram of \mathbb{Z}_8 and find all its subgroups.
Arrange them in a lattice.



Theorem


Every subgroup of a cyclic group is cyclic.

Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.



Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.





Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.



Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.





Let's summarize the sizes of the subgroups of the groups of order 8 that we have seen.

	C_8	Q_8	$C_4 \times C_2$	D_4	C_2^3
# elts. of order 8	4	0	0	0	0
# elts. of order 4	2	6	4	2	0
# elts. of order 2	1	1	3	5	7
# elts. of order 1	1	1	1	1	1
# subgroups	4	6	8	10	16

Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.





Let's summarize the sizes of the subgroups of the groups of order 8 that we have seen.

	C_8	Q_8	$C_4 \times C_2$	D_4	C_2^3
# elts. of order 8	4	0	0	0	0
# elts. of order 4	2	6	4	2	0
# elts. of order 2	1	1	3	5	7
# elts. of order 1	1	1	1	1	1
# subgroups	4	6	8	10	16

Observations?

Groups of order 8

There is one more group of order 8, which is $\mathbb{Z}_4 \times \mathbb{Z}_2$.



Let's summarize the sizes of the subgroups of the groups of order 8 that we have seen.

	C_8	Q_8	$C_4 \times C_2$	D_4	C_2^3
# elts. of order 8	4	0	0	0	0
# elts. of order 4	2	6	4	2	0
# elts. of order 2	1	1	3	5	7
# elts. of order 1	1	1	1	1	1
# subgroups	4	6	8	10	16

Observations?

- Groups that have more elements of small order tend to have more subgroups.
- In all of these cases, the order of each subgroup divides $|G|$.

The end!