

REVIEW

Colour in digital pathology: a review

Emily L Clarke^{1,2} & Darren Treanor^{1,2}

¹Department of Histopathology, Leeds Teaching Hospitals NHS Trust, and ²Section of Pathology and Tumour Biology, Leeds Institute of Cancer and Pathology, University of Leeds, Leeds, UK

Clarke E L & Treanor D

(2017) *Histopathology* 70, 153–163. DOI: 10.1111/his.13079

Colour in digital pathology: a review

Colour is central to the practice of pathology because of the use of coloured histochemical and immunohistochemical stains to visualize tissue features. Our reliance upon histochemical stains and light microscopy has evolved alongside a wide variation in slide colour, with little investigation into the implications of colour variation. However, the introduction of the digital microscope and whole-slide imaging has highlighted the need for further understanding and control of colour. This is because the digitization process itself introduces further colour variation which may affect diagnosis, and image analysis algorithms often use colour or intensity measures to detect or measure

tissue features. The US Food and Drug Administration have released recent guidance stating the need to develop a method of controlling colour reproduction throughout the digitization process in whole-slide imaging for primary diagnostic use. This comprehensive review introduces applied basic colour physics and colour interpretation by the human visual system, before discussing the importance of colour in pathology. The process of colour calibration and its application to pathology are also included, as well as a summary of the current guidelines and recommendations regarding colour in digital pathology.

Keywords: colour calibration, colour standardization, digital pathology, whole slide imaging

Introduction

Colour is central and integral to the practice of pathology; pathologists use coloured histochemical and immunohistochemical stains to identify structures and reach diagnoses. Given the magnitude of these diagnoses to patient management and outcome, it is imperative that pathologists are able to make accurate and reliable assertions.

During recent years, new technology has emerged to enable pathologists to carry out their diagnostic work digitally, rather than with the analogue optical microscope. This technology is often referred to as a 'digital microscope' in which whole slide images (WSIs) of tissue slides are scanned, transmitted and displayed on a computer monitor. While the method of making

diagnoses may be gradually evolving, the need for diagnostic reliability remains a constant. Concerns that WSI may not be an accurate representation of the pathology may, in part, explain why digital pathology has not yet become part of routine clinical practice. To try to resolve these issues, there has been a recent surge in research in the field of WSI validation;¹ however, the impact of colour remains unknown.

The aim of this review is to provide a summary of the issues surrounding colour control in digital microscopy for pathologists. This necessitates an initial brief introduction to the physics of colour and its interpretation by the human visual system, but much of the vast field of colour science is beyond the scope of this work.

Colour basics

Colour is 'an attribute of things that results from the light they reflect, transmit, or emit in so far as this

Address for correspondence: Dr Emily L Clarke, Leeds Institute of Cancer and Pathology, University of Leeds, Beckett Street, Leeds LS9 7TF, UK. e-mail: e.l.clarke@leeds.ac.uk

light causes a visual sensation that depends on its wavelengths² - light with different wavelengths are perceived as different colours. Those colours that are formed by a single wavelength of light are termed 'monochromatic colours' (e.g. red, orange, violet, green, blue, yellow). These make up our colour spectrum including wavelengths from 390 to 700 nm. Light with wavelengths outside this spectrum are not visible to humans, and therefore do not form part of our colour spectrum. These include infrared with wavelengths greater than 1000 nm and ultraviolet with wavelengths less than 300 nm (Figure 1).

The colour of an object is the result of our interpretation of the object's surface, transmission and emission properties. There are three main ways in which objects can affect light. First, opaque objects reflect light; they can do so 'specularly', like a mirror, or 'scattered' with diffuse scattering, which eludes to a 'roughened' surface. Secondly, objects that transmit light appear transparent (no scattering) or translucent (with scattering). Thirdly, those that emit light have excited electrons secondary to a chemical reaction (chemoilluminescence), an elevated temperature (incandescence) or from absorbing light at other frequencies (fluorescence).^{3,4} Pathology tissue slides are translucent when being viewed using light microscopy - in other words, the colours they exhibit are due to transmission of light with scattering.

In 1730, Newton demonstrated that white light, for example from the sun, can be dispersed with a prism into all visible wavelengths. This is a perfect demonstration of additive colour mixing; colours can be added together to make white. Additive colours are produced by objects which themselves create light; for example, computer monitors, rainbows, fireworks. By contrast, subtractive colours combine to

make black and are produced by objects that do not emit light, e.g. colour printing, photographs or fine art. It therefore follows that glass microscope slides are comprised of subtractive colours, whereas the digital microscope displays additive colours. It is important to ensure that diagnostic information contained within the specimen is unaffected by the conversion from a subtractive colour system to an additive colour system.⁵

In order to describe, organize and categorize colours we can employ the use of a colour space (Figure 2). There are various colour spaces, one of the most common being the hue, saturation and value (HSV) colour space (Figure 3) which comprises three qualities for each colour: hue, saturation and value. Each colour will have a defined value in each of these three qualities and can therefore be measured and replicated accurately. A colour's 'hue' is the main discernible attribute of a colour, e.g. red. 'Saturation' is the intensity of a colour and 'value' is the lightness of a colour. An alternative to the HSV colour space is the RGB (red, green, blue) colour space (Figure 2). In this colour space, instead of each colour being defined by hue, value and saturation, each is given a value for red, green and blue (normally from 0 to 256 in an 8-bit-display).⁵

Colour and the human visual system

As highlighted previously, the colour of an object is influenced significantly by human perception and interpretation; indeed, it is likely that individuals view and interpret colour differently.⁶ As a consequence, the role of colour in medical image interpretation is a very complex topic spanning physics, engineering, physiology and psychology.^{7,8} However, a brief




Figure 1. The spectrum of light. Source: https://upload.wikimedia.org/wikipedia/commons/thumb/f/f1/EM_spectrum.svg/2000px-EM_spectrum.svg.png.

Figure 2. Demonstration of additive and subtractive colour mixing. Representation of additive colours are generally portrayed using the red, green, blue (RGB) colourspace. Representation of subtractive colours are demonstrated normally using the cyan–magenta–yellow–key (CMYK) colourspace. Source: <https://commons.wikimedia.org/wiki/File:Subtractive-Additive-Colour-Mixing.jpg>.





Figure 3. Graphical representation of the hue, saturation and value (HSV) colourspace. Assigning a value to each of the qualities enables accurate categorization and replication of each colour. Source: https://upload.wikimedia.org/wikipedia/commons/4/4e/HSV_color_solid_cylinder.png.

summary of how the human visual system interprets images is provided below.

The human eye can identify up to 10 million different colours,⁸ yet the sensitivity for some hues and saturation is stronger than others. This was demonstrated by MacAdam's ellipses in the CIELAB colour-space in 1942. His psychophysical experiments into Just Noticeable Differences (JNDs) highlighted that the Human Visual System (HVS) appears more sensitive to changes in purple than other colours.^{5,9} It is therefore perhaps fortuitous (and perhaps no coincidence) that the most common stain, haematoxylin and eosin (H&E), is purple and pink.

While the HVS is very sensitive to changes to hue and saturation, it is also very adaptive to brightness.

It can adapt to 14 decades (orders of magnitude) of brightness, making use of two mechanisms; light adaption and dark adaption. Light adaption occurs when entering a brightly lit environment from a dark one, and takes approximately 5 min to lower sensitivity to illumination levels. Conversely, dark adaption takes approximately 30 min, increases visual sensitivity and enables some degree of night vision. It is this change in illumination sensitivity that results in a lit candle appearing to be much brighter in a dark room than within a brightly lit environment.

The HVS is also capable of chromatic adaptation. This is the biological equivalence of 'automatic white balance', whereby the surrounding illumination will have an effect on the perceived colour.¹⁰ This can be observed by examining a piece of white paper under many different illuminations, e.g. in daylight, fluorescent and incandescent, and observing the difference. This adaptation allows a phenomenon called 'colour constancy' to occur – this is the way in which the brain adjusts perceived colours based on their surroundings. For example, a banana appears yellow on a bright sunny day and yellow during a candlelit meal, even though the illumination is significantly different.¹¹

A recent study has provided evidence of chromatic adaption occurring in the natural setting. Perception of colour changes with the seasons; during the summer, when we are surrounded by 'green', we compensate for this by shifting our 'unique yellow' settings.¹² It is currently unknown how chromatic adaptation influences the perceived colour of whole slide images, but in order to attempt to minimize variation of interpretation it may be advisable to standardize the illuminance of digital microscope scanners and displays.

Colour appearance is also affected by various visual cognitive mechanisms, including 'memory colour'.¹³ Memory colour is a phenomenon whereby people associate a particular colour with a recognizable object. Interestingly, however, their 'memory' of the colour is often not based in reality, as people often remember colours as more saturated than in actuality. This may have a bearing on pathologists' ability to compare accurately colours from glass slides with the colours of WSIs.

The HVS also changes as part of the natural ageing process, which acts to decrease colour vision. The cones within the retina (the cells responsible for colour vision) decline in sensitivity, rendering images less bright with reduced colour discrimination. Most commonly, blue colours appear more 'washed out' than others,¹⁴ resulting in the majority of colour defects in the elderly population being of blue–yellow type.¹⁵ While the effect of colour-blindness among pathologists has been evaluated in a handful of studies,^{16,17} the effect of ageing on interpreting slides is unknown, and is likely to be more complex than simple physical visual changes.

Does colour matter in pathology?

This question is hotly debated, due primarily to a lack of definitive evidence either way. It is understandable that, given our ability to adapt well to variations in colour, some conclude that colour does not matter in the field of digital pathology. However, evidence of adaptation is not a reason against control of colour variation, as this argument overlooks key points. Firstly, the implications of adaptation to colour variation have not been evaluated, and so lack of colour standardization may be a hindrance to the diagnostic process. Secondly, while we are certainly adept at dealing with variation in colour, our ability to adapt has not been well studied or quantified and, as such, we may only be able to compensate for colour variation to a certain degree.

A further argument against the need for accurate colour representation in digital pathology is the existence of colour blind pathologists. This idea is reinforced by a recent small study by Campbell *et al.*¹⁸ indicating a 92.7% concordance between whole slide images converted to greyscale and the original glass diagnoses in breast biopsy cases. While this may seem a logical argument, a study by Poole *et al.*¹⁷ indicated that colour-blind pathologists had a lower mean score (94 versus 99%) compared to their colleagues with normal colour vision when trying to identify

pathological features. This finding is supported – amusingly – by a recent paper demonstrating that pigeons (*Columba livia*) had a reduced accuracy for detecting breast cancer in using monochrome images compared to full colour.¹⁹ Furthermore, a survey of pathologists by Akman *et al.*¹⁶ reported that 61.3% of pathologists interviewed felt that colour-blind pathologists should not perform pathology – thereby indicating that many pathologists feel that colour interpretation is important in the diagnostic process.

A further argument for the importance of colour in digital pathology is the common laboratory practice of recutting and restaining referred or 'foreign' slides indicating that, at the very least, pathologists prefer working with familiar colours and that this may possibly impact upon their ability, confidence or speed of diagnosis. This perceived preference for certain colours among pathologists has also been demonstrated in our experience with WSIs after a recent external quality assurance (EQA) raised concerns from pathologists that stain irregularities would compromise their diagnostic ability.²⁰ Also, few pathologists would argue that colour is non-essential in the realm of immunostains.

Regardless of this divide in professional opinion, in the field of colour science there is agreement that colour control is a necessary step in digital imaging, and the US Food and Drug Administration (FDA)²¹ have released guidance stating that digital microscope images should be displayed in a consistent and reliable fashion.

Colour variation in pathology

Unfortunately, colour variation is substantial in pathology; this has been acknowledged anecdotally for some time, but has only recently been quantified formally. Disparities in routine H&E staining was evaluated by Gray *et al.*^{22,23} by staining the same piece of tissue, scanning it into a digital slide scanner and performing image deconvolution. While there was high repeatability in the H&E ratio when staining on the same day (mean difference 0.47%), the H&E ratio varied considerably when stained on different days (mean difference 8.32%) (Figure 4).

Unfortunately, even the advent of automated staining does not resolve the issue – further work has demonstrated H&E ratio differences of more than 100% between four different automated staining instruments from the same manufacturer.^{22,23} Despite these known inconsistencies, anecdotally it seems that pathologists are able to make successful




Figure 4. A selection of slides from Gray *et al.*²³ demonstrating routine wide variation in staining. These four different slides of the same piece of appendix were stained on different days. The wide range of different colour stains in routine practice is clearly evident.

diagnoses from slides with a wide variety of staining variation and using a range of optical microscopes.

Colour consistency is much more of an issue with the introduction of the digital microscope, as it does not enable the pathologist to view the tissue directly and colour variation is not confined to irregularities within the staining process. Therefore, digitization of the slide introduces further lack of colour control, compounding the variation already introduced through the slide-staining process.

There are multiple stages involved in making a digital image, each of which may influence slide colour substantially. These stages include sample illumination, magnification, image capture, compression, storage, transmission and finally reproduction on the computer display (Figure 5). To the best of our knowledge, the effect on the slide colour from each of these individual steps in the digitization process has not been quantified. However, the overall effect of colour variation due to scanning the slide was investigated by Gray *et al.*,²² with a mean difference of 7% in H&E ratio when scanning the same slide into the differing scanners on the same day. It should be noted, however, that this measurement does not take into account the effect of colour variation due to the display, so the overall impact of digitization on colour variation from scanning to display is likely to be greater.

Colour constancy in digital imaging is measured in a unit called Delta-E (dE) introduced by the International Commission on Illumination (CIE). This is a number quantifying the change in visual perception between two colours. The smallest perceptual change in colour is generally regarded as 1 dE, with 2–10 dE being perceptible at a glance, 11–49 dE colours are more similar than opposite and 100 dE representing exact opposite colours.²⁴ A study by Shrestha *et al.*²⁵ indicated an average colour discrepancy of 10 dE in uncalibrated scanners alone, so it is easy to imagine the wide variation in colours that are derived from inconsistencies at every stage in digitization of slides.

The effect of colour differences on human interpretation of digitized pathology images has not been studied widely; however, pathologists themselves have raised concerns that colour variability may impact negatively on their diagnostic performance. This was noted formally in feedback from two recent national EQA programmes in the United Kingdom. In one EQA, 14 of 84 comments expressed discontent with the WSI colour and four participants expressly mentioned concern that the variation in colour may impact upon their diagnostic performance negatively.²⁰ However, whether subjective comments reflect diagnostic ability remains unknown.²⁶

Differences in slide colour may also have serious implications for the reproducibility of image analysis




Figure 5. The stages required in capturing and displaying a digital slide. Each of the stages shown may have the capacity to substantially influence digital slide colour.

algorithms. There are numerous papers suggesting algorithms for digital analysis of immunostains,^{27–31} but only a handful consider the huge implications for colour variation in WSI.^{32,33} Gavrielides *et al.*³⁴ reported variation in colour between three different WSI instruments from two manufacturers leading to variation in performance of image analysis algorithm for HER2/*neu*. Approximately 20–30% of cases scored as 2+ expression on one scanner were rescored to a different class on another instrument.

Colour calibration

Colour calibration is an imaging process which seeks to match colours between two or more devices. End-to-end calibration describes the process of controlling colour from source to output through each step of the imaging pathway. Colour calibration is a routine process in the print and photography industry and has been adopted by most digital systems. Even smartphones can be colour-calibrated since the advent of the SpyderGALLERY application from Datacolour.³⁵

Colour calibration works by comparing 'known' colours from a set of colour patches with the colours of those same patches when an image is taken of them with the device. The differences between the 'known' colour values for each of the patches and the colour values acquired through imaging the patches allow for numerical identification of the deficiencies within that specific imaging system and allow for necessary adjustments to be made. Compensation for those differences is afforded with the use of an ICC (International Colour Consortium) colour profile, which can then be used to calibrate subsequent images.

It is important that the coloured patches are representative of the colours encountered by the device,

otherwise colour calibration may not be accurate. Within photography, digital photographs are often calibrated using a Macbeth colour checker (Figure 6),³⁶ which includes 'memory colours'. Memory colours are patches of critical colours often encountered in photographs that would result in a very objectionable outcome if captured incorrectly, e.g. sky or grass.

Colour calibration in pathology

As mentioned previously, the creation of digital slides involves many stages (Figure 5), and so the process of colour calibration in digital pathology is categorized broadly into two main areas: internal and external colour calibration. Internal colour calibration involves standardization and correction of the scanning process itself, while external colour calibration is concerned with standardization of the display,



Figure 6. A Macbeth colour checker. This device is used primarily to colour-calibrate digital cameras. It includes patches for greyscale values, a range of different colour hues (saturated and pastel) and a selection of important 'memory colours', such as the colour of the sky. Source: <http://www.xrte.com/ResourceRoom/category.aspx?CategoryID=16&PartID=1257>.

accounting for the monitor's effect on perceived colour as well as the viewing environment.

External colour calibration is more straightforward, as a spectrophotometer or colorimeter (Figure 7) is used to colour-calibrate externally through measuring the appearance of colours of the display, drawing comparisons with the original image and making necessary adjustments with the use of a monitor ICC profile.

Only a couple of studies have investigated the clinical impact of external colour calibration and variation of the display characteristics on clinical performance in WSI. Krupinski *et al.*³⁹ compared a colour-calibrated monitor with one that was uncalibrated, and demonstrated that while there was no benefit in colour calibration in terms of diagnostic accuracy, there was a statistically significant improvement in time to diagnosis (mean time to diagnosis calibrated = 4.895 s versus uncalibrated = 6.304 s, $P = 0.0460$). Similarly, a small series of experiments by a vendor⁴⁰ has indicated that colour and luminance stability increases diagnostic accuracy and interpathologist agreement while decreasing reading time. However, Hanna *et al.*⁴¹ did not demonstrate an effect of display colour calibration on diagnostic accuracy.

Internal colour calibration, however, presents more difficulties. A review of the issues surrounding colour consistency in digital pathology was discussed at the Summit on Colour in Medical Imaging in May 2013,⁴² which highlighted the need for an established 'gold standard' in slide colour. It was proposed that because the optical microscope is the gold standard viewing modality, virtual slides should mimic the colour of the glass slide as viewed through the microscope. Subsequently, the FDA have recommended in their recent guidance²¹ that 'colour calibration' should be calibrated with a target slide. The test object should contain a set of measurable and representative colour patches, similar to the Macbeth colour chart used in photography, and ideally have similar spectral characteristics to stained tissue.

To meet this need, Bautista *et al.*⁴³ made a nine-patch colour calibration slide made with plastic colour filters and demonstrated a statistically significant reduction in CIELAB variation by 3.42 units between WSIs produced from different scanners of the same tissue slide. Work by Wei *et al.*⁴⁴ presented an alternative colour calibration slide for H&E stain only, with promising laboratory evaluation results. Leica Biosystems (Leica Microsystems (UK) Limited, Milton Keynes, UK) have adopted a different approach




Figure 7. Xrite i1Display Pro Colorimeter (A), a simple spectrophotometer in (B). Colorimeters, as in (A), are small devices, which use red, green and blue filters to measure the absorbance of different colours.³⁷ The colorimeter is hung in front of the screen, while colour patches are displayed on specialist software for use with the colorimeter. The colorimeter measures the colour of the patch as it is displayed and this is compared to the 'known' colour for that patch. By displaying numerous patches, a colour profile for the display can be created, taking into account the deficiencies of the display. Spectrophotometers (B),³⁸ operate slightly differently to colorimeters; they have a self-contained light source used to measure the spectral data (transmittance/ reflectance). They have a wide range of functions, including colorimetry. Source: <https://www.flickr.com/photos/seeminglee/8287570888/> https://upload.wikimedia.org/wikipedia/commons/0/0d/Spectrophotometer_small_for_8_samples-03.jpg.

through creating an ICC profile for use with their AT2 scanners using stain transmittance spectra from published literature, without scanning a colour calibration target. The effect of this colour calibration

profile can be activated or deactivated using their 'colour management' button within ImageScope® (Figure 8).

Building upon this work, our group developed a unique colour calibration test object (Figure 9) in collaboration with a vendor, FFEI Limited (Hemel Hempstead, UK). By using histochemical stains and a tissue mimicking substrate, our target is able to provide a more accurate colour representation than film-based targets with an estimated 60% reduction in colour error, and can be used with scanners from any




Figure 8. (A, B) The same case of a tubulovillous adenoma with high-grade dysplasia and pseudoinvasion as part of the UK Bowel Cancer Screening Programme using the same desktop display. A is viewed using ImageScope® from Leica Biosystems without colour management applied and B is with colour management. Some participants were concerned that the images were 'too dark', 'too blue' or 'too intensely stained', which suggests that they may not have been using colour-managed slides to complete the external quality assurance (EQA). Sources: [http://www.virtualpathology.leeds.ac.uk/eqa/specialist/nbcs/bcsp/bcspcirps.php?circ=N%20\(b15a\)](http://www.virtualpathology.leeds.ac.uk/eqa/specialist/nbcs/bcsp/bcspcirps.php?circ=N%20(b15a)).

manufacturer. Pilot work evaluating its effectiveness in the clinical setting has revealed promising results, indicating that colour calibration of virtual slides may be preferred by pathologists and may result in improved diagnostic confidence.^{45,46}

Guidance and recommendations

In digital radiology, where most images are greyscale, the need for image standardization is recognized as essential to ensure diagnostic consistency and has become a standard part of all radiology imaging workflows. There are clear guidelines regarding calibration requirements and minimum technical standards of displays used for diagnosis.^{47–49} The method used for radiological display calibration is the Digital Imaging and Communications in Medicine (DICOM) Part 14 Grey Scale Display Function (GSDF).⁵⁰ This method ensures that each shade of grey in the transition from white to grey is perceptually linear by taking into account the ability of the human visual system for light, dark and chromatic adaption, preventing two shades of grey from being too close together perceptually. Studies have indicated that calibration using DICOM Part 14 GSDF for radiology displays results in better observer performance,^{51,52} and much of the research in digital radiology is based around compliance to these standards.⁵³

As stated previously, the FDA have produced guidance stating that it is essential to control colour in digital pathology, and have recommended the use of a target slide ideally with similar spectral characteristics to stained tissue.²¹ Furthermore, guidance from the ICC White Paper 44⁵⁴ regarding displays for diagnostic digital pathology indicates that all medical grade displays should be colour-calibrated and the display should be checked for compliance every 50 days, as the displays can change over time. They also state that ambient light must be stable, as calibration of the display takes into account the ambient lighting.

Unfortunately, only a few studies have addressed this topic in digital pathology,^{39,40} and so a lack of primary research has prevented the production of guidance concerning minimum display requirements for diagnostic digital pathology to date.

The future of colour in digital pathology

It is clear that an integral part of digital pathology that has yet to be resolved is colour standardization;




Figure 9. A, A prototype colour calibration slide produced in collaboration with the University of Leeds and FFEI Limited, UK.⁴⁶ The colour patches are created using histochemical stains and a biopolymer which uptakes stains like real tissue. Each of the patches have a 'known' colour. When the colour calibration slide is scanned into a virtual slide scanner, the resultant colour patches on the whole slide image can be used to compare with the 'known' values in much the same way as a MacBeth colour checker (Figure 6). B, The effect of our unique colour calibration test object on the appearance of WSIs. The first case is a Ziehl-Neelsen stain of a lung biopsy with *Mycobacterium* with and without calibration. The second is a normal duodenal biopsy stained with haematoxylin and eosin, again with and without colour calibration.

in order to do so, further work is needed focusing upon fine-tuning colour calibration methods in relation to the effect on diagnosis. Looking to the future, we envisage several ways in which colour accuracy in pathology may be used to improve healthcare:

1. Once a global baseline for colour has been established, this will optimize diagnostic accuracy/reliability as well as facilitating worldwide collaboration. As pathologists become more skilled at using WSIs for routine work they will readjust to the

appearance of the calibrated slides, but they will still probably adapt these colours to best suit themselves and the tissue, activating personal colour profiles.

2. Secondly, digital image analysis algorithms for immunostaining will supersede human semiquantitative analysis. This will be due to improvements in algorithm reliability, facilitated by colour calibration. This might be used alongside colour normalization techniques to account for preprocessing variation in staining.

- Given that tissue staining is a shackle of light microscopy, digital pathology opens new doors with regards to pseudostaining. This involves digital superposition of colour to the WSI as opposed to staining the actual tissue. Some work has already been undertaken in this field, including an interesting paper by Kather *et al.*⁵⁵ which presented altered colours for histology based upon manipulation of colour maps to enhance perceptual contrast. Further work is needed to establish if and how such approaches might improve diagnostic quality.

The incentives to digitize pathology workflow are significant, enhancing working efficiency, global collaboration, teaching/training benefits and improved quality and patient safety. Inherent in the widespread introduction of digital pathology is appropriate colour management of WSIs. However, colour management in this field is made difficult due primarily to the absence of a 'gold standard'; future research should focus upon trying to solve this pivotal issue.

Acknowledgements

We wish to take this opportunity to thank Dr Nick Mapstone, Lead Pathologist for the North West UK Bowel Cancer Screening Programme, for the EQA participant feedback included within this review.

Conflicts of interest

We declare the following interests: D.T. is on the advisory board of Sectra and Leica/Aperio. He receives no personal remuneration for these boards. D.T. has had a collaborative research project with FFEI, where technical staff were funded by them. He received no personal remuneration. D.T. is a co-inventor on a digital pathology patent which has been assigned to Roche-Ventana on behalf of his employer. He will receive no personal remuneration.

References

- Gaucher E, Randell R, Williams B, Treanor D. The diagnostic concordance of whole slide imaging and light microscopy: a systematic review. *Arch. Pathol. Lab. Med.* 2016; doi: 10.5858/arpa.2016-0025-RA
- Collins Dictionary Online [internet] (cited 5 January 2016). Available at: <http://www.collinsdictionary.com/>.
- Physics of Color [internet] (cited 14 April 2016). Available at: <https://en.wikipedia.org/wiki/Color>.
- Sinn P, Wirtz R, Keller M *et al.* Comparison of IHC image analysis, IHC visual scoring, and quantitative real time PCR data for prediction of residual cancer burden after neoadjuvant chemotherapy. *Eur. J. Cancer* 2014; **50**: S198–S199.
- Berns R. *Billmeyer and Saltzman's principles of colour technology*. 3rd ed. Hoboken, NJ: John Wiley & Sons, 2000.
- Robertson D. The woman with superhuman vision [internet] 2014. (cited 27 October 2016). Available at: <http://www.bbc.com/future/story/20140905-the-women-with-super-human-vision>.
- Krupinski EA. Role of color in telemedicine applications (oral presentation). Silver Spring, US: International Colour Consortium Summit on Colour in Medical Imaging, 2013.
- Krupinski EA. Current perspectives in medical image perception. *Atten. Percept. Psychophys.* 2010; **72**: 1205–1217.
- Macadam DL. Visual sensitivities to color differences in daylight. *J. Opt. Soc. Am.* 1942; **32**: 247.
- Kang HR. Chromatic adaptation. In: *Computational color technology* [internet]. p. 43–56. (cited 27 October 2016). Available at: <http://ebooks.spiedigitallibrary.org/content.aspx?bookid=122§ionid=31560914>.
- Fairchild M. *Color appearance models*. 2nd ed. Hoboken, NJ: John Wiley & Sons Ltd, 2005.
- Wellbourne LE, Morland A, Wade A. Human colour perception changes between seasons. *Curr. Biol.* 2015; **25**: 646–647.
- Bartels A, Zeki S. The architecture of the human color center. *Eur. J. Neurosci.* 2000; **12**: 172–193.
- Heiting G. How Your Vision Changes as You Age [internet] (cited 14 April 2016). Available at: <http://www.allaboutvision.com/over60/vision-changes.htm>.
- Schneck M, Haegerstrom-Portnoy G, Lott L, Brabyn J. Comparison of panel D-15 tests in a large older population. *Optom. Vis. Sci.* 2014; **91**: 284–290.
- Akman O, Kosemehmetoglu K. Ocular diseases among pathologists and pathologists' perceptions on ocular diseases: a survey study. *Turk. J. Pathol.* 2015; **31**: 194–199.
- Poole CJM, Hill DJ, Christie JL, Birch J. Deficient colour vision and interpretation of histopathology slides: cross sectional study. *BMJ* 1997; **315**: 1279–1281.
- Campbell WS, Talmon GA, Foster KW, Lele SM, Kozel JA, West WW. Sixty-five thousand shades of gray: importance of color in surgical pathology diagnoses. *Hum. Pathol.* 2015; **46**: 1945–1950.
- Levenson RM, Krupinski EA, Navarro VM, Wasserman EA. Pigeons (*Columba livia*) as trainable observers of pathology and radiology breast cancer images. *PLoS One* 2015; **10**: e0141357.
- Bowel Cancer Screening Programme External Quality Assurance Feedback from Dr Nic Mapstone [internet]. (cited 27 October 2016). Available at: <http://www.gieqa.org.uk/>.
- Food and Drug Administration (FDA). *Technical performance assessment of digital pathology whole slide imaging devices; guidance for industry and food and drug administration staff*. Silver Spring, MD: FDA, 2016.
- Gray A, Wright A, Jackson P, Hale M, Treanor D. *Quality control of histochemical stains: repeatable and objective quantification using virtual slides (oral presentation)*. San Antonio, US: Pathology Visions, 2013.
- Gray A, Wright A, Jackson P, Hale M, Treanor D, James S. Quantification of histochemical stains using whole slide imaging: development of a method and demonstration of its usefulness in laboratory quality control. *J. Clin. Pathol.* 2015; **68**: 192–199.
- Schuessler Z, Delta E – color difference algorithms [internet] (cited 14 April 2016). Available at: <http://zschuessler.github.io/DeltaE/learn/>.

25. Shrestha P, Hulsken B. Colour accuracy and reproducibility in whole slide imaging scanners. *SPIE Digit. Libr.* 2014; **1**: 027501.
26. Ng CS, Palmer CR. Analysis of diagnostic confidence and diagnostic accuracy: a unified framework. *Br. J. Radiol.* 2007; **80**: 1550–1562.
27. Ali HR, Irwin M, Morris L *et al.* Astronomical algorithms for automated analysis of tissue protein expression in breast cancer. *Br. J. Cancer* 2013; **108**: 602–612.
28. Ayad E, Mansy M, Elwi D, Salem M, Salama M, Kayser K. Comparative study between quantitative digital image analysis and fluorescence *in situ* hybridization of breast cancer equivocal HER2 2+ cases. *J. Pathol. Inform.* 2015; **6**: 31.
29. Helin HO, Tuominen VJ, Ylinen O, Helin HJ, Isola J. Free digital image analysis software helps to resolve equivocal scores in HER2 immunohistochemistry. *Virchows Arch.* 2016; **468**: 191–198.
30. Holten-Rossing H, Moller Talman M-L, Kristensson M *et al.* Quantitative immunohistochemical analysis reveals association between sodium iodide symporter and estrogen receptor expression in breast cancer. *Am. J. Clin. Pathol.* 2013; **8**: 2123–2131.
31. Laurinaviciene A, Dasevicius D, Ostapenko V, Jarmalaite S, Lazutka J. Membrane connectivity estimated by digital image analysis of HER2 immunohistochemistry is concordant with visual scoring and fluorescence *in situ* hybridization results: algorithm evaluation on breast cancer tissue microarrays. *Diagn. Pathol.* 2011; **6**: 87.
32. Magee D, Treanor D, Crellin D *et al.* Colour normalisation in digital histopathology images. *Opt. Tissue Image. Anal. Microsc. Histopathol. Endosc.* 2009; 100–111.
33. Reinhard E, Ashikhmin M, Gooch B, Shirley P. Color transfer between images. *IEEE Comput. Graph. Appl.* 2001; **21**: 34–41.
34. Gavrielides MA, Gallas BD, Lenz P, Badano A, Hewitt SM. Observer variability in the interpretation of HER2/*neu* immunohistochemical expression with unaided and computer-aided digital microscopy. *Arch. Pathol. Lab. Med.* 2011; **135**: 233–242.
35. SpyderGALLERY. DATACOLOR. Lawrenceville, NJ: SpyderGALLERY, 2016.
36. McCamy C, Marcus H, Davidson J. A color-rendition chart. *J. App. Photog. Eng.* 1976; **2**: 95–99.
37. Colorimeters [internet] (cited 18 April 2016). Available at: <http://www.xrite.com/colorimeter>.
38. Spectrophotometry [internet] (cited 18 April 2016). Available at: <http://www.xrite.com/spectrophotometer>.
39. Krupinski EA, Silverstein LD, Hashimi SF, Graham AR, Weinstein RS, Roehrig H. Observer performance using virtual pathology slides: impact of LCD color reproduction accuracy. *J. Digit. Imaging* 2012; **25**: 738–743.
40. Kimpe T, Avanski A, Espig K, Rostang J, Van Hoey G, Xthona A. Influence of display characteristics on clinical performance in digital pathology. *Diagn. Pathol.* 2016; **8**: 7–8.
41. Hanna MG, Monaco SE, Ahmed I, Parwani AV, Pantanowitz L. *Impact of monitor color calibration on digital pathology interpretation*. Pittsburgh, PA: Pathology Informatics, 2015.
42. Badano A, Revie WC, Casertano A *et al.* Consistency and standardization of color in medical imaging: a consensus report. *J. Digit. Imaging* 2015; **28**: 41–52.
43. Bautista P, Hashimoto N, Yagi Y. Color standardization in whole slide imaging using a color calibration slide. *J. Pathol. Inform.* 2014; **5**: 4.
44. Wei H, Brill MH, Park T. Evaluation of targets for color calibrating digital images from an optical bright-field transmission microscope. *Colour Res. Appl.* 2014; **40**: 577–584.
45. Clarke E, Revie C, Brett D, Wilson R, Mello-Thoms C, Treanor D. Colour calibration in digital pathology: the clinical impact of a novel test object (oral presentation). 13th European Congress on Digital Pathology, Berlin. Boston, USA, 2016.
46. Clarke E, Revie C, Brett D, Wilson R, Mello-Thoms C, Treanor D. Colour calibration in digital pathology: the clinical impact of a novel test object (abstract). *J. Clin. Pathol.* 2016; **238** Suppl.: S1–S20.
47. Hirschorn DS, Krupinski EA, Flynn MJ. *IT Reference guide for the practicing radiologist*. Reston, US: American College of Radiology, 2013; 1–12.
48. Compton BK, Oosterwijk H. Requirements for medical imaging monitors (part I) O Tech Incorporated. Denton, US: OTech Incorporated. 1–13.
49. National Quality Assurance Coordinating Group for Radiography. Quality assurance guidelines for mammography including radiographic quality control. NHSBSP Publication Number 63. *Natl Qual. Assur. Coord. Gr. Radiogr.* 2006.
50. Rosslyn S. Digital imaging and communications in medicine (DICOM) Part 14: grayscale Standard display function. *Medicine (Balt.)* 2004; **10**: 3–4.
51. Krupinski E, Roehrig H. The influence of a perceptually linearized display on observer performance and visual search. *Acad. Radiol.* 2000; **7**: 8–13.
52. Johnston R, Zimmerman J, Rogers D, Pizer S. Perceptual standardization. In 3rd International Conference on Picture Archiving and Communication Systems. Proc SPIE 0536. California, 1985.
53. Butt A, Mahoney M, Savage N. The impact of computer display performance on the quality of digital radiographs: a review. *Aust. Dent. J.* 2012; **57**: 16–23.
54. International Color Consortium. White Paper no. 44 Visualization of medical content on color display systems. Reston, VA: International Color Consortium, 2016.
55. Kather JN, Weis C-A, Marx A, Schuster AK, Schad LR, Zöllner FG. New colors for histology: optimized bivariate color maps increase perceptual contrast in histological images. *PLoS One* 2015; **10**: e0145572.