

4 Multivariate Distributions

4.1 Introduction

Experimenters will often measure more than one quantity, and are often interested in the distribution of all observed quantities. As an example, a naturalist measures the height and weight of chimpanzees. They might be interested in the distribution of height-weight pairs. Since the distribution involves two quantities, we call it a *bivariate distribution*. One question might be whether or not these two quantities are independent. (We suspect they are not, since a reasonable assumption is that taller chimpanzees tend to weigh more.)

Another important application of multivariate distributions is statistical sampling. Suppose Y_1, Y_2, \dots, Y_n are n successive trials of an experiment. Statisticians are interested in the distribution of (Y_1, Y_2, \dots, Y_n) , and can use information about this distribution to infer characteristics of the experiment or the population from which the experiment sampled.

In this section, we will primarily be interested in bivariate distributions, the probability distribution of two random variables. As before, we start with the discrete case and then consider the continuous case.

4.2 Distribution of Two Discrete Random Variables

First, let's define the joint probability distribution for a pair of discrete random variables.

Joint probability distribution, discrete case

Let Y_1 and Y_2 be two discrete random variables. Then the *joint distribution* of Y_1 and Y_2 is given by the function of two variables:

$$p(y_1, y_2) = \mathbb{P}(Y_1 = y_1, Y_2 = y_2) \quad \text{for all possible pairs } (y_1, y_2)$$

Sometimes this is called the *joint probability mass function* (joint pmf). Note that $\mathbb{P}(Y_1 = y_1, Y_2 = y_2)$ means $\mathbb{P}(Y_1 = y_1 \cap Y_2 = y_2)$. This is standard notation for expressing the joint probability of two random variables.

We have already encountered one example of a bivariate distribution. Recall the distribution of the rolls of two standard, six-sided dice which we discussed several times in the section on discrete random variables. Let X_1 be the roll of the first die and X_2 the roll of the second die. Then since we have a discrete uniform distribution, the joint distribution of X_1 and X_2 is given by:

$$p(x_1, x_2) = \frac{1}{36} \quad x_1, x_2 = 1, 2, 3, 4, 5, 6$$

Just as in the case for a single discrete random variable, for a joint distribution of two discrete random variables, all the possible probabilities are nonnegative and they sum to 1.

Let Y_1 and Y_2 be discrete random variables with joint probability distribution $p(y_1, y_2)$. Then

$$0 \leq p(y_1, y_2) \leq 1 \text{ for all } y_1, y_2$$

$$\sum_{\text{all } (y_1, y_2)} p(y_1, y_2) = 1$$

where the sum is taken over all possible pairs (y_1, y_2) .

Just as in the case of a single discrete random variable, we can construct a valid joint probability distribution of two discrete random variables by assigning probabilities that add up to 1. Let Y_1 be a discrete random variable with m possible output values, and Y_2 a discrete random variable with n possible output values. Then there are mn possible joint outputs of the pair of random variables. Each of the outputs is an ordered pair of the form (y_1, y_2) . If we make an $m \times n$ table and assign probabilities to each of the mn possible joint outputs so they add up to 1, we have constructed a joint probability distribution for the two discrete random variables Y_1 and Y_2 .

Let's consider an example.

Example. Imagine we surveyed Brown undergraduates and asked them two questions.

1. Do you have an exam this week?
2. How many cups of coffee have you drunk today?

Let X_1 be the discrete random variable with values **{yes, no}** indicating whether or not a student has an exam this week. Let X_2 be the number of cups of coffee a student has drunk today. For simplicity, we will let X_2 take only the values **{0, 1, 2}** (Whether or not this is a realistic simplification is beyond the scope of this course!)

We can display the joint probability distribution for the pair (X_1, X_2) in a 2 x 3 table. There are 6 possible values for the pair (x_1, x_2) . We can choose any probabilities for the six pairs as long as they sum to 1. One possible choice is shown in the table below.

| | | X_2 | | |
|-------|-----|-------|------|------|
| | | 0 | 1 | 2 |
| X_1 | yes | 2/20 | 3/20 | 3/20 |
| | no | 6/20 | 4/20 | 2/20 |

You can verify that the six probabilities do indeed sum to 1.

4.2.1 Marginal distribution

Consider again a joint distribution (Y_1, Y_2) of two discrete random variables with pmf $p(y_1, y_2)$. (You can think of the exam-coffee example above). Y_1 and Y_2 are themselves discrete random variables. What are their distributions?

Suppose we wish to find the distribution for Y_1 by itself. We call this the *marginal distribution* of Y_1 . Essentially what we want to do is take Y_2 out of the picture entirely. How do we do that? All we have to do is sum over all the possible values of Y_2 ! The probability that $Y_1 = y_1$ is the sum of $p(y_1, y_2)$ over all possible values y_2 that Y_2 can take; this is the marginal distribution of Y_1 , and is written $p_1(y_1)$. Similarly, we can sum over all possible values Y_1 to get $p_2(y_2)$, the marginal distribution of Y_2 . This is summarized below.

Marginal distribution, discrete random variables

Let Y_1 and Y_2 be discrete random variables with joint probability distribution $p(y_1, y_2)$. Then the marginal distribution of Y_1 is given by:

$$p_1(y_1) = \sum_{\text{all } y_2} p(y_1, y_2)$$

and the marginal distribution of Y_2 is given by.

$$p_2(y_2) = \sum_{\text{all } y_1} p(y_1, y_2)$$

In both cases, we just sum the joint distribution over all the possibilities of the other random variable.

Let's return to our example above.

Example. In the exam-coffee example above, compute the marginal distributions for X_1 and X_2 .

To find the marginal distributions for each variable, we sum over all the possibilities of the other variable. If the joint distribution is presented in a two-dimensional table, this is easy. To find the marginal distribution of X_2 , we sum the values in each column. The bottom row, which we will label “total”, is the marginal distribution of X_2 . Similarly, we can find the marginal distribution for X_1 by summing each row. The rightmost column, also labeled “total”, is the marginal distribution for X_1 . In fact, the marginal distribution is called “marginal” because its values lie in the margins of the joint distribution table.

| | | X_2 | | | |
|-------|-------|-------|------|------|-------|
| | | 0 | 1 | 2 | total |
| X_1 | yes | 2/20 | 3/20 | 3/20 | 8/20 |
| | no | 6/20 | 4/20 | 2/20 | 12/20 |
| | total | 8/20 | 7/20 | 5/20 | |

You can check that the two marginal distributions sum to 1 and are thus valid probability distributions for discrete random variables.

4.2.2 Conditional distribution

Suppose again we have a joint distribution (Y_1, Y_2) of two discrete random variables with joint pmf $p(y_1, y_2)$. Another question we might ask is what is the distribution of Y_1 given that $Y_2 = y_2$. In other words, what is the conditional distribution of Y_1 given that Y_2 takes a specific value.

Let's look once more at the exam-coffee example to see how we can do this.

Example. In the exam-coffee example above, what is the distribution of the number of cups of coffee drunk today (X_2) given that a student has a midterm this week ($X_1 = \text{yes}$)?

To do this, we look at the first row of the table, which corresponds to $X_1 = \text{yes}$. This is not a valid probability mass function, because the elements do not sum to 1. But we can fix that! All we have to do is divide by the marginal probability $p_1(\text{yes}) = \mathbb{P}(X_1 = \text{yes})$, which is conveniently located just to the right in the “total” column. If we do that, we get the conditional probability for X_2 given $X_1 = \text{yes}$, which we can write as $p(y_2|\text{yes})$ or $p(y_2|Y_1 = \text{yes})$:

| y | $p(y \text{midterm})$ |
|---|-----------------------|
| 0 | 2/8 |
| 1 | 3/8 |
| 2 | 3/8 |

Now that we have seen an example, we will give the formal definition of the conditional distribution of two discrete random variables.

Conditional distribution, discrete random variables

Let Y_1 and Y_2 be discrete random variables with joint probability distribution $p(y_1, y_2)$. Let $p_2(y_2)$ be the marginal distribution of Y_2 . Then the conditional distribution of Y_1 given $Y_2 = y_2$ is:

$$p(y_1|y_2) = \mathbb{P}(Y_1 = y_1|Y_2 = y_2) = \frac{\mathbb{P}(Y_1 = y_1, Y_2 = y_2)}{\mathbb{P}(Y_2 = y_2)} = \frac{p(y_1, y_2)}{p_2(y_2)}$$

where $p_2(y_2) > 0$. In words, the conditional distribution is the joint distribution divided by the marginal distribution. We can similarly define the conditional distribution of Y_2 given $Y_1 = y_1$.

4.2.3 Independence

The final question to settle is independence. Roughly speaking, two random variables are independent if the probabilities of each one are not affected by the value of the other one. The following will serve as our definition for independence of two discrete random variables.

Independence of discrete random variables

Let Y_1 and Y_2 be discrete random variables with joint probability distribution $p(y_1, y_2)$. Let $p_1(y_1)$ and $p_2(y_2)$ be the marginal distributions of Y_1 and Y_2 . Then Y_1 and Y_2 are independent if

$$p(y_1, y_2) = p_1(y_1)p_2(y_2) \quad \text{for all } y_1, y_2$$

In other words, two random variables are independent if their joint distribution is the product of the two marginal distributions.

In the exam-coffee example above, using just about any pair of y_1 and y_2 , we can show that Y_1 and Y_2 are not independent. Did we really expect them to be independent?

4.3 Distribution of Two Discrete Continuous Variables

We will essentially repeat the same discussion for a pair of continuous random variables. Since working with continuous random variables requires integration, this will require integration in two dimensions, i.e. multivariable calculus. Since it is likely that many of you have not taken multivariable calculus, all multivariable techniques will be taught as they are needed.

4.3.1 Joint Probability Density

Recall that in the discrete case, a probability distribution was described by a probability density function (pdf). For the joint distribution of two continuous random variable, we have a joint density function, which is the continuous analogue of the joint distribution function in the discrete case.

Joint probability density, continuous case

Let Y_1 and Y_2 be two continuous random variables. Then the *joint density* of Y_1 and Y_2 is a function of two variables $f(y_1, y_2)$ with the properties that:

1. $f(y_1, y_2) \geq 0$ for all y_1, y_2 .

2.

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(y_1, y_2) dy_1 dy_2 = 1$$

In other words, $f(y_1, y_2)$ is nonnegative and integrates to 1.

In the bivariate case, instead of finding the probability that a single variable lies in an interval $[a, b]$, we find the probability that a pair of random variables lies within a region of the plane. To do this, we integrate the joint density over that region.

Probability of an event, continuous bivariate case

Let Y_1 and Y_2 be two continuous random variables with joint density $f(y_1, y_2)$. Let A be a region of the plane. Then

$$\mathbb{P}((Y_1, Y_2) \in A) = \int \int_A f(y_1, y_2) dy_1 dy_2$$

This notation may not be precise. Don't worry about it for now, we will do plenty of examples. Just remember the key idea: to find the probability that a pair of random variables lie in a region, integrate the joint density over that region.

The extra complication here is the double integral. Whereas a single integral is defined on a closed interval, a double integral is defined on a two-dimensional region of the plane. The key to success for any double integral problem is to draw the region of integration before doing anything else. Since this is so important, I will repeat it: always draw the region of integration!

We will learn how to handle this through a series of examples. We will keep coming back to this first example throughout this section.

Example. Let X and Y be random variables with joint distribution function $f(x, y)$ given by:


$$f(x, y) = \begin{cases} kxy & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

1. Find the value of k such that $f(x, y)$ is a valid joint probability density function.

As defined above (and similar to the one-dimensional case), for a joint probability density function to be valid, its integral must be 1 over the region of integration, i.e.


$$\int \int f(x, y) dx dy = 1$$

The bounds of the density function are the following: $y \geq 0, y \leq x$, and $x \leq 1$. This describes the triangular region illustrated below.



Whenever we have a double integral, we have two choices when we do our integration. We can integrate in x direction first, or we can integrate in the y direction first. Both ways give the same answer, but sometimes one is easier than the other. We will show both of them here.

Let's start by integrating in the y direction first.




Following the directions on the picture above, we integrate y first; y goes from 0 to the diagonal line $y = x$, so those are the limits for the integral with respect to y (inner integral). Note that the upper limit is a function of x . Each integral in y is a vertical "slice" of our region. Then we integrate with respect to x . Imagining this as stacking our vertical slices side-by-side in the horizontal direction, x goes from 0 to 1, so those are the limits for the integral with respect to x (outer integral). Putting this together,

we have:

$$\begin{aligned}
 1 &= \int_0^1 \int_0^x kxy \, dy \, dx \\
 &= k \int_0^1 x \frac{y^2}{2} \Big|_0^x \, dx \\
 &= \frac{k}{2} \int_0^1 x^3 \, dx \\
 &= \frac{k}{2} \frac{x^4}{4} \Big|_0^1 = \frac{k}{8}
 \end{aligned}$$

Multiplying by 8 gives us $k = 8$.

Let's do the integral the other way and verify to see if we get the same result. This time we integrate in the x direction first.



Once again following the directions on the picture above, we integrate x first; x starts at the diagonal line $y = x$ and goes to $x = 1$. The diagonal line has the equation $y = x$, which we solve for x to get $x = y$. Thus the lower limit is the function $x = y$. The upper limit is 1, so the limits of the integral with respect to x (inner integral) are y and 1. Each integral in x is a horizontal "slice" of our region. Then we integrate with respect to y . Imagining this as stacking our horizontal slices one on top of the other in the vertical direction, y goes from 0 to 1, so those are the limits for the integral with


respect to y (outer integral). Putting this together, we have:

$$\begin{aligned}
 1 &= \int_0^1 \int_y^1 kxy \, dx dy \\
 &= k \int_0^1 y \frac{x^2}{2} \Big|_y^1 dy \\
 &= \frac{k}{2} \int_0^1 y(1 - y^2) dy \\
 &= \frac{k}{2} \int_0^1 (y - y^3) dy \\
 &= \frac{k}{2} \left(\frac{y^2}{2} - \frac{y^4}{4} \right) \Big|_0^1 = \frac{k}{8}
 \end{aligned}$$

We get the same answer! Which one was easier?

2. Find $\mathbb{P}(X < 0.6 \cap Y > 0.2)$.

Here we are finding the probability that the pair (X, Y) falls in a specific region of the plane. The first step (as always) is to draw the region.



From this picture, we get the limits of integration. Let's integrate in the y direction first. Recall that we found that $k = 8$ above. This gives us the integral:

$$\mathbb{P}(X < 0.6 \cap Y > 0.2) = \int_{0.2}^{0.6} \int_{0.2}^x 8xy \, dy dx$$

This integral can be evaluated like the one above. The most important thing to know is how to set the problem up with the correct limits, but for completeness we will do

the computation below.

$$\begin{aligned}
\mathbb{P}(X < 0.6 \cap Y > 0.2) &= \int_{0.2}^{0.6} \int_{0.2}^x 8xy \, dy dx \\
&= 8 \int_{0.2}^{0.6} x \frac{y^2}{2} \Big|_{0.2}^x dx \\
&= 4 \int_{0.2}^{0.6} (x^3 - 0.04x) dx \\
&= 4 \left(\frac{x^4}{4} - 0.04 \frac{x^2}{2} \right) \Big|_{0.2}^{0.6} \\
&= (0.6^4 - 0.2^4) - 0.08 (0.6^2 - 0.2^2) \\
&= 0.1024
\end{aligned}$$

We could also have integrated in the x direction first.

4.3.2 Marginal Distribution

Consider a joint distribution (Y_1, Y_2) of two continuous random variables with joint density $f(y_1, y_2)$. Y_1 and Y_2 are themselves continuous random variables, and their distributions are called the *marginal distributions* of Y_1 and Y_2 . The probability densities of Y_1 and Y_2 are the *marginal densities* of Y_1 and Y_2 .

How do we find the marginal densities? Recall that for the discrete case, we summed over the other variable, e.g. to find the marginal density of Y_1 , we summed the joint distribution over all the values Y_2 can take. Here we do the exact same thing, except we replace summation with integration. To get the marginal density of Y_1 , we integrate the joint density $f(y_1, y_2)$ over y_2 . The marginal density of Y_1 is written $f_1(y_1)$, and is a function of y_1 alone. (If after the integration you still have terms involving y_2 , something went wrong.) Similarly, we can find the marginal density of Y_2 .

Marginal density, continuous random variables

Let Y_1 and Y_2 be continuous random variables with joint density $f(y_1, y_2)$. Then the marginal density of Y_1 is given by:

$$f_1(y_1) = \int f(y_1, y_2) dy_2$$

and the marginal density of Y_2 is given by.

$$f_2(y_2) = \int f(y_1, y_2) dy_1$$

In both cases, we integrate the joint density over the other random variable to remove it from the picture entirely. Sometimes we call this “integrating out” the other random variable.

Let's return to the first example from the joint density section.

Example. Let X and Y be random variables with joint distribution function $f(x, y)$ given by:

$$f(x, y) = \begin{cases} 8xy & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

1. Find the marginal densities for X and Y .

Since our random variable are X and Y , we will denote the two marginal densities by $f_X(x)$ and $f_Y(y)$. First, we find the marginal density for X by integrating over y . Since we are integrating only in a single variable, there is only one choice for limits. Refer back to the picture of the region above. To integrate in y , we start at $y = 0$ and integrate until we reach the line $y = x$. Thus the limits of integration are 0 and x .

$$\begin{aligned} f_X(x) &= \int_0^x 8xy \, dy \\ &= 8x \frac{y^2}{2} \Big|_0^x \\ &= 4x^3 \end{aligned}$$

With Y out of the picture, the random variable X is free to take values from 0 to 1. (The marginal distribution is one-dimensional, so there are no pesky regions to deal with!) The above expression for the marginal density is only valid for $0 \leq x \leq 1$. Outside that range, the marginal density is 0. Thus we write the marginal density $f_X(x)$ as:

$$f_X(x) = \begin{cases} 4x^3 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

It is important that we write the marginal density in this way so that we know that the X is 0 outside $[0, 1]$. Since the marginal density is a valid probability density, you can check that it does in fact integrate to 1. Also note that the marginal density for X is a function of x alone; y does not appear anywhere since we integrated it out!

Now we find the marginal density for Y by integrating over x . The limits for x are $x = 0$ and $x = y$. (See the picture of the region above if this is not clear.)

$$\begin{aligned} f_Y(y) &= \int_y^1 8xy \, dx \\ &= 8y \frac{x^2}{2} \Big|_y^1 \\ &= 4y(1 - y^2) \end{aligned}$$

With X out of the picture, the random variable Y can take values from 0 to 1, so we write the marginal density of Y as

$$f_Y(y) = \begin{cases} 4y(1 - y^2) & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

2. Find the expected values for X and Y .

How do we find the expected values of X and Y . We use the marginal densities and do the same thing we do every night, Pinky - try to take over the world! The marginal densities are just standard probability densities for continuous random variables; thus we find the expected values of X and Y using the marginal densities and the formula for the expected value of a continuous random variable.

$$\begin{aligned} \mathbb{E}(X) &= \int_{-\infty}^{\infty} x f_X(x) dx \\ \mathbb{E}(Y) &= \int_{-\infty}^{\infty} y f_Y(y) dy \end{aligned}$$

First we find the expected value for X . Notice that the limits of integration become 0 and 1, since that is the region where the marginal density is nonzero.

$$\begin{aligned} \mathbb{E}(X) &= \int_{-\infty}^{\infty} x f_X(x) dx \\ &= \int_0^1 x f_X(x) dx \\ &= \int_0^1 x 4x^3 dx \\ &= 4 \int_0^1 x^4 dx \\ &= 4/5 \end{aligned}$$

Similarly, we find the expected value of Y .

$$\begin{aligned} \mathbb{E}(Y) &= \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= \int_0^1 y f_Y(y) dy \\ &= \int_0^1 y 4y(1 - y^2) dy \\ &= 4 \int_0^1 (y^2 - y^4) dy \\ &= 8/15 \end{aligned}$$

4.3.3 Conditional Distribution

Just as in the discrete case, we can talk about conditional distributions. In the continuous case, we will have a continuous density function. Suppose we have a joint distribution (Y_1, Y_2) of two continuous random variables with conditional density $f(y_1, y_2)$. We might be interested in the distribution of Y_1 given that $Y_2 = y_2$. Since Y_1 is a continuous random variable, we will express this as a conditional probability density.

Conditional distribution, continuous random variables

Let Y_1 and Y_2 be continuous random variables with joint probability density $f(y_1, y_2)$. Let $f_2(y_2)$ be the marginal density of Y_2 . Then the conditional density of Y_1 given $Y_2 = y_2$ is:

$$f(y_1|y_2) = \frac{f(y_1, y_2)}{f_2(y_2)}$$

where $f_2(y_2) > 0$. As in the discrete case, the conditional density is the joint density divided by the marginal density. We can similarly define the conditional density of Y_2 given $Y_1 = y_1$.

Note that the conditional density can (and often will) depend on the value of the random variable we are conditioning on. For example, $f(y_1|y_2)$ may depend on the value y_2 . Furthermore, the range of values y_1 can take may also depend on y_2 (we will see this in the example below). Contrast this to the marginal density, where we eliminate the other random variable entirely.

Once again, let's return to our example.

Example. Let X and Y be random variables with joint distribution function $f(x, y)$ given by:

$$f(x, y) = \begin{cases} 8xy & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

1. Find the conditional density of X given $Y = y$.

To find the conditional density $f(x|y)$, we divide the joint density $f(x, y)$ by the marginal density $f_Y(y)$. We found the marginal density above, so we have:

$$f(x|y) = \frac{f(x, y)}{f_Y(y)} = \frac{8xy}{4y(1 - y^2)} = \frac{2x}{1 - y^2}$$

Note that this density does in fact depend on the value of y we are conditioning on. However, we are not done. The bounds on the conditional density are important and must be specified. Referring back to the picture of the region, we note that if $Y = y$,

then X can only range from the diagonal line $y = x$ to 1, i.e. X must be between y and 1. Thus the conditional density of X given $Y = y$ is given by:

$$f(x|y) = \begin{cases} \frac{2x}{1-y^2} & y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

2. Find the conditional density of Y given $X = x$.

Again, we divide the joint density by the marginal density, where we computed the marginal density $f_X(x)$ above.

$$f(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{8xy}{4x^3} = \frac{2y}{x^2}$$

As with the other conditional density, this depends on x . The bounds of the conditional density will also depend on x . Looking at the picture of the region, if $X = x$, then y can only range from 0 to the diagonal line $y = x$, i.e. Y must be between 0 and x . Thus the conditional density is:

$$f(y|x) = \begin{cases} \frac{2y}{x^2} & 0 \leq y \leq x \\ 0 & \text{otherwise} \end{cases}$$

3. Find the expected value of X given $Y = y$.

A conditional density is just a probability density of a continuous random variable, so we can find its expected value using the standard formula for the expected value of a continuous random variable.

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} xf(x|y)dx \\ &= \int_y^1 x \frac{2x}{1-y^2} dx \\ &= \frac{2}{1-y^2} \int_y^1 x^2 \\ &= \frac{2}{1-y^2} \left. \frac{x^3}{3} \right|_y^1 \\ &= \frac{2(1-y^3)}{3(1-y^2)} \end{aligned}$$

Note that we used the bounds on the conditional density in the second line above. Unsurprisingly, this depends on y .

4.3.4 Independence

We have a similar definition for independence in the case of continuous random variables.

Independence of continuous random variables

Let Y_1 and Y_2 be continuous random variables with joint density $f(y_1, y_2)$. Let $f_1(y_1)$ and $f_2(y_2)$ be the marginal densities of Y_1 and Y_2 . Then Y_1 and Y_2 are independent if

$$f(y_1, y_2) = f_1(y_1)f_2(y_2) \quad \text{for all } y_1, y_2$$

In other words, two continuous random variables are independent if their joint density is the product of the two marginal densities.

In the example we have been looking at for the past three sections, the joint density is not the product of the marginal densities, so X and Y are not independent.

4.3.5 Another Example


Bivariate densities are important enough that we will provide more examples problems involving them.

Example. Let X and Y be random variables with joint distribution function $f(x, y)$ given by:

$$f(x, y) = \begin{cases} cx & 0 \leq x \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

1. Find the value of c such that $f(x, y)$ is a valid joint probability density function.

The first step is always to draw the region.



To find the value of c , we integrate the joint density over the region and set the result equal to 1. We will integrate in the x direction first since there's already an x in the integrand and integrating in the x direction first allows both lower limits to be 0 (the other choice would work fine too):

$$\begin{aligned}
 1 &= \int_0^1 \int_0^y c x dx dy \\
 &= c \int_0^1 \frac{x^2}{2} \Big|_0^y dy \\
 &= c \int_0^1 \frac{y^2}{2} dy \\
 &= c \frac{y^3}{6} \Big|_0^1 \\
 &= \frac{c}{6}
 \end{aligned}$$

Thus we find that $c = 6$ for this to be a valid joint probability density function.

2. Find the marginal densities for X and Y .

To do this, we integrate out each random variable in turn. Be careful to get the correct limits of integration (refer back to the picture of the region early and often.) For the marginal density of X we first integrate out y .

$$\begin{aligned}
 f_X(x) &= \int_x^1 6x dy \\
 &= 6xy \Big|_x^1 \\
 &= 6x(1 - x)
 \end{aligned}$$

With Y removed from the picture, X can freely range from 0 to 1, so the marginal density of X is:

$$f_X(x) = \begin{cases} 6x(1 - x) & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

For the marginal density of Y , we first integrate out x :

$$\begin{aligned}
 f_Y(y) &= \int_0^y 6x dx \\
 &= 3x^2 \Big|_0^y \\
 &= 3y^2
 \end{aligned}$$

Then, with X removed from the picture, Y can freely range from 0 to 1, so the marginal density of Y is:

$$f_Y(y) = \begin{cases} 3y^2 & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

It is important that you give the correct bounds for the marginal densities.

3. What is the expected value of X ?

To find $\mathbb{E}(X)$, we use the formula for expected value of a continuous random variable with the marginal density for X .

$$\begin{aligned} \mathbb{E}(X) &= \int_{-\infty}^{\infty} f_X(x) dx \\ &= \int_0^1 x 6x(1-x) dx \\ &= 6 \int_0^1 (x^2 - x^3) dx \\ &= 6 \left(\frac{x^3}{3} - \frac{x^4}{4} \right) \Big|_0^1 \\ &= 6 \left(\frac{1}{3} - \frac{1}{4} \right) \\ &= \frac{1}{2} \end{aligned}$$

Similarly we can find the expected value of Y using the marginal density for Y .

4. What is the conditional density for X given $Y = y$?

To get the conditional density $f(x|y)$, we first divide the joint density by the marginal density of Y :

$$f(x|y) = \frac{f(x, y)}{f_Y(y)} = \frac{6x}{3y^2} = \frac{2x}{y^2}$$

What are the bounds on the conditional density? Refer back to the picture of the region above. Given that $Y = y$, X can range from 0 to y , thus the conditional density is:

$$f(x|y) = \begin{cases} \frac{2x}{y^2} & 0 \leq x \leq y \\ 0 & \text{otherwise} \end{cases}$$

5. What is the expected value of X given $Y = y$?

Here, we use the formula for the expected value of a continuous random variable with the conditional density $f(x|y)$.

$$\begin{aligned}
 \mathbb{E}(X|Y = y) &= \int_{-\infty}^{\infty} f(x|y)dx \\
 &= \int_0^y x \frac{2x}{y^2} dx \\
 &= \int_0^y \frac{2x^2}{y^2} dx \\
 &= \frac{2x^3}{3y^2} \Big|_0^y \\
 &= \frac{2y}{3}
 \end{aligned}$$