

# A REFORMULATED KREIN MATRIX FOR STAR-EVEN POLYNOMIAL OPERATORS WITH APPLICATIONS\*

TODD KAPITULA<sup>†</sup>, ROSS PARKER<sup>‡</sup>, AND BJÖRN SANDSTEDÉ<sup>§</sup>

**Abstract.** In its original formulation the Krein matrix was used to locate the spectrum of first-order star-even polynomial operators where both operator coefficients are nonsingular. Such operators naturally arise when considering first-order-in-time Hamiltonian PDEs. Herein the matrix is reformulated to allow for operator coefficients with nontrivial kernel. Moreover, it is extended to allow for the study of the spectral problem associated with quadratic star-even operators, which arise when considering the spectral problem associated with second-order-in-time Hamiltonian PDEs. In conjunction with the Hamiltonian-Krein index (HKI) the Krein matrix is used to study two problems: conditions leading to Hamiltonian-Hopf bifurcations for small spatially periodic waves, and the location and Krein signature of small eigenvalues associated with, e.g.,  $n$ -pulse problems. For the first case we consider in detail a first-order-in-time fifth-order KdV-like equation. In the latter case we use a combination of Lin's method, the HKI, and the Krein matrix to study the spectrum associated with  $n$ -pulses for a second-order-in-time Hamiltonian system which is used to model the dynamics of a suspension bridge.

**Key words.** Krein matrix, star-even operators,  $n$ -pulses

**AMS subject classifications.** 35P30, 47A55, 47A56, 70H14

**1. Introduction.** Consider the star-even operator polynomial,

$$\mathcal{P}_n(\lambda) := \sum_{j=0}^n \lambda^j \mathcal{A}_j.$$

On some Hilbert space,  $X$ , endowed with inner-product,  $\langle \cdot, \cdot \rangle$ , we assume the operator coefficients  $\mathcal{A}_{2\ell}$  are Hermitian,  $\mathcal{A}_{2\ell}^a = \mathcal{A}_{2\ell}$ , and the operator coefficients  $\mathcal{A}_{2\ell+1}$  are skew-Hermitian,  $\mathcal{A}_{2\ell+1}^a = -\mathcal{A}_{2\ell+1}$ . Here we let  $\mathcal{T}^a$  denote the adjoint of the operator  $\mathcal{T}$ . If  $n = 1$ ,

$$\mathcal{P}_1(\lambda)\psi = 0 \rightsquigarrow (\mathcal{A}_0 + \lambda\mathcal{A}_1)\psi = 0.$$

Assuming  $\mathcal{A}_1$  is invertible, this spectral problem is equivalent to,

$$\mathcal{A}_1^{-1}\mathcal{A}_0\psi = \gamma\psi, \quad \gamma = -\lambda,$$

which, since  $\mathcal{A}_1^{-1}$  is skew-Hermitian and  $\mathcal{A}_0$  is Hermitian, is the canonical form for a Hamiltonian eigenvalue problem. Indeed, while we will not go into the details here, it is possible via a change of variables to put any star-even problem into canonical form, see [15, Section 3] and the references therein. For our purposes it is best to leave the problem in its original formulation.

Values  $\lambda_0$  for which the polynomial  $\mathcal{P}_n(\lambda_0)$  is singular will be called *polynomial eigenvalues*. Because of these assumed coefficient properties, the polynomial eigenvalues are symmetric with respect to the imaginary axis of the complex plane. The

---

\*Submitted to the editors 01/30/2019.

**Funding:** TK was partially funded by the NSF under DMS-1108783. RP was partially funded by the NSF under DMS-1148284. BS was partially funded by the NSF under DMS-1714429.

<sup>†</sup>Department of Mathematics and Statistics, Calvin University, Grand Rapids, MI 49546 (tmk5@calvin.edu).

<sup>‡</sup>Division of Applied Mathematics, Brown University, Providence, RI 02912 (ross.parker@brown.edu).

<sup>§</sup>Division of Applied Mathematics, Brown University, Providence, RI 02912 (bjorn.sandstede@brown.edu).

eigenvalue symmetry follows from,

$$\mathcal{P}_n(\lambda)^a = \mathcal{P}_n(-\bar{\lambda}),$$

so  $\lambda$  being a polynomial eigenvalue implies  $-\bar{\lambda}$  is also a polynomial eigenvalue. In order to ensure there are no polynomial eigenvalues at infinity, we assume  $\mathcal{A}_n$  is invertible.

More can be said about the set of polynomial eigenvalues under compactness assumptions (which will henceforth be assumed, except for the example considered in [section 6](#)). Suppose the Hermitian operator  $\mathcal{A}_0$  has compact resolvent, so the eigenvalues for this operator coefficient are real, semi-simple, and have finite multiplicity. Let  $P_{\mathcal{A}_0} : X \mapsto \ker(\mathcal{A}_0)$  be the orthogonal projection, and set  $P_{\mathcal{A}_0}^\perp = \mathcal{I} - P_{\mathcal{A}_0} : X \mapsto \ker(\mathcal{A}_0)^\perp$ . Assuming the operators,

$$(P_{\mathcal{A}_0}^\perp \mathcal{A}_0 P_{\mathcal{A}_0}^\perp)^{-1} P_{\mathcal{A}_0}^\perp \mathcal{A}_j P_{\mathcal{A}_0}^\perp : \ker(\mathcal{A}_0)^\perp \mapsto \ker(\mathcal{A}_0)^\perp, \quad j = 1, \dots, n,$$

are compact, the spectrum for  $\mathcal{P}_n(\lambda)$  is point spectra only [\[2, Remark 2.2\]](#). Moreover, each polynomial eigenvalue has finite multiplicity, and infinity is the only possible accumulation point for the polynomial eigenvalues.

Regarding the number of unstable polynomial eigenvalues, i.e., those polynomial eigenvalues with positive real part, the total number can be bounded above via the Hamiltonian-Krein index (HKI). Let  $k_r$  denote the total number (counting multiplicity) of real and positive polynomial eigenvalues, and let  $k_c$  be the total number (counting multiplicity) of polynomial eigenvalues with positive real part and nonzero imaginary part. The total number of unstable polynomial eigenvalues is  $k_r + k_c$ .

The HKI also takes into account a subset of purely imaginary polynomial eigenvalues; namely, those with negative Krein signature. For each purely imaginary and nonzero eigenvalue,  $i\lambda_0$  with  $\lambda_0 \in \mathbb{R}$ , with associated eigenspace  $\mathbb{E}_{i\lambda_0}$ , set

$$k_i^-(i\lambda_0) = n(-\lambda_0 [iP'_n(i\lambda_0)]|_{\mathbb{E}_{i\lambda_0}}).$$

Here  $n(\mathbf{S})$  denotes the number of negative eigenvalues for the Hermitian matrix  $\mathbf{S}$ , and  $iP'_n(i\lambda_0)|_{\mathbb{E}_{i\lambda_0}}$  is the Hermitian matrix formed by the representation of the Hermitian operator  $-i\lambda_0 P'_n(i\lambda_0)$  restricted to the eigenspace  $\mathbb{E}_{i\lambda_0}$ . If the polynomial eigenvalue is simple with associated eigenvector  $u_{i\lambda_0}$ , then

$$k_i^-(i\lambda_0) = n(\lambda_0 \langle -iP'_n(i\lambda_0)u_{i\lambda_0}, u_{i\lambda_0} \rangle);$$

in particular, if  $n = 1$  then it takes the more familiar form,

$$k_i^-(i\lambda_0) = n(\langle \mathcal{A}_0 u_{i\lambda_0}, u_{i\lambda_0} \rangle).$$

The nonnegative integer  $k_i^-(i\lambda_0)$  is the negative Krein index associated with the purely imaginary eigenvalue. If  $k_i^-(i\lambda_0) = 0$ , the polynomial eigenvalue is said to have positive Krein signature; otherwise, it has negative Krein signature. The total negative Krein index is the sum of the individual Krein indices,

$$k_i^- = \sum k_i^-(i\lambda_0).$$

Regarding  $k_i^-$ , consider the collision of two simple polynomial eigenvalues on the imaginary axis. If they both have the same signature, then after the collision they will each remain purely imaginary. On the other hand, if they have opposite Krein

signature, then it will generically be the case that after the collision the pair will have nonzero real part, which due to the spectral symmetry means that one of the polynomial eigenvalues will have positive real part. This is the so-called Hamiltonian-Hopf bifurcation. In the case of  $n = 1$  the interested reader should consult [20, Chapter 7.1] for more details regarding the case of the collision of two simple polynomial eigenvalues, and [14, 37] for the case of higher-order collisions. Note that if  $k_i^- = 0$ , then no polynomial eigenvalues will leave the imaginary axis.

The HKI is defined to be the sum of the three indices,

$$K_{\text{Ham}} = k_r + k_c + k_i^-.$$

The HKI is intimately related to the operator coefficients. If  $X = \mathbb{C}^N$ , i.e., the operator is actually a star-even matrix polynomial with  $nN$  polynomial eigenvalues, then under the assumption  $\mathcal{A}_0, \mathcal{A}_n$  are nonsingular,

$$K_{\text{Ham}} = \begin{cases} n(\mathcal{A}_0) + (\ell - 1)N, & n = 2\ell - 1 \\ n(\mathcal{A}_0) + n((-1)^{\ell-1} \mathcal{A}_n) + (\ell - 1)N, & n = 2\ell \end{cases}$$

[15, Theorem 3.4]. If  $n \geq 3$  the upper bound for the total number of unstable polynomial eigenvalues depends upon the dimension of the space; consequently, taking the limit  $N \rightarrow +\infty$  provides no meaningful information regarding the limiting case of operator coefficients which are compact operators. Consequently, we henceforth assume  $n \in \{1, 2\}$ .

Now, suppose that the highest-order coefficient is nonsingular, but  $\mathcal{A}_0$  has a nontrivial kernel. If  $n = 1$ , then under the widely applicable assumptions,

- (a)  $\mathcal{A}_1 : \ker(\mathcal{A}_0) \mapsto \ker(\mathcal{A}_0)^\perp$
- (b)  $\mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1|_{\ker(\mathcal{A}_0)}$  is invertible,

we know,

$$(1.1) \quad K_{\text{Ham}} = n(\mathcal{A}_0) - n(-\mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1|_{\ker(\mathcal{A}_0)}),$$

see [12, 27] and the references therein. Regarding the operator  $\mathcal{A}_1$ , the case where

- (a) there is a nontrivial kernel, but where the rest of spectrum is otherwise uniformly bounded away from the origin, is covered in [7, 19] and [20, Chapter 5.3]
- (b) where the spectrum which is not bounded away from the origin is considered in [21, 28].

If  $n = 2$ , then upon replacing condition (b) above with,

- (b)  $(\mathcal{A}_2 - \mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1)|_{\ker(\mathcal{A}_0)}$  is invertible,

we know,

$$(1.2) \quad K_{\text{Ham}} = n(\mathcal{A}_0) + n(\mathcal{A}_2) - n([\mathcal{A}_2 - \mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1]|_{\ker(\mathcal{A}_0)}),$$

see [2].

The goal of this paper is to construct a square matrix-valued function, say  $\mathbf{K}(\lambda)$ , which has the properties that for  $\lambda \in i\mathbb{R}$ ,

- (a)  $\mathbf{K}(\lambda)$  is Hermitian and meromorphic
- (b)  $\det \mathbf{K}(\lambda) = 0$  only if  $\lambda$  is a polynomial eigenvalue
- (c)  $\mathbf{K}(\lambda)$  can be used to determine the Krein signature of a polynomial eigenvalue.

The matrix  $\mathbf{K}(\lambda)$  is known as the Krein matrix. The properties (a) and (b) listed above are reminiscent of those possessed by the Evans matrix, except that the Evans matrix is analytic [20, Chapters 8-10]. Regarding (b) and (c), since the determinant of a matrix is equal to the product of its eigenvalues, property (b) is satisfied if at least one of the eigenvalues is zero. Henceforth, we will call the eigenvalues of the Krein matrix, say  $r_j(\lambda)$ , the Krein eigenvalues. The determination of the Krein signature of a purely imaginary polynomial eigenvalue takes place through the Krein eigenvalues. If  $r_j(\lambda_0) = 0$  for some  $\lambda_0 \in i\mathbb{R}$ , the Krein signature is found by considering the sign of  $r'_j(\lambda_0)$ . Thus, via a plot of the Krein eigenvalues one can graphically determine the signature of a purely imaginary polynomial eigenvalue through the slope of the curve at a zero. The interested reader should consult [24] for a graphical connection between the Krein signature and an extended Evans function.

The Krein matrix was first constructed for linear polynomials of the canonical form,

$$\mathcal{P}_1(\lambda) = \begin{pmatrix} \mathcal{L}_+ & 0 \\ 0 & \mathcal{L}_- \end{pmatrix} + \lambda \begin{pmatrix} 0 & \mathcal{I} \\ -\mathcal{I} & 0 \end{pmatrix},$$

where  $\mathcal{L}_{\pm}$  are invertible Hermitian operators with compact resolvent, and  $\mathcal{I}$  denotes the identity operator, see [14, 25]. Recent applications of the Krein matrix include a new proof of the Jones-Grillakis instability criterion,

$$k_r \geq |\mathbf{n}(\mathcal{L}_-) - \mathbf{n}(\mathcal{L}_+)|,$$

as well as a study of the spectral problem for waves to a mathematical model for Bose-Einstein condensates [18, 19].

The paper is organized as follows. In section 2 the Krein matrix is constructed for star-even polynomial operators of any degree. In particular, the previous invertibility assumption on  $\mathcal{A}_0$  is removed. In section 3 the properties of the Krein eigenvalues are deduced; in particular, their relation to the Krein signature of purely imaginary polynomial eigenvalues is given. In section 4 the Krein eigenvalues are used to study the Hamiltonian-Hopf bifurcation problem associated with small periodic waves. While the underlying wave is small, it is possible for the polynomial eigenvalues to have  $\mathcal{O}(1)$  imaginary part (see [8, 23, 35] for a similar study using a different approach). In section 5 we show how the Krein matrix can be used to locate small eigenvalues which arise from some type of bifurcation. However, the analysis does not use perturbation theory, so it is consequently possible to use the resulting Krein matrix to consider spectral stability for multi-pulse problems, where the small eigenvalues arise from the exponentially small tail-tail interactions of a translated base pulse. Finally, in section 6 we use the Krein matrix to study the spectral problem associated with  $n$ -pulse solutions to the suspension bridge equation, which is a second-order-in-time Hamiltonian PDE.

**Acknowledgements.** The authors would like to thank the referees for their careful reading of the original manuscript, and their helpful suggestions and constructive critique. We believe that this revision is a substantial improvement over the original because of their work.

**2. The Krein matrix.** The Krein matrix allows us to reduce the infinite-dimensional eigenvalue problem,

$$\mathcal{P}_n(\lambda)\psi = 0,$$

to a finite-dimensional problem,

$$\mathbf{K}_S(\lambda)\mathbf{x} = \mathbf{0}.$$

Here  $\mathbf{K}_S(\lambda)$  is the (square) Krein matrix. Whereas the original star-even operator is analytic in the spectral parameter, the Krein matrix is meromorphic with poles on the imaginary axis. The presence of these poles is the key to using the Krein matrix to determine the Krein signature of a purely imaginary eigenvalue.

**2.1. General construction.** Let  $S \subset X$  be a finite-dimensional subspace of dimension  $n_S$  with orthonormal basis  $\{s_j\}$ , and let  $P_S : X \mapsto X$  be the orthogonal projection, i.e.,

$$P_S u = \sum_{j=1}^{n_S} \langle u, s_j \rangle s_j.$$

Denote the complementary orthogonal projection as

$$P_{S^\perp} := \mathcal{I} - P_S,$$

and write

$$u = s + s^\perp, \text{ with } P_S u = s, \ P_{S^\perp} u = s^\perp.$$

In constructing the subspace-dependent Krein matrix,  $\mathbf{K}_S(\lambda)$ , for the polynomial eigenvalue problem, we will extensively use the orthogonal projections. We first rewrite the polynomial eigenvalue problem,

$$(2.1) \quad \mathcal{P}_n(\lambda)s + \mathcal{P}_n(\lambda)s^\perp = 0.$$

Applying the complementary projection to (2.1) yields

$$(2.2) \quad P_{S^\perp} \mathcal{P}_n(\lambda)s + P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp} s^\perp = 0.$$

The operator  $P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp} : S^\perp \mapsto S^\perp$  is a star-even polynomial operator. Consequently, it has the same spectral properties as the original star-even operator; in particular, it is invertible except for a countable number of spectral values. If  $\lambda$  is not a polynomial eigenvalue for the operator  $P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp}$ , then we can invert to write

$$s^\perp = -(P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(\lambda)s,$$

which leads to,

$$(2.3) \quad s^\perp = P_{S^\perp} s^\perp = -P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(\lambda)s.$$

If we take the inner-product of (2.1) with a basis element  $s_j$ , we get

$$\langle s_j, \mathcal{P}_n(\lambda)s \rangle + \langle s_j, \mathcal{P}_n(\lambda)s^\perp \rangle = 0.$$

Substitution of the expression in (2.3) into the above provides,

$$\langle s_j, \mathcal{P}_n(\lambda)s \rangle - \langle s_j, \mathcal{P}_n(\lambda)P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(\lambda)P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(\lambda)s \rangle = 0.$$

Writing

$$s = \sum_{j=1}^{n_S} x_j s_j,$$

the above expression becomes

$$(2.4) \quad \mathbf{K}_S(\lambda)\mathbf{x} = \mathbf{0},$$

where the Krein matrix  $\mathbf{K}_S(\lambda) \in \mathbb{C}^{n_S \times n_S}$  has the form

$$\mathbf{K}_S(\lambda) = \mathcal{P}_n(\lambda)|_S - \mathcal{P}_n(\lambda)P_{S^\perp}(P_{S^\perp}\mathcal{P}_n(\lambda)P_{S^\perp})^{-1}P_{S^\perp}\mathcal{P}_n(\lambda)|_S,$$

where we use the notation

$$(\mathcal{T}|_S)_{ij} = \langle s_i, \mathcal{T}s_j \rangle.$$

In conclusion, polynomial eigenvalues for the original problem are found via solving (2.4), which means

$$\det \mathbf{K}_S(\lambda) = 0, \quad \text{or} \quad \mathbf{x} = \mathbf{0}.$$

What does it mean if  $\lambda_0$  is a polynomial eigenvalue with  $\mathbf{x} = \mathbf{0}$ ? In this case the associated eigenfunction for the polynomial eigenvalue,  $u_0$ , satisfies

$$P_S u_0 = 0, \quad P_{S^\perp} u_0 = u_0.$$

Going back to (2.1) and (2.2) we see

$$P_n(\lambda_0)P_{S^\perp}u_0 = 0 \quad \rightsquigarrow \quad P_{S^\perp}P_n(\lambda_0)P_{S^\perp}u_0 = 0.$$

In other words,  $\lambda_0$  is also a polynomial eigenvalue for the operator  $P_{S^\perp}P_n(\lambda_0)P_{S^\perp}$ . Thus, if  $\lambda_0$  is a polynomial eigenvalue for which the associated eigenfunction resides in  $S^\perp$ ,  $\lambda_0$  is also a pole for the Krein matrix. Consequently, we cannot expect to capture such polynomial eigenvalues by solving  $\det \mathbf{K}_S(\lambda) = 0$ . This fact will motivate our later choice for the subspace  $S$ , as we need to know that the polynomial eigenvalues being missed by considering the zero set of the determinant of the Krein matrix are somehow unimportant.

The choice of the subspace is determined by looking at the Krein index of a purely imaginary polynomial eigenvalue,  $\lambda = i\lambda_0$ . Letting  $\mathbb{E}_{i\lambda_0}$  denote the generalized eigenspace, the negative Krein index is

$$k_i^-(i\lambda_0) := n(-\lambda_0[iP_n'(i\lambda_0)]|_{\mathbb{E}_{i\lambda_0}})$$

(see [2]). Since the goal is to have the Krein matrix capture all possible polynomial eigenvalues with negative Krein index through its determinant, we then want it to be the case that if  $i\lambda_0$  is a polynomial eigenvalue whose associated eigenfunction is in  $S^\perp$ , then the negative Krein index is zero. In other words, we want it to be the case that the Hermitian matrix,  $-\lambda_0[iP_n(i\lambda_0)]|_{\mathbb{E}_{i\lambda_0}}$ , is positive definite whenever  $i\lambda_0$  is also a polynomial eigenvalue for the operator  $P_{S^\perp}P_n(\lambda)P_{S^\perp}$ .

*Remark 2.1.* In practice, mapping  $\mathbf{K}(\lambda) \mapsto \lambda^\ell \mathbf{K}(\lambda)$  for some  $\ell \in \mathbb{N}$  does not change the above property of the Krein matrix. However, as we will see, an appropriate choice of  $\ell$  gives better graphical properties regarding the determination of those polynomial eigenvalues with negative Krein signature.

*Remark 2.2.* Note that if  $\lambda = i\lambda_0 \in i\mathbb{R}$ , so that the operator  $\mathcal{P}_n(i\lambda_0)$  is Hermitian, then for  $\lambda \in i\mathbb{R}$  the elements in the second matrix can be rewritten,

$$\left( (P_{S^\perp}\mathcal{P}_n(\lambda)P_{S^\perp})^{-1}|_{P_{S^\perp}\mathcal{P}_n(\lambda)S} \right)_{ij} = \langle P_{S^\perp}\mathcal{P}_n(\lambda)s_i, (P_{S^\perp}\mathcal{P}_n(\lambda)P_{S^\perp})^{-1}P_{S^\perp}\mathcal{P}_n(\lambda)s_j \rangle.$$

**2.2. Subspace selection.** We now see how the operator coefficients may dictate the choice of the subspace  $S$ . First consider the first-order operator,

$$\mathcal{P}_1(\lambda) = \mathcal{A}_0 + \lambda\mathcal{A}_1,$$

where  $\mathcal{A}_0$  is Hermitian, and  $\mathcal{A}_1$  is skew-Hermitian. Regarding the term associated with the calculation of the negative Krein index,

$$-\lambda_0[i\mathcal{P}'_1(i\lambda_0)] = -\lambda_0(i\mathcal{A}_1).$$

If  $\psi_0$  is an eigenfunction associated with the polynomial eigenvalue, so  $\mathcal{P}_1(i\lambda_0)\psi_0 = 0$ , then

$$-\lambda_0(i\mathcal{A}_1)\psi_0 = \mathcal{A}_0\psi_0,$$

so we recover the “standard” definition of the negative Krein index for first-order star-even operators,

$$k_i^-(i\lambda_0) = n(-\lambda_0[i\mathcal{P}'_1(i\lambda_0)]|_{\mathbb{E}_{i\lambda_0}}) = n(\mathcal{A}_0|_{\mathbb{E}_{i\lambda_0}}).$$

We want the matrix  $\mathcal{A}_0|_{\mathbb{E}_{i\lambda_0}}$  to be positive definite if  $\mathbb{E}_{i\lambda_0} \subset S^\perp$ . If we choose,

$$S := N(\mathcal{A}_0) \oplus \ker(\mathcal{A}_0),$$

where  $N(\mathcal{A}_0)$  is the finite-dimensional negative subspace of  $\mathcal{A}_0$ , and  $\ker(\mathcal{A}_0)$  is the finite-dimensional kernel, then the fact that  $\mathcal{A}_0$  is positive definite on  $S^\perp$  implies that if  $i\lambda_0$  is a polynomial eigenvalue whose associated eigenfunction resides in  $S^\perp$ , then the negative Krein index will be zero. Note that in this case  $P_S$  and  $P_{S^\perp}$  will be spectral projections. Further note that with this choice of subspace that if a pole of the Krein matrix corresponds to purely imaginary polynomial eigenvalue, then it will necessarily have positive Krein index. Consequently, all purely imaginary polynomial eigenvalues with negative Krein index will be captured by solving  $\det \mathbf{K}_S(\lambda) = 0$ .

Now, consider the second-order operator

$$\mathcal{P}_2(\lambda) = \mathcal{A}_0 + \lambda\mathcal{A}_1 + \lambda^2\mathcal{A}_2,$$

where  $\mathcal{A}_0, \mathcal{A}_2$  are Hermitian, and  $\mathcal{A}_1$  is skew-Hermitian. We have

$$-\lambda_0[i\mathcal{P}'_2(i\lambda_0)] = -\lambda_0(i\mathcal{A}_1) + 2\lambda_0^2\mathcal{A}_2.$$

If  $\psi_0$  is an eigenfunction associated with the polynomial eigenvalue, so  $\mathcal{P}_2(i\lambda_0)\psi_0 = 0$ ,

$$(-\lambda_0(i\mathcal{A}_1) + 2\lambda_0^2\mathcal{A}_2)\psi_0 = (\mathcal{A}_0 + \lambda_0^2\mathcal{A}_2)\psi_0.$$

The negative Krein index can be alternatively defined,

$$k_i^-(i\lambda_0) = n(-\lambda_0[i\mathcal{P}'_2(i\lambda_0)]|_{\mathbb{E}_{i\lambda_0}}) = n((\mathcal{A}_0 + \lambda_0^2\mathcal{A}_2)|_{\mathbb{E}_{i\lambda_0}}).$$

In order for it to be the case that the matrix  $(\mathcal{A}_0 + \lambda_0^2\mathcal{A}_2)|_{\mathbb{E}_{i\lambda_0}}$  is guaranteed to be positive definite, it must be true that the eigenspace  $\mathbb{E}_{i\lambda_0}$  resides in the positive space of the operator  $\mathcal{A}_0 + \lambda_0^2\mathcal{A}_2$ . In the applications we consider the operator  $\mathcal{A}_2$  will be positive definite. In this case, if we again choose,

$$S := N(\mathcal{A}_0) \oplus \ker(\mathcal{A}_0),$$

then the operator,

$$P_{S^\perp}(\mathcal{A}_0 + \lambda_0^2\mathcal{A}_2)P_{S^\perp} = P_{S^\perp}\mathcal{A}_0P_{S^\perp} + \lambda_0^2P_{S^\perp}\mathcal{A}_2P_{S^\perp},$$

will be positive definite. Consequently, if  $i\lambda_0$  is a polynomial eigenvalue whose associated eigenfunction resides in  $S^\perp$ , then the negative Krein index will be zero.

**3. The Krein eigenvalues.** Since  $P_n(\lambda)$  is a star-even polynomial operator, the Krein matrix is a self-adjoint meromorphic family of operators in the spectral parameter,  $\lambda$ . In particular, the Krein matrix is Hermitian for purely imaginary  $\lambda$ . Henceforth, write  $\lambda = iz$  for  $z \in \mathbb{R}$ , and write the Krein matrix as

$$\mathbf{K}_S(z) = \mathcal{P}_n(iz)|_S - \mathcal{P}_n(iz)P_{S^\perp}(P_{S^\perp}\mathcal{P}_n(iz)P_{S^\perp})^{-1}P_{S^\perp}\mathcal{P}_n(iz)|_S.$$

Since the Krein matrix is Hermitian for real  $z$ , for each value of  $z$  there are  $n_S$  real-value eigenvalues,  $r_j(z)$ . These eigenvalues of the Krein matrix are called the *Krein eigenvalues*. The Krein eigenvalues are real meromorphic, as are the associated spectral projections. In particular, if the Krein eigenvalues are simple, the associated eigenvectors are real meromorphic. See Kato [22, Chapter VII.3] for the details.

Since

$$\det \mathbf{K}_S(z) = \prod_{j=1}^{n_S} r_j(z),$$

finding the zeros of the determinant of the Krein matrix is equivalent to finding the zero set of each of the Krein eigenvalues. One of the most important properties of the Krein eigenvalues is that the sign of the derivative at a simple zero is related to the Krein index of that polynomial eigenvalue. In order to see this, we start with

$$(3.1) \quad \mathbf{K}_S(z)\mathbf{v}_j(z) = r_j(z)\mathbf{v}_j(z) \quad \rightsquigarrow \quad r'_j(z) = \frac{\mathbf{v}_j(z)^a \mathbf{K}'_S(z) \mathbf{v}_j(z)}{|\mathbf{v}_j(z)|^2}.$$

The latter equality is a solvability condition which follows upon noting that both the Krein eigenvalue and its associated eigenvector are meromorphic and consequently have convergent Taylor expansions. If  $r_j(z) = 0$ , then the components of the associated eigenvector correspond to the various basis elements in the subspace  $S$ ; namely, the associated eigenfunction is given by

$$(3.2) \quad \psi = \sum_{k=1}^{n_S} v_k^j s_k + s^\perp, \quad \mathbf{v}_j = \begin{pmatrix} v_1^j \\ v_2^j \\ \vdots \\ v_{n_S}^j \end{pmatrix},$$

where the element  $s^\perp$  is determined via (2.3),

$$s^\perp = - \sum_{k=1}^{n_S} v_k^j (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_k.$$

We now compute  $\mathbf{K}'(z)$ . For the first term in the Krein matrix,

$$\frac{d}{dz} \langle s_i, \mathcal{P}_n(iz) s_j \rangle = \langle s_i, [i\mathcal{P}'_n(iz)] s_j \rangle.$$

The operator  $i\mathcal{P}'_n(iz)$  is Hermitian. Differentiating the second term requires repeated applications of the product rule, as well as using the fact that the operator  $\mathcal{P}_n(iz)$  is Hermitian. Since

$$\frac{d}{dz} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} = -(P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} [i\mathcal{P}'_n(iz)] (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1},$$



upon some simplification we can write

$$\begin{aligned} \frac{d}{dz} \langle s_i, P_{S^\perp} \mathcal{P}_n(iz) (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_j \rangle = \\ \langle s_i, [\mathbf{i} \mathcal{P}'_n(iz)] P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_j \rangle \\ + \langle s_i, \mathcal{P}_n(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} [\mathbf{i} \mathcal{P}'_n(iz)] s_j \rangle \\ - \langle s_i, \mathcal{P}_n(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} [\mathbf{i} \mathcal{P}'_n(iz)] (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_j \rangle. \end{aligned}$$

The right-hand side has the compact form

$$\frac{d}{dz} \langle s_i, P_{S^\perp} \mathcal{P}_n(iz) (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_j \rangle = \langle s_i, (\mathcal{R} + \mathcal{R}^a) s_j \rangle - \langle s_i, \mathcal{S} s_j \rangle,$$

where

$$\begin{aligned} \mathcal{R} &:= \mathcal{P}_n(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} [\mathbf{i} \mathcal{P}'_n(iz)] \\ \mathcal{S} &:= \mathcal{P}_n(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} [\mathbf{i} \mathcal{P}'_n(iz)] (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz). \end{aligned}$$

In conclusion, the derivative of the Krein matrix is

$$(3.3) \quad \mathbf{K}'(z) = [\mathbf{i} \mathcal{P}'_n(iz)]_S + \mathcal{S}|_S - (\mathcal{R} + \mathcal{R}^a)|_S,$$

where the operators  $\mathcal{R}, \mathcal{S}$  are defined above.

We now compute the Krein index using our decomposition of an eigenfunction.

For the sake of exposition, let us assume that the polynomial eigenvalue is simple.

Using the decomposition (3.2) with  $\mathbf{K}_s(z) \mathbf{v}_j(z) = \mathbf{0}$ , we have

$$[\mathbf{i} \mathcal{P}'_n(iz)] \psi = \sum_{k=1}^{n_S} v_k^j [\mathbf{i} \mathcal{P}'_n(iz)] s_k - \sum_{k=1}^{n_S} v_k^j [\mathbf{i} \mathcal{P}'_n(iz)] (P_{S^\perp} \mathcal{P}_n(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_n(iz) s_k.$$

Upon taking the inner product with  $\psi$ , and using the fact that  $\mathcal{P}_n(iz)$  is Hermitian,

$$\langle \psi, [\mathbf{i} \mathcal{P}'_n(iz)] \psi \rangle = \mathbf{v}_j(z)^a ([\mathbf{i} \mathcal{P}'_n(iz)]|_S + \mathcal{S}|_S - (\mathcal{R} + \mathcal{R}^a)|_S) \mathbf{v}_j(z).$$

Upon comparing with (3.3) we conclude

$$\langle \psi, [\mathbf{i} \mathcal{P}'_n(iz)] \psi \rangle = \mathbf{v}_j(z)^a \mathbf{K}'(z) \mathbf{v}_j(z),$$

where the eigenfunction  $\psi$  has the expansion provided for in (3.2).

Going back to (3.1), we have that the derivative of the Krein eigenvalue can be

expressed in terms of the eigenfunction as

$$r'_j(z) = \frac{\langle \psi, [\mathbf{i} \mathcal{P}'_n(iz)] \psi \rangle}{|\mathbf{v}_j(z)|^2}.$$

Going further back to the definition of the negative Krein index, we can conclude the

desired result. If  $iz$  is a polynomial eigenvalue with  $r_j(z) = 0$ , then the Krein index

is related through the derivative via

$$k_i^-(iz) = \begin{cases} 0, & z r'_j(z) < 0 \\ 1, & z r'_j(z) > 0. \end{cases}$$

Since our goal is to quickly and easily read off the Krein signature via a graph of the Krein eigenvalues, we will redefine the Krein matrix as

$$\mathbf{K}_S(z) = -z [\mathcal{P}_n(\mathrm{i}z)|_S - \mathcal{P}_n(\mathrm{i}z)P_{S^\perp}(P_{S^\perp}\mathcal{P}_n(\mathrm{i}z)P_{S^\perp})^{-1}P_{S^\perp}\mathcal{P}_n(\mathrm{i}z)|_S].$$

The Krein eigenvalues for the new matrix are related to the original matrix via  $r_j(z) \mapsto -zr_j(z)$ . Thus, at a zero of the Krein eigenvalue we have the mapping  $r'_j(z) \mapsto -zr'_j(z)$ , so for the new Krein matrix we have the relationship

$$k_i^-(\mathrm{i}z) = \begin{cases} 0, & r'_j(z) > 0 \\ 1, & r'_j(z) < 0. \end{cases}$$

A positive slope of a Krein eigenvalue at a zero corresponds to a polynomial eigenvalue with positive signature, whereas a negative slope shows that the polynomial eigenvalue has negative Krein signature.

If the zero of a Krein eigenvalue is not simple, then the corresponding polynomial eigenvalue has a Jordan chain, and the negative Krein index depends upon the length of the chain, see [14, Section 2.2] and the references therein. For example, if  $r_j(z) = r'_j(z) = 0$  with  $r''_j(z) \neq 0$ , then there will be a Jordan chain of length two; moreover, the negative Krein index associated with the Jordan chain will be one. In general, a zero of order  $m$  implies a Jordan chain of length  $m$ , and the negative Krein index associated with that chain will be roughly half the length of the chain. We will not provide any more details here, as in our examples the polynomial eigenvalues will be simple. In summary, we have the following result:

**THEOREM 3.1.** *Consider the star-even polynomial,*

$$\mathcal{P}_n(\lambda) = \sum_{j=1}^n \lambda^j \mathcal{A}_j,$$

where on a Hilbert space,  $X$ , with inner-product,  $\langle \cdot, \cdot \rangle$ , the operator coefficients satisfy:

- (a)  $\mathcal{A}_0$  has compact resolvent with  $\mathfrak{n}(\mathcal{A}_0)$  being finite
- (b) setting  $P_{\mathcal{A}_0} : X \mapsto \ker(\mathcal{A}_0)^\perp$  to be the spectral projection, and  $P_{\mathcal{A}_0}^\perp = \mathcal{I} - P_{\mathcal{A}_0}$ , the operators,

$$(P_{\mathcal{A}_0}^\perp \mathcal{A}_0 P_{\mathcal{A}_0}^\perp)^{-1} P_{\mathcal{A}_0}^\perp \mathcal{A}_j P_{\mathcal{A}_0}^\perp : \ker(\mathcal{A}_0)^\perp \mapsto \ker(\mathcal{A}_0)^\perp, \quad j = 1, \dots, n,$$

are compact.

For a given finite-dimensional subspace,  $S \subset X$ , and orthogonal projection,  $P_{S^\perp} : X \mapsto S^\perp$ , the associated Krein matrix is

$$\mathbf{K}_S(z) = -z [\mathcal{P}_n(\mathrm{i}z)|_S - \mathcal{P}_n(\mathrm{i}z)P_{S^\perp}(P_{S^\perp}\mathcal{P}_n(\mathrm{i}z)P_{S^\perp})^{-1}P_{S^\perp}\mathcal{P}_n(\mathrm{i}z)|_S].$$

For  $z \in \mathbb{R}$  the Krein eigenvalues,  $r_j(z)$  for  $j = 1, \dots, \dim[S]$ , are meromorphic and have the properties that if  $\lambda = \mathrm{i}z$  is a polynomial eigenvalue with  $\mathcal{P}_n(\mathrm{i}z)\psi = 0$ :

- (a) then either
  - $r_j(z) = 0$  for at least one  $j$ , or
  - $\psi \in S^\perp$ .
- (b) if  $z \in \mathbb{R}$ , and if  $r_j(z) = 0$  for some  $j$ , then the Krein signature is determined by the slope of the graph of the Krein eigenvalue,

$$k_i^-(\mathrm{i}z) = \begin{cases} 0, & r'_j(z) > 0 \\ 1, & r'_j(z) < 0. \end{cases}$$

*Remark 3.2.* Recall that the choice,

$$S = N(\mathcal{A}_0) \oplus \ker(\mathcal{A}_0),$$

ensures that all polynomial eigenvalues with negative Krein signature are seen as zeros of one or more Krein eigenvalues.

In its general form the Krein matrix looks to be complicated, and does not appear to have an underlying intuitively understood structure. However, as we shall see in our subsequent examples, the Krein matrix can have intimate connections with dispersion relations, the Hale-Sandstede-Lin's method for constructing multi-pulses, etc.

**4. First application: modulational instabilities for small amplitude periodic solutions.** For our first application we show how the Krein matrix can be used to understand the existence of instability bubbles for small spatially periodic waves to dispersive systems. The instabilities will not necessarily be associated with high-frequency (long wavelength) perturbations. Without loss of generality we will assume the spatial period is  $2\pi$ .

Regarding the existence problem we will assume it is of the form,

$$(4.1) \quad \mathcal{L}u - cu + f(u) = 0,$$

where

$$(a) \quad \mathcal{L} = \sum_{j=0}^N a_j \ell^{2j} \partial_x^{2j} \text{ with } \ell, (-1)^N a_{2N} > 0$$

$$(b) \quad c \in \mathbb{R} \text{ is a free parameter}$$

$$(c) \quad f(u) \text{ is a smooth nonlinearity with } f(0) = f'(0) = 0.$$

The parameter  $\ell$  can be adjusted via a rescaling of  $x$ . The operator  $\mathcal{L}$  is self-adjoint under the inner-product,

$$\langle f, g \rangle = \int_0^{2\pi} f(x) \overline{g(x)} dx.$$

*Remark 4.1.* The nonlinearity could be more general,  $f = f(u, \partial_x u, \dots)$ . All that is required is that it be smooth and (at least) quadratic in the arguments near the origin, and that it be unchanged under reversibility,  $x \mapsto -x$ .

We briefly sketch the argument leading to the existence of a family of small spatially periodic solutions. The details can be found in [11, Theorem 3.15]. The characteristic polynomial associated with the ordinary differential operator  $\mathcal{L}$  is

$$p_{\mathcal{L}}(r, \ell) = \sum_{j=0}^N a_j \ell^{2j} r^{2j}.$$

Regarding the characteristic polynomial we assume there is an  $\ell_0$  such that,

$$(a) \quad \partial_r p_{\mathcal{L}}(i, \ell_0) \neq 0$$

$$(b) \quad \text{upon setting,}$$

$$(4.2) \quad c_0 := p_{\mathcal{L}}(i, \ell_0) = \sum_{j=0}^N (-1)^j a_j \ell_0^{2j},$$

$$\text{there is no positive real } k \neq 1 \text{ such that } p_{\mathcal{L}}(ik, \ell_0) - c_0 = 0.$$

There will then exist a family of  $2\pi$ -periodic solutions, say  $U(x)$ , with the properties:

- (a)  $U(x) = U(-x)$
- (b)  $U(x) = \epsilon A \cos(x) + \mathcal{O}(\epsilon^2)$  for  $A > 0$
- (c)  $\ell = \ell_0 + \mathcal{O}(\epsilon)$  (the  $\mathcal{O}(\epsilon)$  terms depend on  $A$ ).

If (4.2) above does not hold, i.e., if there are other purely imaginary roots to  $p_{\mathcal{L}}(r, \ell_0) - \beta_0 = 0$ , then the equations on the center-manifold will still be reversible. However, the dimension of the manifold (equal to the number of purely imaginary roots, counting multiplicity) increases, and since the reduced system is no longer planar it is not clear if there are still periodic (versus quasi-periodic) solutions. The case of a second additional imaginary root,  $\pm iq$  with  $q > 1$ , is discussed by [11, Chapter 4.3.4]. If  $q$  is irrational, or if  $q \geq 5$ , only KAM tori are expected, and consequently only quasi-periodic solutions. In the case of strong resonance,  $q = 2$ , the equations on the center manifold are completely integrable, and there can be periodic orbits, homoclinic orbits, and orbits homoclinic to periodic orbits. The other resonant case of  $q = 3$  is still open. In conclusion, we can safely assume the existence of small  $2\pi$ -periodic solutions to (4.1).

We now consider the spectral stability of these spatially periodic solutions. Consider the KdV-like and first-order-in-time Hamiltonian system,

$$(4.3) \quad \partial_t u + \partial_x (\mathcal{L}u + f(u)) = 0.$$

While we will not do it here, much of what follows can easily be adapted to cover second-order-in-time systems. The nonlinearity  $f(u)$  satisfies the assumption (c) above, while

$$\mathcal{L}u = \sum_{j=0}^N a_j \partial_x^{2j} u, \quad (-1)^N a_{2N} > 0.$$

In traveling coordinates,  $z := x - ct$ , the equation becomes,

$$\partial_t u + \partial_z (\mathcal{L}u - cu + f(u)) = 0, \quad \partial_x^{2j} \mapsto \partial_z^{2j}.$$

Upon rescaling of time and space,

$$\tau = \ell t, \quad y = \ell z,$$

we have the PDE to be studied,

$$(4.4) \quad \partial_\tau u + \partial_y (\mathcal{L}u - cu + f(u)) = 0,$$

where

$$\mathcal{L}u = \sum_{j=0}^N a_j \ell^{2j} \partial_y^{2j} u, \quad (-1)^N a_{2N} > 0.$$

Following the previous discussion, upon setting,

$$c_0 := p_{\mathcal{L}}(i, \ell_0),$$

where  $\ell_0$  is chosen so that  $p_{\mathcal{L}}(ik, \ell_0) - c_0 = 0$  has no integral solutions for  $k > 1$ , we know there is a family of small  $2\pi$ -periodic solutions,  $U(x) = \mathcal{O}(\epsilon)$ , for  $0 < \epsilon \ll 1$ .

We now consider the spectral stability of such solutions. The linearized problem is,

$$\partial_\tau v + \partial_y (\mathcal{L}v - c_0 v + f'(U)v) = 0, \quad |f'(U)| = \mathcal{O}(\epsilon).$$

Using separation of variables,  $v(y, \tau) = e^{\lambda \tau} v(y)$ , we arrive at the spectral problem,

$$(4.5) \quad \lambda v + \partial_y (\mathcal{L}v - c_0 v + f'(U)v) = 0, \quad |f'(U)| = \mathcal{O}(\epsilon).$$

We use a Bloch decomposition to understand the spectral problem, see [20, Chapter 3.3]. Writing for  $-1/2 < \mu \leq 1/2$ ,

$$v(y) = e^{i\mu y} w(y), \quad w(y + 2\pi) = w(y),$$

the problem (4.5) becomes,

$$(4.6) \quad \lambda w + (\partial_y + i\mu) (\mathcal{L}_\mu w - c_0 w + f'(U)w) = 0, \quad |f'(U)| = \mathcal{O}(\epsilon),$$

where

$$\mathcal{L}_\mu = \sum_{j=0}^N a_{2j} \ell_0^{2j} (\partial_y + i\mu)^{2j}.$$

Because the underlying wave is even in  $x$ , it is sufficient to consider  $0 \leq \mu \leq 1/2$ ; in particular, if  $\lambda$  is an eigenvalue associated with  $\mu$ , then  $\bar{\lambda}$  is an eigenvalue associated with  $-\mu$ , see [12, Section 4]. For fixed  $\mu$  the spectrum will be discrete, countable, and have an accumulation point only at  $\infty$ . The full spectrum, which is essential spectra only, will be the union of all the point spectra as  $\mu$  is varied over the range.

We are henceforth interested only in sideband instabilities,  $\mu > 0$ . Set,

$$\mathcal{A}_0 := \mathcal{L}_\mu - c_0 + f'(U).$$

The operator  $\mathcal{A}_0$  is self-adjoint on the space of  $2\pi$ -periodic functions endowed with the natural  $L^2[0, 2\pi]$  inner product. The invertible operator  $\partial_y + i\mu$  is skew-Hermitian. Since  $\mathcal{A}_0$  is self-adjoint with smooth dependence on parameters, the spectrum of this operator spectrum is smooth in  $(\mu, \epsilon)$ . The same can be said of the composition,  $(\partial_y + i\mu)\mathcal{A}_0$ , except at possibly the finite number of points where there are Jordan chains. Consequently, we will first consider the spectral problem when  $\epsilon = 0$ . Afterwards, we will make generic statements about what will happen for  $\epsilon > 0$  small.

For  $0 < \mu \leq 1/2$  we rewrite the spectral problem in the star-even form,

$$(4.7) \quad \mathcal{A}_0 w + \lambda \mathcal{A}_1 w = 0, \quad \mathcal{A}_1 := (\partial_y + i\mu)^{-1}.$$

The boundary conditions associated with this problem are periodic,  $w(y + 2\pi) = w(y)$ . We will analyze the spectral problem using the Krein matrix. First assume  $\epsilon = 0$ , so that  $f'(U) \equiv 0$ . The spectrum for (4.7) is straightforward to compute using a Fourier analysis. Letting  $w(y) = e^{iny}$  for  $n \in \mathbb{Z}$  we get a sequence of problems,

$$(4.8) \quad d(n, \mu) + \lambda \frac{1}{i(n + \mu)} = 0,$$

where the first term is the dispersion relation associated with the steady-state problem,

$$d(n, \mu) := \sum_{j=0}^N (-1)^j a_{2j} \ell_0^{2j} (n + \mu)^{2j} - c_0.$$

The existence assumption implies  $d(\pm 1, 0) = 0$ . For small  $\mu$  we have the expansions,

$$(4.9) \quad d(\pm 1, \mu) = \pm \left( 2 \sum_{j=1}^N (-1)^j j a_j \right) \mu + \mathcal{O}(\mu^2).$$

Consequently,  $d(+1, \mu)d(-1, \mu) < 0$  for small  $\mu$ , so  $n(\mathcal{A}_0) \geq 1$  for  $\mu$  sufficiently small. The assumption  $(-1)^N a_{2N} > 0$  implies there is an  $N_0$  such that  $d(n, \mu) > 0$  for  $|n| \geq N_0$ . Consequently,  $n(\mathcal{A}_0) < \infty$ . By continuity this will remain true for small  $\epsilon > 0$ .

We now construct the Krein matrix. Assume there is a sequence  $n_1, n_2, \dots, n_q$  such that  $d(n, \mu) < 0$  for  $n \in \{n_1, n_2, \dots, n_q\}$ , and  $d(n, \mu) > 0$  for  $n \notin \{n_1, n_2, \dots, n_q\}$ . Clearly,  $n(\mathcal{A}_0) = q$ . We take as our space  $S = N(\mathcal{A}_0)$ ,

$$S = \text{span}\{e^{in_1 y}, e^{in_2 y}, \dots, e^{in_q y}\}.$$

Since

$$P_1(iz)S = S, \quad P_1(iz)S^\perp = S^\perp,$$

the Krein matrix as described in [Theorem 3.1](#) collapses to

$$\begin{aligned} K_S(z) &= -z\mathcal{P}_1(iz)|_S \\ &= -z \text{diag} \left( d(n_1, \mu) + \frac{z}{n_1 + \mu}, \dots, d(n_q, \mu) + \frac{z}{n_q + \mu} \right). \end{aligned}$$

The expected poles, which are the eigenvalues of the sandwiched operator,

$$P_{S^\perp} \mathcal{P}_1(iz) P_{S^\perp} = P_1(iz)S^\perp,$$

are located at  $z_n^p = -(n + \mu)d(n, \mu)$  for  $n \notin \{n_1, n_2, \dots, n_q\}$ , and are removable singularities. All of the poles are polynomial eigenvalues for the spectral problem. Since they correspond to removable singularities, the polynomial eigenvalues all have positive Krein signature. Generically, the poles will not be removable for  $\epsilon > 0$ .

The Krein eigenvalues are

$$r_j(z) = -z \left( d(n_j, \mu) + \frac{z}{n_j + \mu} \right), \quad j = 1, \dots, q.$$

The nonzero zeros of the Krein eigenvalues,

$$z_j^n = -(n_j + \mu)d(n_j, \mu), \quad j = 1, \dots, q,$$

satisfy

$$r'_j(z_j^n) = d(n_j, \mu) < 0,$$

so these zeros correspond to polynomial eigenvalues with negative Krein signature. In conclusion, via Fourier analysis we have located all of the polynomial eigenvalues, and through the Krein eigenvalues we have identified those which have a negative Krein index.

*Remark 4.2.* For constant states the Krein signature can be directly computed from the dispersion relation, see [\[8\]](#).

When  $\epsilon = 0$  the wave is spectrally stable, and all of the spectra is purely imaginary. For  $\epsilon > 0$  a spectral instability arises through the collision of a purely imaginary polynomial eigenvalue with positive Krein index and one with negative Krein index. This collision generically leads to a Hamiltonian-Hopf bifurcation, see [\[20, Chapter 7.1.2\]](#) and the references therein. If for a fixed  $\mu_0$  there is a polynomial eigenvalue with positive real part, then such polynomial eigenvalues will exist for  $\mu$  in a neighborhood of  $\mu_0$ . If for  $\mu_0$  the polynomial eigenvalue with positive real part is simple, then the

union of all polynomial eigenvalues for  $\mu$  in a neighborhood of  $\mu_0$  will form a smooth curve. We will call this curve an instability bubble. In our example any instability bubble will have an  $\mathcal{O}(1)$  imaginary part; consequently, they will not be related to instability curves coming from the origin which arise due to a long wavelength modulational instability. A bubble intersects the imaginary axis, and because of the  $\{\lambda, -\bar{\lambda}\}$  reflection symmetry about the imaginary axis, the curve on the left of the imaginary axis is a mirror image of that on the right.

The Krein eigenvalues reflect this collision of polynomial eigenvalues with opposite index in one of two possible ways. The first is that a Krein eigenvalue has a double zero at the time of collision, see [14, Lemma 2.8]. For small waves this cannot happen, as the explicit form of the Krein eigenvalues shows that all of the zeros are simple for the limiting zero amplitude wave.

As for the other possible collision scenario, recall that when  $\epsilon = 0$  a zero of a Krein eigenvalue corresponds to a polynomial eigenvalue with negative Krein signature, while all the removable singularities, i.e., polynomial eigenvalues of the operator  $P_{S^\perp} P_1(iz) P_{S^\perp}$ , correspond to polynomial eigenvalues with positive Krein signature. If a simple zero is isolated, then the Krein matrix being meromorphic implies via a winding number calculation that the zero remains simple for small perturbations. Moreover, the spectral symmetry implies the polynomial eigenvalue must remain purely imaginary. Now, suppose that a simple zero coincides with a simple removable singularity, so when  $\epsilon = 0$  the winding number is again one. For the problem at hand this situation is realized when a zero of one of the Krein eigenvalues intersects one of the removable singularities,  $z_n^p$ . In general, this intersection must be computed numerically. Assume that upon perturbation the singularity is no longer removable - it will remain simple. In this case the invariance of the winding number to small perturbation implies there must now be two zeros. The spectral symmetry implies these correspond to either two purely imaginary polynomial eigenvalues, or a pair of polynomial eigenvalues with nonzero real part. In the former case, the invariance of the HKI to small perturbation implies that one polynomial eigenvalue will have positive Krein signature, whereas the other will have negative Krein signature. The latter case corresponds to the onset of a Hamiltonian-Hopf bifurcation. An analytic argument which leads to the same conclusion is presented in [14, Section 2.4].

In conclusion, the total number of bubbles that can form is bounded above by the number of intersections of Krein eigenvalues with poles. Supposing that the HKI is fixed for all  $\mu$ , this leaves open the possibility that the number of bubbles is greater than  $K_{\text{Ham}}$ . For example, suppose  $K_{\text{Ham}} = 2$ , so that for each  $\mu$  there can be at most two polynomial eigenvalues with positive real part. Since there will be two Krein eigenvalues, for each  $\mu$  there can be at most two associated bubbles. However, overall there can be more than two bubbles. Suppose there is a sequence  $0 < \mu_1 < \mu_2 < \dots < \mu_N$  for which a Krein eigenvalue intersects a pole. A Hamiltonian-Hopf bifurcation is then possible for  $\mu$  near each  $\mu_j$ , which leaves open the possibility of having up to  $N$  bubbles.

*Remark 4.3.* More generally, if  $k$  polynomial eigenvalues with negative signature coincide with a removable singularity for the Krein matrix of order  $\ell$ , then upon perturbation the invariance of the winding number implies that  $k + \ell$  polynomial eigenvalues will be created via the collision. The invariance of the HKI implies that  $k = k_c + k_i^-$ , where here  $k_i^-$  corresponds to the number of purely imaginary polynomial eigenvalues with negative Krein signature which are close to the unperturbed eigenvalue, and  $k_c$  is the number of polynomial eigenvalues with positive real part which

are close to the unperturbed eigenvalue. As for the number of polynomial eigenvalues associated with the order of the removable singularity,  $\ell = k_c + k_i^+$ , where here  $k_i^+$  corresponds to the number of purely imaginary polynomial eigenvalues with positive Krein signature which are close to the unperturbed eigenvalue.

*Remark 4.4.* For small amplitude wave the intersection of a Krein eigenvalue with a (potentially) removable singularity of the Krein matrix can be noted without actually computing a Krein eigenvalue. For fixed  $\mu$  the Krein eigenvalues are dispersion curves that correspond to polynomial eigenvalues with negative Krein index, and the poles correspond to dispersion curves with positive Krein index. If the two curves intersect, then there is a collision of polynomial eigenvalues with opposite Krein signature. This graphical approach is the one taken by [8, 23, 35].




FIG. 1. (color online) Plots of the dispersion relations,  $z_n(\mu)$ , for the linearization of (4.11) for relevant values of  $n$  when  $b = -8/15$ . A dotted curve corresponds to an eigenvalue with negative Krein index, while a solid curve shows an eigenvalue with positive index. Not only is a Hamiltonian-Hopf bifurcation possible for small  $\mu$ , it is possible for  $\mu \sim 0.21$  and  $\mu \sim 0.37$ .

For a particular example, consider the fifth-order KdV-like equation,

$$(4.10) \quad \partial_t u + \partial_x \left( \frac{2}{15} \partial_x^4 u - b \partial_x^2 u + \frac{3}{2} u^2 + \frac{1}{2} [\partial_x u]^2 + u \partial_x^2 u \right) = 0.$$

This weakly nonlinear long-wave equation arises as an approximation to the classical gravity-capillary water-wave problem [5]. Here  $u(x, t)$  is the surface elevation with respect to the underlying normal water height, and  $b \in \mathbb{R}$  is the offset of the Bond number (a measure of surface tension) from the value  $1/3$ . In traveling coordinates,  $z = x - ct$ , the equation (4.10) becomes

$$(4.11) \quad \partial_t u + \partial_z \left( \frac{2}{15} \partial_z^4 u - b \partial_z^2 u - cu + \frac{3}{2} u^2 + \frac{1}{2} [\partial_z u]^2 + u \partial_z^2 u \right) = 0.$$

The wavespeed  $c$  is a free parameter. To the best of our knowledge the spectral stability of small periodic waves to equation (4.10) has not yet been studied. However,



the spectral stability of small spatially periodic waves to the Kawahara equation, which is (4.10) with the last two terms in the open brackets removed, was recently studied by [35].




FIG. 2. (color online) Plots of the absolute value of the real part of the spectrum for various values of  $\mu$  for a wave with approximate amplitude  $2.3 \times 10^{-2}$ . The plot on the left is for  $\mu$  values near the zero/pole collision point  $\mu \sim 0.207$ , and the plot on the right is for  $\mu$  values near the zero/pole collision point  $\mu \sim 0.368$ . The  $\mu$  value for which the collision occurs is marked by a (red) cross.

First consider the existence problem. As discussed by [32, Section 4] (also see [5]), the fourth-ODE,

$$\frac{2}{15}\partial_z^4 u - b\partial_z^2 u + \frac{3}{2}u^2 + \frac{1}{2}[\partial_z u]^2 + u\partial_z^2 u = 0,$$

is a reversible Hamiltonian system. The position and momentum variables are

$$q_1 = u, \quad q_2 = \partial_z u, \quad p_1 = -\frac{2}{15}\partial_z^3 u + b\partial_z u - u\partial_z u, \quad p_2 = \frac{2}{15}\partial_z^2 u,$$

and the (analytic) Hamiltonian is

$$H = -\frac{1}{2}q_1^3 - \frac{1}{2}cq_1^2 + p_1q_1 - \frac{1}{2}bq_2^2 + \frac{15}{4}p_2^2 + \frac{1}{2}q_1q_2^2.$$

The symplectic matrix for the system is the canonical one. Setting

$$c = c_0 := \frac{2}{15} + b,$$

the eigenvalues for the linearization of this Hamiltonian system about the origin satisfy

$$r^2 = -1, \quad r^2 = 1 + \frac{15}{2}b.$$

If  $b > -2/15$ , then the center-manifold is two-dimensional, and the existence of a family of periodic orbits follows from reversibility. If  $b < -2/15$ , but  $b \neq -2(1 +$

595  $m^2)/15$  for  $m = 1, 2, \dots$  (the non-resonance condition), then one can invoke the  
 596 Lyapunov center theorem to conclude the existence of a family of small periodic  
 597 orbits with period close to  $2\pi$  (see [3, 38] for a discussion). In either case, the period  
 598 can be fixed to be  $2\pi$  via a rescaling of the spatial variable. We will assume for that  
 599 sake of exposition that  $b = -8/15$ , so  $c_0 = -6/15$ . For this value of  $b$  the ODE system  
 600 is not in resonance.




FIG. 3. (color online) Plots of the Krein eigenvalues for the trivial state (left figures) and for a wave with approximate amplitude  $2.3 \times 10^{-2}$  (right figures). The top two figures show the situation at the zero/pole collision point,  $\mu \sim 0.368$ . The (red) circles correspond to polynomial eigenvalues, and the (red) cross is the spurious zero of the Krein eigenvalues. The (green) vertical lines are poles of the Krein matrix. In each quadrant the bottom figure is a blow-up of the top figure near the polynomial eigenvalues of interest. Upon perturbation the zeros of the Krein eigenvalues remain purely real.

601 We now consider the spectral stability of the periodic wave. For the unperturbed  
 602 problem the dispersion relationship is,

$$603 \quad d(n, \mu) = \frac{2}{15}(n + \mu)^4 - \frac{8}{15}(n + \mu)^2 + \frac{6}{15}.$$

604 It is straightforward to check that  $d(n, \mu) > 0$  for  $\mu \notin \{-2, +1\}$ . Moreover, we have  
 605  $d(+1, \mu) < 0$ , and

$$606 \quad d(-2, \mu) \begin{cases} > 0, & 0 < \mu < \mu_{\text{ch}} \\ < 0, & \mu_{\text{ch}} < \mu < 1/2, \end{cases}$$

607 where

$$608 \quad \mu_{\text{ch}} := 2 - \sqrt{3} \sim 0.26795.$$

Consequently,

$$n(\mathcal{A}_0) = \begin{cases} 1, & 0 < \mu < \mu_{\text{ch}} \\ 2, & \mu_{\text{ch}} < \mu < 1/2. \end{cases}$$

Since the negative index of an invertible operator is unchanged for small perturbations, we know there is a  $0 < \mu_0 \ll 1$  such that if  $\mu > \mu_0$ , then  $n(\mathcal{A}_0)$  remains unchanged for sufficiently small  $\epsilon$ . Going back to equation (1.1), we then know that for small  $\epsilon$  the HKI is,

$$k_r + k_c + k_i^- = \begin{cases} 1, & \mu_0 < \mu < \mu_{\text{ch}} \\ 2, & \mu_{\text{ch}} < \mu < 1/2. \end{cases}$$

If there are instability bubbles for the perturbed problem, there can be at most one for  $\mu < \mu_{\text{ch}}$ , and at most two for  $\mu_{\text{ch}} < \mu < 1/2$ . For  $0 \leq \mu < \mu_0$  a curve of unstable spectra may arise from the origin. We will not consider that here, but an example calculation for the KdV with general nonlinearity is provided in [12, Section 4].




FIG. 4. (color online) Plots of the Krein eigenvalues for the trivial state (left figures) and for a wave with approximate amplitude  $2.3 \times 10^{-2}$  (right figures). The top two figures show the situation at the zero/pole collision point,  $\mu = 0.3585$ . The (red) circles correspond to polynomial eigenvalues, and the (red) cross is the spurious zero of the Krein eigenvalues. The (green) vertical lines are poles of the Krein matrix. In each quadrant the bottom figure is a blow-up of the top figure near the polynomial eigenvalues of interest. Note the existence of a Hamiltonian-Hopf bifurcation upon the perturbation.

A picture of the dispersion curves for the full problem,

$$z_n(\mu) = -(n + \mu)d(n, \mu), \quad n \in \mathbb{Z},$$

is provided in Figure 1 for relevant values of  $n$ . If the curve is dotted, then for fixed  $\mu$  that corresponds to a polynomial eigenvalue with negative Krein signature. The solid curves correspond to polynomial eigenvalues with positive Krein signature. There are two possible values for which a bubble may appear:

$$z_{-2}(\mu) = z_{+1}(\mu) \rightsquigarrow \mu = \frac{1}{10} \left( 5 - \sqrt{5(2\sqrt{129} - 21)} \right) \sim 0.20711,$$

and

$$z_0(\mu) = z_{-2}(\mu) \rightsquigarrow \mu = 1 - \frac{1}{5}\sqrt{10} \sim 0.36754.$$

Consequently, for small waves there are at most two instability bubbles. For a wave with approximate amplitude  $2.3 \times 10^{-2}$  we have the spectral magnitude plots of Figure 2. There we show the maximal value of the absolute value of the real part of a polynomial eigenvalue for various values of  $\mu$  near the predicted bifurcation points,  $\mu \sim 0.207$  and  $\mu \sim 0.368$ . In both cases the range of  $\mu$  values for which there is an instability is  $\mathcal{O}(10^{-3})$ .

We conclude by showing plots of the Krein eigenvalues for the situation in the right panel,  $\mu \sim 0.36$ . In Figure 3 we see a plot of the Krein eigenvalues for  $\mu \sim 0.368$ . The panel on the left shows the plot for the trivial state, and the panel on the right shows the plot for a small wave. Since this value of  $\mu$  is not associated with an instability (see right panel of Figure 2), the zeros of the Krein eigenvalues are purely real. One of the zeros corresponds to a polynomial eigenvalue with negative Krein index. In Figure 4 we see a plot of the Krein eigenvalues for  $\mu = 0.3585$ . The panel on the left shows the plot for the trivial state, and the panel on the right shows the plot for a small wave. Here there is not a zero/pole collision for the Krein eigenvalues. Since this value of  $\mu$  is associated with an instability (see right panel of Figure 2), the zeros of the Krein eigenvalues are purely real.

**5. Application: location of small eigenvalues.** The goal here is to use the Krein matrix to locate small polynomial eigenvalues. We start by assuming that the operator  $\mathcal{A}_0$  has a collection of arbitrarily small eigenvalues. These eigenvalues may arise, e.g., when looking at,

- (a) modulational stability problems for spatially periodic waves
- (b) sideband stability problems for uni-directional waves
- (c) interaction stability problems for multi-pulses.

*Assumption 5.1.* For each  $\epsilon > 0$  there exist  $N$  eigenvalues of  $\mathcal{A}_0 = \mathcal{A}_0(\epsilon)$ , say  $\mu_1, \dots, \mu_N$ , which satisfy  $|\mu_j| < \epsilon$ . The number  $N$  is independent of  $\epsilon$ . Moreover, there exists a positive constant  $C$ , independent of  $\epsilon$ , such that all other eigenvalues of  $\mathcal{A}_0$  satisfy  $|\mu| > C$ .

We will let  $s_1, \dots, s_N$  be the normalized set of associated eigenfunctions,

$$\mathcal{A}_0 s_j = \mu_j s_j, \quad \langle s_j, s_k \rangle = \delta_{jk},$$

and the subspace  $S$  used in the construction of the Krein matrix will be spectral subspace,  $S = \text{span}\{s_1, \dots, s_N\}$ . Letting  $P_S$  represent the spectral projection for  $\mathcal{A}_0$ , we have,

$$P_S \mathcal{A}_0 = \mathcal{A}_0 P_S, \quad P_{S^\perp} \mathcal{A}_0 = \mathcal{A}_0 P_{S^\perp}.$$

The Krein matrix,  $\mathbf{K}_S(z)$  for  $z = -i\lambda$ , associated with this subspace is given in Theorem 3.1, and the eigenvalues for the star-even operator are found by solving,

$$(5.1) \quad \mathbf{K}_S(z) \mathbf{x} = 0.$$

We start with a preliminary result concerning the part of the Krein matrix which generates poles.

LEMMA 5.2. *There exists a constant  $C_0 > 0$ , independent of  $\epsilon$ , such that for  $n = 1, 2$  and  $|z| < 1/C_0$ ,  $P_{S^\perp} P_n(i z) P_{S^\perp}$  is invertible. Moreover, for  $|z|$  sufficiently small there is the expansion,*

$$(P_{S^\perp} P_n(i z) P_{S^\perp})^{-1} = [\mathcal{I} + \mathcal{O}(|z|)] (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1}.$$

*Proof.* First suppose  $n = 1$ . Because  $P_{S^\perp}$  is a spectral projection,

$$P_{S^\perp} P_1(i z) P_{S^\perp} = P_{S^\perp} \mathcal{A}_0 P_{S^\perp} \left[ \mathcal{I} + z (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1) P_{S^\perp} \right].$$

The operator  $P_{S^\perp} \mathcal{A}_0 P_{S^\perp}$  is invertible with bounded inverse, as  $S$  is a spectral subspace associated with the small eigenvalues. Since  $(P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1) P_{S^\perp}$  is a compact operator, it too is uniformly bounded. Setting,

$$C_0 = \| (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1) P_{S^\perp} \|,$$

the operator  $\mathcal{I} + z (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1) P_{S^\perp}$  is invertible for  $|z| < 1/C_0$ . Moreover, a first-order Taylor expansion provides,

$$\left( \mathcal{I} + z (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1) P_{S^\perp} \right)^{-1} = \mathcal{I} + \mathcal{O}(|z|).$$

Taking the inverse yields the desired result.

If  $n = 2$  a similar argument gives the same result once one writes,

$$P_{S^\perp} P_2(i z) P_{S^\perp} = P_{S^\perp} \mathcal{A}_0 P_{S^\perp} \left[ \mathcal{I} + z (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i \mathcal{A}_1 - z \mathcal{A}_2) P_{S^\perp} \right],$$

and then notes that by assumption  $(P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_2 P_{S^\perp}$  is also compact.  $\square$

Since  $P_{S^\perp} P_n(i z) P_{S^\perp}$  is invertible for small  $z$ , we know through the argument in [subsection 2.1](#) that the following holds:

COROLLARY 5.3.  $\lambda_0$  is a small polynomial eigenvalue if and only if  $\det \mathbf{K}_S(z_0) = 0$  for  $z_0 = -i \lambda_0$ .

We now use the result of [Lemma 5.2](#) to find an approximation of the Krein matrix for small  $z$ .

LEMMA 5.4. *Suppose that  $n = 1$ . The Krein matrix is analytic for  $|z| < C_0$ . Moreover, if  $|z|$  is sufficiently small the Krein matrix has the expansion,*

$$\begin{aligned} \mathbf{K}_S(z) = & -z \left[ \text{diag}(\mu_1, \dots, \mu_N) + \bar{z} (i \mathcal{A}_1|_S) \right. \\ & \left. - \bar{z}^2 \left\{ -\mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1|_S \right\} + \mathcal{O}(|z|^3) \right]. \end{aligned}$$

*Proof.* Analyticity follows from the fact that  $P_{S^\perp} P_1(i z) P_{S^\perp}$  is invertible for  $|z| < C_0$ . Regarding the expansion, we first note that for the first term in the Krein matrix,

$$(\mathcal{P}_1(i z)|_S)_{jk} = \langle s_j, [\mathcal{A}_0 + z(i \mathcal{A}_1)] s_k \rangle = \mu_k \langle s_j, s_k \rangle + \bar{z} \langle s_j, (i \mathcal{A}_1) s_k \rangle,$$

so upon using the fact the eigenfunctions for  $\mathcal{A}_0$  form an orthonormal basis,

$$\mathcal{P}_1(i z)|_S = \text{diag}(\mu_1, \dots, \mu_N) + \bar{z} (i \mathcal{A}_1|_S).$$

Regarding the second term of the Krein matrix, first recall that we saw in the proof of [Lemma 5.2](#) that for small  $|z|$ ,

$$\begin{aligned} P_{S^\perp} P_1(iz) P_{S^\perp} &= P_{S^\perp} \mathcal{A}_0 P_{S^\perp} \left[ \mathcal{I} + z (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i\mathcal{A}_1) P_{S^\perp} \right] \\ &= P_{S^\perp} \mathcal{A}_0 P_{S^\perp} [\mathcal{I} + \mathcal{O}(|z|)], \end{aligned}$$

so upon using a Taylor expansion in  $z$ ,

$$(P_{S^\perp} P_1(iz) P_{S^\perp})^{-1} = [\mathcal{I} + \mathcal{O}(|z|)] (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1}.$$

Second, since  $P_{S^\perp}$  is a spectral projection, for any  $s \in S$ ,

$$P_{S^\perp} \mathcal{P}_1(iz) s = z P_{S^\perp} (i\mathcal{A}_1) s.$$

Combining these two facts,

$$\begin{aligned} &(\mathcal{P}_1(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_1(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_1(iz)|_S)_{jk} \\ &= \langle s_j, \mathcal{P}_1(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_1(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_1(iz) s_k \rangle \\ &= \langle P_{S^\perp} \mathcal{P}_1(-i\bar{z})^a s_j, (P_{S^\perp} \mathcal{P}_1(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_1(iz) s_k \rangle \\ &= \langle \bar{z} P_{S^\perp} (i\mathcal{A}_1) s_j, z [\mathcal{I} + \mathcal{O}(|z|)] (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i\mathcal{A}_1) s_k \rangle \\ &= \bar{z}^2 \langle s_j, (i\mathcal{A}_1) P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i\mathcal{A}_1) s_k \rangle + \mathcal{O}(|z|^3), \end{aligned}$$

which provides,

$$\begin{aligned} &\mathcal{P}_1(iz) P_{S^\perp} (P_{S^\perp} \mathcal{P}_1(iz) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{P}_1(iz)|_S = \\ &\bar{z}^2 (i\mathcal{A}_1) P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} (i\mathcal{A}_1)|_S + \mathcal{O}(|z|^3). \end{aligned}$$

The final result follows upon combining the above two calculations.  $\square$

Upon setting  $\gamma = i\bar{z}$  the bracketed part of the Krein matrix is approximated by a quadratic star-even polynomial matrix,

$$\text{diag}(\mu_1, \dots, \mu_N) + \gamma (\mathcal{A}_1|_S) + \gamma^2 \left[ -\mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1|_S \right].$$

Since  $|\mu_j| = \mathcal{O}(\epsilon)$ , the polynomial eigenvalues for this matrix will be  $\mathcal{O}(\epsilon^{1/2})$ ; consequently, the smallness assumption of [Lemma 5.2](#) regarding the polynomial eigenvalues is satisfied. Moreover, to leading order the polynomial eigenvalues are found by ignoring the middle term, so the small polynomial eigenvalues are found by solving the generalized linear eigenvalue problem, (5.2)

$$\text{diag}(\mu_1, \dots, \mu_N) \mathbf{v} = \alpha \left[ -\mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1|_S \right] \mathbf{v}, \quad \alpha = -\gamma^2 = \bar{z}^2.$$

In conclusion, the  $N$  small eigenvalues for  $\mathcal{A}_0$  will generate  $2N$  small polynomial eigenvalues, and to leading order these small polynomial eigenvalues are realized as the eigenvalues for the generalized eigenvalue problem (5.2). Since  $\det \mathbf{K}_S(\gamma)$  is analytic, and the winding number is invariant under small perturbations, the result is robust; in other words, we can conclude that there will be precisely  $2N$  small polynomial eigenvalues for  $\mathcal{P}_1(iz)$ , and these polynomial eigenvalues will be  $\mathcal{O}(\epsilon^{1/2})$ .

*Remark 5.5.* If  $S = \ker(\mathcal{A}_0)$ , then under the assumption  $\mathcal{A}_1|_{\ker(\mathcal{A}_0)}$  is the zero matrix,

$$-\mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1|_S = -\mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1|_{\ker(\mathcal{A}_0)},$$

which is precisely the constraint matrix associated with the Hamiltonian-Krein index calculation for linear star-even problems, see equation (1.1).

If  $n = 2$ , then an argument similar to that provided for Lemma 5.4 provides the approximate Krein matrix for small  $|z|$ . The details of the proof will be left for the interested reader.

LEMMA 5.6. *Suppose that  $n = 2$ . If  $|z|$  is sufficiently small the Krein matrix can be written,*

$$\begin{aligned} \mathbf{K}_S(z) = & -z \left[ \text{diag}(\mu_1, \dots, \mu_N) + \bar{z} (i\mathcal{A}_1|_S) \right. \\ & \left. - \bar{z}^2 \left( \mathcal{A}_2 - \mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1 \right)|_S + \mathcal{O}(|z|^3) \right]. \end{aligned}$$

*Remark 5.7.* If  $S = \ker(\mathcal{A}_0)$ , then under the assumption  $\mathcal{A}_1|_{\ker(\mathcal{A}_0)}$  is the zero matrix,

$$\left( \mathcal{A}_2 - \mathcal{A}_1 P_{S^\perp} (P_{S^\perp} \mathcal{A}_0 P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1 \right)|_S = \left( \mathcal{A}_2 - \mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1 \right)|_{\ker(\mathcal{A}_0)},$$

which is precisely the constraint matrix associated with the Hamiltonian-Krein index calculation for quadratic star-even problems, see equation (1.2).

**6. Example: suspension bridge equation.** Motivated by observations of traveling waves on suspension bridges, McKenna and Walter [26] proposed the model,

$$(6.1) \quad \partial_t^2 u + \partial_x^4 u + u^+ - 1 = 0,$$

to describe waves propagating on an infinitely long suspended beam, where  $u^+ = \max(u, 0)$ . To reduce the complexity due to the nonsmooth term  $u^+$ , Chen and McKenna [6] introduced the regularized equation,

$$(6.2) \quad \partial_t^2 u + \partial_x^4 u + e^{u-1} - 1 = 0.$$

Making the change of variables  $u - 1 \mapsto u$  in (6.2), so that localized solutions will decay to a baseline of 0, we will consider the equation,

$$(6.3) \quad \partial_t^2 u + \partial_x^4 u + e^u - 1 = 0.$$

Writing this in a co-moving frame with speed  $c$  by letting  $\xi = x - ct$ , equation (6.3) becomes

$$(6.4) \quad \partial_t^2 u - 2c\partial_{xt}^2 u + \partial_x^4 u + c^2\partial_x^2 u + e^u - 1 = 0,$$

where we have renamed the independent variable back to  $x$ .

An equilibrium solution to (6.4) satisfies the ODE,

$$(6.5) \quad \partial_x^4 u + c^2\partial_x^2 u + e^u - 1 = 0.$$

Smets and van den Berg [34, Theorem 11] prove the existence of a localized, symmetric solution  $U(x)$  to (6.5) for almost all wavespeeds  $c \in (0, \sqrt{2})$ . van den Berg et.al. [36, Theorem 1] use a computer-assisted proof technique to show existence of such solutions to (6.5) for all speeds  $c$  with  $c^2 \in [0.5, 1.9]$ . We take the following hypothesis.

*Hypothesis 6.1.* An exponentially localized primary pulse solution  $U(x; c)$  exists to (6.5) for each wavespeed  $c \in (0, \sqrt{2})$  and is smooth in the wavespeed.

For the remainder of this section, we will fix  $c \in (0, \sqrt{2})$  and write the primary pulse solution corresponding to wavespeed  $c$  as  $U(x)$ . We are interested in the existence and stability of multi-pulse equilibrium solutions to (6.4). A multi-pulse is a localized, multi-modal solution  $U_n(x)$  to (6.5) which resembles multiple, well-separated copies of the primary pulse  $U(x)$ .

**6.1. Existence of pulses.** First, we look at the existence of such pulses. The linearization of (6.5) about a given solution  $U_*$  of (6.5) is,

$$(6.6) \quad \mathcal{A}_0(U_*) = \partial_x^4 + c^2 \partial_x^2 + e^{U_*}.$$

We take the following hypothesis concerning  $\mathcal{A}_0(U)$ .

*Hypothesis 6.2.* On the space  $L^2(\mathbb{R})$ , the point spectrum of  $\mathcal{A}_0(U)$  consists of exactly two eigenvalues:

- (a)  $\mathcal{A}_0(U)$  has a one-dimensional kernel spanned by  $\partial_x U(x)$ .
- (b)  $n[\mathcal{A}_0(U)] = 1$ , i.e.  $\mathcal{A}_0(U)$  has a unique, simple negative eigenvalue  $\lambda_-$ .

*Remark 6.3.* While the results below hold if we only make assumptions on the nonnegative point spectrum of  $\mathcal{A}_0(U)$ , Hypothesis 6.2 is supported by numerical evidence. For simplicity, we make assumptions on the entire point spectrum of  $\mathcal{A}_0(U)$ .

From this hypothesis, it follows that the primary pulse  $U(x)$  is transversely constructed in the ODE phase space. Before stating our theorem, we will also need the following result. For  $c \in (0, \sqrt{2})$ , the solutions to the constant-coefficient equation,

$$\mathcal{A}_0(0)v = \partial_x^4 v + c^2 \partial_x^2 v + 1 = 0,$$

are given by  $v = e^{\mu x}$ , where

$$(6.7) \quad \mu = \pm \sqrt{\frac{-c^2 \pm \sqrt{c^4 - 4}}{2}} = \pm \alpha \pm i\beta,$$

for  $\alpha, \beta > 0$ . The constant  $\alpha$  is the exponential decay rate of localized solutions to (6.5). We now have the following theorem, which is adapted from [30, Theorem 3.6]. In all that follows, the norm  $\|\cdot\|_\infty$  is the supremum norm on  $C(\mathbb{R})$  and  $\langle \cdot, \cdot \rangle$  is the inner product on  $L^2(\mathbb{R})$ .

**THEOREM 6.4.** Assume Hypothesis 6.1 and Hypothesis 6.2. Fix a wavespeed  $c$ , and let  $U(x)$  be an exponentially localized solution to (6.5). Then for any  $n \geq 2$  and any sequence of nonnegative integers  $k_1, \dots, k_{n-1}$  with at least one of the  $k_j \in \{0, 1\}$ , there exists a nonnegative integer  $m_0$  and  $\delta > 0$  such that:

- (a) For any integer  $m$  with  $m \geq m_0$ , there exists a unique  $n$ -modal solution  $U_n(x)$  to (6.5) which is of the form

$$(6.8) \quad U_n(x) = \sum_{j=1}^n U^j(x) + r(x),$$

where each  $U^j(x)$  is a translate of the primary pulse  $U(x)$ . The distance between the peaks of  $U^j$  and  $U^{j+1}$  is  $2X_j$ , where

$$X_j \approx \frac{\pi}{\beta}(2m + k_j) + \tilde{X},$$



$\beta$  is defined in (6.7), and  $\tilde{X}$  is a constant. The remainder term  $r(x)$  satisfies

$$(6.9) \quad \|r\|_\infty \leq C e^{-\alpha X_{\min}},$$

where  $\alpha$  is defined in (6.7), and  $X_{\min} = \min\{X_1, \dots, X_{n-1}\}$ . This bound holds for all derivatives with respect to  $x$ .

(b) The point spectrum of the linear operator  $\mathcal{A}_0(U_n)$  on  $L^2(\mathbb{R})$  consists of exactly  $2n$  eigenvalues, which are as follows:

(1) There are  $n$  real eigenvalues  $\nu_1, \dots, \nu_n$  with  $|\nu_j| < \delta$ , where  $\nu_n = 0$  is a simple eigenvalue, and for  $j = 1, \dots, n-1$ ,

$$\begin{aligned} \nu_j &< 0 \text{ if } k_j \text{ is odd} \\ \nu_j &> 0 \text{ if } k_j \text{ is even.} \end{aligned}$$

We will refer to these as the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$ . For  $j = 1, \dots, n-1$ ,  $\nu_j = \mathcal{O}(e^{-2\alpha X_{\min}})$ , and the corresponding eigenfunctions  $s_j$  are given by

$$(6.10) \quad s_j = \sum_{k=1}^n d_{jk} \partial_x U^k + w_j,$$

where  $d_{jk} \in \mathbb{C}$  are constants, and the remainder terms  $w_j$  satisfy

$$(6.11) \quad \|w_j\|_\infty \leq C e^{-2\alpha X_{\min}}.$$

This bound holds for all derivatives with respect to  $x$ . The eigenfunction corresponding to  $\nu_n$  is  $s_n = \partial_x U_n$ .

(2) There are  $n$  negative eigenvalues which are  $\delta$ -close to  $\lambda_-$ .

(c) The essential spectrum of  $\mathcal{A}_0(U)$  is

$$(6.12) \quad \sigma_{\text{ess}}(\mathcal{A}_0(U)) = [1 - c^4/4, \infty).$$

which is positive and bounded away from 0.

*Proof.* Equation (6.5) is Hamiltonian with energy

$$(6.13) \quad H(u) = (\partial_x u) (\partial_x^4 u) - \frac{1}{2} (\partial_x^2 u)^2 + \frac{c^2}{2} (\partial_x u)^2 + e^u - u.$$

Using (6.7), (6.13), the assumption that the kernel is simple, and the fact that the Melnikov integral  $M = \int_{-\infty}^{\infty} (\partial_x U)^2 dx$  is positive, (a) follows from [30, Theorem 3.6], except for the bound on  $r(x)$ , which follows from [29] and [31]. All eigenvalues are real since  $\mathcal{A}_0(U_n)$  is self-adjoint on  $L^2(\mathbb{R})$ . From Hypothesis 6.2,  $\mathcal{A}_0(U)$  has a simple eigenvalue at 0 and a simple negative eigenvalue at  $\lambda_-$ . It follows from [1] that  $\mathcal{A}_0(U_n)$  has exactly  $n$  eigenvalues near each eigenvalue of  $\mathcal{A}_0(U)$ , i.e.  $n$  eigenvalues near 0 and  $n$  negative eigenvalues near  $\lambda_-$ . This proves the eigenvalue count and part (b2). Part (b1) follows from [31]. We can verify directly that  $\mathcal{A}_0(U_n) \partial_x U_n = 0$ . Part (c) follows from the Weyl Essential Spectrum Theorem [20, Theorem 2.2.6] and [20, Theorem 3.1.11], since  $\mathcal{A}_0(U_n)$  is exponentially asymptotic to  $\mathcal{A}_0(0)$ .  $\square$

**6.2. Stability of pulses.** Now that we know about the existence of single and multiple pulses, we consider their spectral stability. To determine linear PDE stability of the multi-pulse solutions constructed in Theorem 6.4, we look at the linearization

of the PDE (6.4) about  $U_n(x)$ , which is the quadratic operator polynomial  $\mathcal{P}_2(\lambda; U_n) : H^4(\mathbb{R}, \mathbb{C}) \subset L^2(\mathbb{R}, \mathbb{C}) \rightarrow L^2(\mathbb{R}, \mathbb{C})$  given by

$$(6.14) \quad \mathcal{P}_2(\lambda; U_n) = \mathcal{I}\lambda^2 + \mathcal{A}_1\lambda + \mathcal{A}_0(U_n)$$

where  $\mathcal{A}_0(U_n)$  is defined in (6.6),  $\mathcal{I}$  refers to the identity, and  $\mathcal{A}_1 = -2c\partial_x$ .

First, we consider the essential spectrum. Since  $U_n$  is exponentially localized,  $\mathcal{P}_2(\lambda; U_n)$  is exponentially asymptotic to the operator

$$(6.15) \quad \mathcal{P}_2(\lambda; 0) = \partial_x^4 + c^2\partial_x^2 - 2c\lambda\partial_x + (\lambda^2 + 1).$$

By [20, Theorem 3.1.11],  $\mathcal{P}_2(\lambda; U_n)$  is a relatively compact perturbation of  $\mathcal{P}_2(\lambda; 0)$ , thus by the Weyl essential spectrum theorem [20, Theorem 2.2.6],  $\mathcal{P}_2(\lambda; U_n)$  and  $\mathcal{P}_2(\lambda; 0)$  have the same essential spectrum. To find the essential spectrum of  $\mathcal{P}_2(\lambda; 0)$ , consider the related first-order operator  $\mathcal{T}(\lambda) : H^1(\mathbb{R}, \mathbb{C}^4) \subset L^2(\mathbb{R}, \mathbb{C}^4) \rightarrow L^2(\mathbb{R}, \mathbb{C}^4)$  given by

$$(6.16) \quad \mathcal{T}(\lambda) = \frac{d}{dx} - \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 - \lambda^2 & 2c\lambda & -c^2 & 0 \end{pmatrix},$$

which we obtain by writing  $\mathcal{P}_2(\lambda; 0)$  as a first order system. By a straightforward adaptation of [33, Theorem A.1] (the only difference being the presence of the fourth-order differential operator), the operators  $\mathcal{T}(\lambda)$  and  $\mathcal{P}_2(\lambda; 0)$  have the same Fredholm properties, thus the same essential spectrum. By a straightforward calculation,

$$(6.17) \quad \sigma_{\text{ess}}(\mathcal{P}_2(\lambda; U_n)) = \sigma_{\text{ess}}(\mathcal{T}(\lambda)) = \{ir : |r| \geq \rho\},$$

where  $\rho > 0$  is the minimum of the function  $\lambda(r) = cr + \sqrt{1 + r^4}$ . The value of  $\rho$  is positive for  $c \in (0, \sqrt{2})$ , and  $\rho \rightarrow 0$  as  $c \rightarrow \sqrt{2}$ , so the essential spectrum is purely imaginary and bounded away from 0. Spectral stability thus depends entirely on the point spectrum.

**6.2.1. Single pulse.** Before considering the spectral stability of the  $n$ -pulse, we must show the stability of the primary pulse,  $U(x)$ . In addition to Hypothesis 6.1 and Hypothesis 6.2, our assumptions are:

*Hypothesis 6.5.* Regarding the PDE (6.4) and the base solution  $U(x)$ ,  
 (a) for every initial condition  $u(x, 0)$  and  $\partial_t u(x, 0)$  there exists a solution  $u(x, t)$  to (6.4) on the interval  $I = [0, T]$ , where

$$(6.18) \quad T = T(\max\{\|u(x, 0)\|, \|\partial_t u(x, 0)\|\})$$

(b) the constrained energy evaluated on the wave,  $d(c)$  (see [9, Equation (2.16)] for the exact expression), is concave up,

$$(6.18) \quad d''(c) = -\partial_c(c\|\partial_x U\|^2) > 0, \quad 0 < c^2 < 2.$$

We will provide numerical evidence that these hypotheses are met in subsection 6.3.

Under these assumptions, we will prove the spectral and orbital stability of the single pulse using the HKI. However, there are first two issues that must be resolved. First, the HKI as discussed in section 1 assumes that  $\mathcal{A}_0$  has a compact resolvent,

which is certainly not true for the operator associated with this problem. This compactness assumption is taken primarily for the sake of convenience, and to remove the possibility of point spectrum being embedded in the essential spectrum. However, as seen in the original formulation of the HKI for solitary waves, see [16, 17], this is not a necessary condition. It is sufficient to assume that the origin is an isolated eigenvalue, and  $\mathcal{A}_0$  is a higher-order differential operator than  $\mathcal{A}_1$  with  $n[\mathcal{A}_0] < +\infty$ . The interested reader should consult [21] for the case where the origin is not isolated. The second difficulty is that these previous results for solitary waves do not immediately apply to quadratic eigenvalue problems. However, as seen in [2, Section 4.1] one can easily convert a quadratic star-even eigenvalue problem into a linear star-even eigenvalue problem, and then apply the index theory to the reformulated problem. Thus, we can conclude the index theory is applicable to the problem at hand, which allows for the following stability result.

LEMMA 6.6. *Let  $c^2 \in (0, 2)$ , and let  $U(x)$  be the primary pulse solution to (6.5). Then  $U(x)$  is spectrally and orbitally stable if and only if*

$$d''(c) = -\partial_c (c \|\partial_x U\|^2) > 0,$$

where  $d(c)$  is defined in [9, equation (2.16)].

*Proof.* We first check that the origin is an isolated eigenvalue. The essential spectrum of  $\mathcal{A}_0(U)$  is the same as that of  $\mathcal{A}_0(U_n)$ , and is given by (6.12), which is positive and bounded away from 0. By assumption,  $\mathcal{A}_0(U)$  has a single negative eigenvalue.

We now use the HKI to complete the proof; in particular, the formulation as presented in equation (1.2). First, we note that,

$$\mathcal{A}_1|_{\text{span}\{\partial_x U\}} = \langle -2c\partial_x (\partial_x U), \partial_x U \rangle = 0,$$

where the equality follows from the fact that the primary pulse is even. Since  $\mathcal{A}_2 = \mathcal{I}$  is positive definite, we can write,

$$\begin{aligned} K_{\text{Ham}} &= n(\mathcal{A}_0) - n\left(\left[\mathcal{I} - \mathcal{A}_1\mathcal{A}_0^{-1}\mathcal{A}_1\right]|_{\text{span}\{\partial_x U\}}\right) \\ &= 1 - n\left(\left[\mathcal{I} - \mathcal{A}_1\mathcal{A}_0^{-1}\mathcal{A}_1\right]|_{\text{span}\{\partial_x U\}}\right), \end{aligned}$$

for by assumption,  $n(\mathcal{A}_0) = 1$ .

Regarding the second term,

$$\begin{aligned} \left[\mathcal{I} - \mathcal{A}_1\mathcal{A}_0^{-1}\mathcal{A}_1\right]|_{\text{span}\{\partial_x U\}} &= \|\partial_x U\|^2 - \langle (-2c\partial_x)\mathcal{A}_0^{-1}(-2c\partial_x)\partial_x U, \partial_x U \rangle \\ &= \|\partial_x U\|^2 + 2c\langle \partial_x \mathcal{A}_0^{-1}(-2c\partial_x^2 U), \partial_x U \rangle. \end{aligned}$$

Going back to the existence equation (6.5) and differentiating with respect to  $c$  yields,

$$\mathcal{A}_0(U)\partial_c U + 2c\partial_x^2 U = 0 \rightsquigarrow \mathcal{A}_0(U)^{-1}(-2c\partial_x^2 U) = \partial_c U.$$

Substitution and changing the order of differentiation provides,

$$\langle \partial_x \mathcal{A}_0(U)^{-1}(-2c\partial_x^2 U), \partial_x U \rangle = \langle \partial_c \partial_x U, \partial_x U \rangle = \frac{1}{2} \partial_c \|\partial_x U\|^2.$$

In conclusion,

$$\left[\mathcal{I} - \mathcal{A}_1\mathcal{A}_0^{-1}\mathcal{A}_1\right]|_{\text{span}\{\partial_x U\}} = \|\partial_x U\|^2 + c \partial_c \|\partial_x U\|^2 = \partial_c (c \|\partial_x U\|^2).$$

We now have for the primary pulse,

$$K_{\text{Ham}} = 1 - n [\partial_c (c \|\partial_x U\|^2)].$$

If  $d''(c) < 0$ , then  $K_{\text{Ham}} = 1$ , and there is one positive real polynomial eigenvalue. If  $d''(c) > 0$ , the HKI is zero. Consequently, the wave is spectrally stable. Appealing to [2, Theorem 4.1] we can further state that the wave is orbitally stable.  $\square$

**6.2.2.  $n$ -pulse.** We now locate all potentially unstable eigenvalues of (6.14) for an  $n$ -pulse. These include polynomial eigenvalues with positive real part, as well as purely imaginary polynomial eigenvalues with negative Krein signature. To accomplish this task we use the HKI in combination with the Krein matrix. First, we compute the HKI for (6.14), so that we have an exact count of the number of potentially unstable polynomial eigenvalues. We then use the Krein matrix to find  $(n - 1)$  pairs of eigenvalues close to 0; each pair is either real or purely imaginary with negative Krein signature. We refer to these as small magnitude polynomial eigenvalues, or interaction polynomial eigenvalues, since heuristically they result from interactions between neighboring pulses. We then show that the number of potentially unstable interaction polynomial eigenvalues is exactly the same as the HKI, from which we conclude that we have found all of the potentially unstable eigenvalues. By Hamiltonian reflection symmetry, all other point spectrum must be purely imaginary with positive Krein signature.

We start with the calculation of the HKI. By Theorem 6.4 we know that  $\mathcal{A}_0(U_n)$  has precisely  $n$  eigenvalues near the origin. Let  $0 \leq n_s \leq n - 1$  represent the number of these eigenvalues which are negative. We have the following result concerning the HKI for the  $n$ -pulse:

LEMMA 6.7. *Assume Hypotheses 6.1, 6.2, and 6.5, and let  $U_n(x)$  be an  $n$ -modal solution to (6.5). Then*

$$K_{\text{Ham}} = n + n_s - 1.$$

*Proof.* From Theorem 6.4 part (b) and the definition of  $n_s$ ,  $n[\mathcal{A}_0(U_n)] = n + n_s$ , so for the HKI,

$$K_{\text{Ham}} = n + n_s - n \left( [\mathcal{I} - \mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1] \big|_{\text{span}\{\partial_x U_n\}} \right),$$

where  $\mathcal{A}_0 = \mathcal{A}_0(U_n)$ . In the proof of Lemma 6.6 we saw that when the wave depends smoothly on  $c$ ,

$$[\mathcal{I} - \mathcal{A}_1 \mathcal{A}_0^{-1} \mathcal{A}_1] \big|_{\text{span}\{\partial_x U_n\}} = \partial_c (c \|\partial_x U_n\|^2).$$

Since to leading order the  $n$ -pulse is  $n$  copies of the original pulse, we have

$$\|\partial_x U_n\|^2 = n \|\partial_x U\|^2 + \mathcal{O}(e^{-\alpha X_{\min}}).$$

Consequently, we can write

$$\begin{aligned} \partial_c (c \|\partial_x U_n\|^2) &= n \partial_c (c \|\partial_x U\|^2) + \mathcal{O}(e^{-\alpha X_{\min}}) \\ &= -n d''(c) + \mathcal{O}(e^{-\alpha X_{\min}}). \end{aligned}$$

Since  $d''(c) > 0$  by assumption, we have to leading order,

$$\partial_c (c \|\partial_x U_n\|^2) < 0.$$

For sufficiently well-separated pulses the sign will not change even when incorporating the higher-order terms in the asymptotic expansion. The result now follows.  $\square$

We now locate the potentially unstable polynomial eigenvalues of the quadratic eigenvalue problem (6.14). This will be accomplished through the Krein matrix. For the sake of exposition only we will henceforth assume that each of the small magnitude eigenvalues  $\nu_1, \dots, \nu_n$  of  $\mathcal{A}_0(U_n)$  is simple. For each of these eigenvalues, denote the associated normalized eigenfunctions as  $s_1, \dots, s_n$ . Since  $\mathcal{A}_0(U_n)$  is self-adjoint, these eigenfunctions are pairwise orthogonal. In the construction of the Krein matrix the relevant subspace for the spectral problem is the span of this set of eigenfunctions associated with the small magnitude eigenvalues of  $\mathcal{A}_0$ ,

$$(6.19) \quad S = \text{span}\{s_1, \dots, s_n\}.$$

We now present the following theorem, which is the main result of this section.

**THEOREM 6.8.** *Assume Hypotheses 6.1, 6.2, and 6.5. Let  $U_n(x)$  be an  $n$ -pulse solution to (6.5), and let  $\nu_1, \dots, \nu_n$  be the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$ , as defined in Theorem 6.4. Under a suitable normalization of the eigenfunctions  $s_j$ , near the origin the Krein matrix has the asymptotic expansion,*

$$(6.20) \quad -\frac{K_S(z)}{z} = \|\partial_x U\|^2 \text{diag}(\nu_1, \dots, \nu_n) + d''(c) \mathbf{I}_n \bar{z}^2 + \mathcal{O}(e^{-(3\alpha/2)X_{\min}} |z| + |z|^3),$$

which is diagonal to leading order.

The proof of this result is left to subsection 6.4. As a corollary, we have the following criteria for spectral stability and instability of the multi-pulse solutions  $U_n(x)$ .

**COROLLARY 6.9.** *Let  $U_n(x)$  be an  $n$ -pulse solution to (6.5) constructed as in Theorem 6.4 using the sequence of nonnegative integers  $\{k_1, \dots, k_{n-1}\}$ . Assume the same hypotheses as in Theorem 6.8. Let  $\nu_1, \dots, \nu_n$  be the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$ , where  $\nu_n = 0$ . Then there are  $(n-1)$  pairs of eigenvalues of (6.14) close to 0, which we will term interaction polynomial eigenvalues. These are described as follows. For each  $j = 1, 2, \dots, n-1$ ,*

(a) *if  $k_j$  is odd (equivalently,  $\nu_j < 0$ ), there is a corresponding pair of purely imaginary interaction polynomial eigenvalues,*

$$(6.21) \quad \lambda_j^\pm = \pm i \left( \|\partial_x U\| \sqrt{\frac{|\nu_j|}{d''(c)}} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}) \right),$$

*each of which has negative Krein signature*

(b) *if  $k_j$  is even (equivalently,  $\nu_j > 0$ ), there is a corresponding pair of real interaction polynomial eigenvalues,*

$$\lambda_j = \pm \left( \|\partial_x U\| \sqrt{\frac{\nu_j}{d''(c)}} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}) \right).$$

*In particular, there exists a positive, real eigenvalue.*

*In addition, there is a geometrically simple polynomial eigenvalue at  $\lambda = 0$  with corresponding eigenfunction  $\partial_x U_n$ . All other point spectra is purely imaginary, and has positive Krein signature.*

**Remark 6.10.** In other words, if all the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$  are negative, and if the individual pulses are sufficiently well-separated, then the  $n$ -pulse is spectrally stable; otherwise, it is unstable.

While we can find the interaction polynomial eigenvalues using Lin's method as in [31], using the Krein matrix allows us to also determine the Krein signatures of any purely imaginary interaction polynomial eigenvalues. This additional information is needed to ensure that via the HKI all of the potentially unstable point spectrum has small magnitude.

*Proof.* By Corollary 5.3 the small polynomial eigenvalues are found by solving  $\det \mathbf{K}_S(z) = 0$ . This is equivalent to finding zeros of the Krein eigenvalues. For  $j = 1, 2, \dots, n$  set,

$$-\frac{r_j(z)}{z} = \|\partial_x U\|^2 \nu_j + d''(c)\bar{z}^2 + \tilde{r}_j(z),$$

where

$$\tilde{r}_j(z) = \mathcal{O}(e^{-(3\alpha/2)X_{\min}}|z| + |z|^3).$$

Note that the first two terms in  $-r_j(z)/z$  are the diagonal entries of the Krein matrix. Since to leading order the Krein matrix is diagonal, by [13] these are valid asymptotic expressions for the Krein eigenvalues. The small and nonzero polynomial eigenvalues are found by solving,

$$(6.22) \quad \|\partial_x U\|^2 \nu_j + d''(c)\bar{z}^2 + \tilde{r}_j(z) = 0, \quad j = 1, 2, \dots, n.$$

First suppose that  $z$  is real, so the Krein matrix is Hermitian. The Krein eigenvalues are then real-valued; in particular, the error term,  $\tilde{r}_j(z)$ , is real-valued. Recall that  $d''(c) > 0$ . Suppose that  $\nu_j < 0$ , and set,

$$(6.23) \quad \epsilon_j^2 = -\frac{\|\partial_x U\|^2 \nu_j}{d''(c)} > 0.$$

Equation (6.22) can then be rewritten,

$$(6.24) \quad z^2 - \epsilon_j^2 + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}|z| + |z|^3) = 0.$$

Letting  $y = \epsilon_j z$  and noting that  $\epsilon_j = \mathcal{O}(e^{-\alpha X_{\min}})$ , equation (6.24) becomes,

$$(6.25) \quad y^2 - 1 + \mathcal{O}(\epsilon_j^{1/2}|y| + \epsilon_j|y|^3) = 0.$$

For sufficiently small  $\epsilon_j$ , equation (6.25) has two roots,  $y = \pm 1 + \mathcal{O}(\epsilon_j^{1/2})$ . Thus, for sufficiently large  $X_{\min}$ , equation (6.22) has two solutions,

$$z_j^\pm = \pm \|\partial_x U\| \sqrt{-\frac{\nu_j}{d''(c)}} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

The Krein eigenvalue,  $r_j(z)$ , has a simple zero at  $z_j^\pm$ . Since to leading order,

$$r'_j(z_j^\pm) = -\|\partial_x U\|^2 \nu_j - 3d''(c)(z_j^\pm)^2 = 2\|\partial_x U\|^2 \nu_j < 0,$$

each of these polynomial eigenvalues has negative Krein signature.

Now suppose  $\nu_j > 0$ , and assume  $z$  is purely imaginary,  $z = i\tilde{z}$ . In this case the Krein matrix is no longer Hermitian, which implies that the remainder term associated with each Krein eigenvalue is no longer necessarily real-valued. Define  $\epsilon_j^2$  as in (6.23), but this time  $\epsilon_j^2 < 0$ . The two zeros of the Krein eigenvalue are now,

$$\tilde{z}_j^\pm = \pm \|\partial_x U\| \sqrt{\frac{\nu_j}{d''(c)}} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}),$$

which to leading order are purely real. Going back to the original problem, there are two interaction polynomial eigenvalues given by,

$$\lambda_j^\pm = \tilde{z}_j^\pm.$$

To leading order these eigenvalues are real-valued. Under the assumption that the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$  are simple, via the asymptotic expansion  $\lambda_j^\pm$  will also then be simple. By the Hamiltonian reflection symmetry of the polynomial eigenvalues about the real axis, the fact they are real-valued to leading order implies they are truly real-valued and come in opposite-sign pairs.

Since the kernels of (6.14) and  $\mathcal{A}_0(U_n)$  are the same, we can verify directly that  $\lambda = 0$  is an eigenvalue of (6.14) with eigenfunction  $\partial_x U_n$ . We now show that all other point spectra is purely imaginary. We have for the small magnitude polynomial eigenvalues,  $k_i^- = 2n_s$ , and  $k_r = n - 1 - n_s$ . Thus, for the small magnitude polynomial eigenvalues,

$$k_r + k_i^- = (n - 1 - n_s) + (2n_s) = n - 1 + n_s.$$

By Lemma 6.7 this is the HKI for the  $n$ -pulse. Consequently, there are no other point polynomial eigenvalues which have positive real part, or which are purely imaginary and have negative Krein signature.  $\square$

**6.3. Numerical results.** In this section, we show numerical results to illustrate the theoretical results of the previous section. First, we can construct a primary pulse solution  $U(x)$  numerically using the string method from [4]. The top two panels of Figure 5 show these solutions for the same values of  $c$  as in [6, Figure 3]. Next, we compute the spectrum of the operator  $\mathcal{A}_0(U)$  numerically using Matlab's `eig` function. In the bottom panel of Figure 5 we note the presence of a simple eigenvalue at the origin and a simple negative eigenvalue, which supports our hypotheses on the spectrum of  $\mathcal{A}_0(U)$ . As expected, we also see that the essential spectrum is positive and bounded away from 0.

We can construct multi-pulse solutions numerically by joining together multiple copies of the primary pulse and using Matlab's `fsolve` function. Consecutive distances between peaks given by Theorem 6.4. The first four double pulse solutions are shown in the top two panels of Figure 6. These double pulses are numbered using the integer  $k_1$  from Theorem 6.4. We verify Theorem 6.4(b) numerically by computing the spectrum of  $\mathcal{A}_0(U_2)$ . The spectrum of  $\mathcal{A}_0(U_2)$  for double pulses 0 and 1 are shown in the bottom two panels of Figure 6. In both cases, there is an eigenvalue at 0. For double pulse 0, there is an additional positive eigenvalue near 0, and for double pulse 1, there is an additional negative eigenvalue near 0.

We verify Corollary 6.9 by computing the polynomial eigenvalues of (6.14) directly using the Matlab package `quadeig` from [10]. For double pulse 0,  $\mathcal{A}_0(U_2)$  has one positive small magnitude eigenvalue; thus, by Corollary 6.9, equation (6.14) has a polynomial eigenvalue with positive real part. For double pulse 1, the small magnitude eigenvalue of  $\mathcal{A}_0(U_2)$  is negative; thus by Corollary 6.9, since the distance between the two peaks is sufficiently large, the polynomial eigenvalues of (6.14) are purely imaginary. These are shown in Figure 7.

**6.4. Proof of Theorem 6.8.** Using Theorem 6.4, let  $U_n(x)$  be an  $n$ -modal solution to (6.5), and let  $\{\nu_1, \dots, \nu_n\}$  be the small magnitude eigenvalues of  $\mathcal{A}_0(U_n)$  with corresponding eigenfunctions  $\{s_1, \dots, s_n\}$ . Since  $\mathcal{A}_0(U_n)$  is self-adjoint, the  $s_i$  are orthogonal, and for the sake of convenience scale them so that,

$$(6.26) \quad \langle s_i, s_j \rangle = \|\partial_x U\|^2 \delta_{ij}.$$






FIG. 5. Primary pulse solutions  $U(x)$  to (6.5) for  $c = 1.354$  (top left) and  $c = 1.40$  (top right). In the bottom panel there is the spectrum of  $\mathcal{A}_0(U)$ , the linearization of (6.5) about a single pulse  $U(x)$  for  $c = 1.3$ . For the spectral plot we use finite difference methods with  $N = 512$  and periodic boundary conditions. The left boundary of the essential spectrum is  $\lambda \sim 0.286$ . The spectrum to the right of the boundary is discrete instead of continuous because of the boundary conditions.

Typically, we assume these eigenfunctions also have unit length. However, this is not important in the construction of the Krein matrix, nor in the derived properties.

By Lemma 5.6, and using the normalization of (6.26), for small  $|z|$  the Krein matrix is the  $n \times n$  matrix,

$$(6.27) \quad -\frac{\mathbf{K}_S(z)}{z} = \|\partial_x U_n\|^2 \text{diag}(\nu_1, \dots, \nu_n) + \bar{z} \mathbf{K}_1 - \bar{z}^2 (\|\partial_x U_n\|^2 \mathbf{I}_n - \mathbf{K}_2) + \mathcal{O}(|z|^3),$$

where

$$(6.28) \quad (\mathbf{K}_1)_{jk} = \langle s_j, i\mathcal{A}_1 s_k \rangle,$$

and

$$(6.29) \quad (\mathbf{K}_2)_{jk} = \langle \mathcal{A}_1 s_j, P_{S^\perp} (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \mathcal{A}_1 s_k \rangle.$$

This is, to leading order, a matrix-valued quadratic polynomial in  $z$  (and its complex conjugate). We now prove Theorem 6.8 in a series of lemmas. In all that follows,  $C$  refers to a constant independent of  $x$ , but it may have a different value each time it is used. The first lemma is a bound on the product of exponentially separated pulses.

LEMMA 6.11. Let  $U_+(x)$  and  $U_-(x)$  be localized pulses which decay exponentially with rate  $\alpha$  and whose peaks are separated by a distance  $2X$ . We have the following






FIG. 6. Double pulse solutions  $U_2(x)$  to (6.5) for  $c = 1.2$ . The top left panel shows double pulse 0, and the top right panel shows double pulse 1. In the bottom two panels we see the associated spectrum for  $\mathcal{A}_0(U_2)$ : double pulse 0 on the left, and double pulse 1 on the right.

1087 bounds,

$$1088 \quad (6.30) \quad \sup_{x \in \mathbb{R}} |U_-(x)U_+(x)| \leq Ce^{-2\alpha X},$$

1089 and

$$1090 \quad (6.31) \quad |\langle U_-(x)U_+(x) \rangle| \leq Ce^{-(3\alpha/2)X}.$$

1091

1092 *Proof.* Without loss of generality, let  $U_{\pm}(x)$  be exponentially localized peaks  
 1093 centered at  $\pm X$ , thus  $|U_-(x)| \leq Ce^{-\alpha|x+X|}$  and  $|U_+(x)| \leq Ce^{-\alpha|x-X|}$ . For  $x \in$   
 1094  $(-\infty, -X]$ ,

$$1095 \quad (6.32) \quad |U_-(x)U_+(x)| \leq Ce^{\alpha(x+X)}e^{\alpha(x-X)} = Ce^{2\alpha x} \leq Ce^{-2\alpha X}$$

1097 and for  $x \in [-X, 0]$ ,

$$1098 \quad (6.33) \quad |U_-(x)U_+(x)| \leq Ce^{-\alpha(x+X)}e^{\alpha(x-X)} = Ce^{-2\alpha X}$$

1100 Bounds on  $[0, X]$  and  $[X, \infty)$  are similar. Since these are independent of  $x$ , we obtain  
 1101 the bound (6.30).




FIG. 7. Polynomial eigenvalues of (6.14) for double pulses 0 (left) and 1 (right) for  $c = 1.2$ . The eigenvalues are marked with a filled (blue) circle, and the edge of the essential spectrum is marked with a (red) cross. The essential spectrum is discrete instead of continuous because of the boundary conditions. For the right panel the two purely imaginary polynomial eigenvalues nearest the origin have negative Krein signature. Here we use finite difference methods with  $N = 512$  and periodic boundary conditions.

For the bound (6.31), we split the integral into four pieces.

$$\begin{aligned}
 |\langle U_-(x), U_+(x) \rangle| &\leq \int_{-\infty}^{-X} |U_-(x)U_+(x)|dx + \int_{-X}^0 |U_-(x)U_+(x)|dx \\
 &+ \int_0^X |U_-(x)U_+(x)|dx + \int_X^{\infty} |U_-(x)U_+(x)|dx
 \end{aligned}
 \tag{6.34}$$

For the first integral, we use (6.32) to get

$$\int_{-\infty}^{-X} |q_-(x)q_+(x)|dx \leq C \int_{-\infty}^{-X} e^{2\alpha x}dx = Ce^{-2\alpha X}$$

For the second integral, we use (6.33) to get

$$\begin{aligned}
 \int_{-X}^0 |q_-(x)q_+(x)|dx &\leq C \int_{-X}^0 e^{-\alpha(x+X)} e^{\alpha(x-X)}dx \leq C \int_{-X}^0 e^{-\alpha(x+X)/2} e^{\alpha(x-X)}dx \\
 &\leq Ce^{-(3\alpha/2)X} \int_{-X}^0 e^{(\alpha/2)x}dx \leq Ce^{-(3\alpha/2)X}
 \end{aligned}$$

The third and fourth integrals are similar. Combining these, we obtain (6.31).  $\square$

*Remark 6.12.* If the hypotheses of Lemma 6.11 are satisfied, we say that  $U_+(x)$  and  $U_-(x)$  are exponentially separated by  $2X$ .

Next, we obtain a bound on the matrix  $\mathbf{K}_1$ .

1115 LEMMA 6.13. For the matrix  $\mathbf{K}_1$  in (6.27),

1116 (6.35)  $\mathbf{K}_1 = \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$

1117 *Proof.* Substituting  $\mathcal{A}_1 = -2c\partial_x$  into (6.28),  $(\mathbf{K}_1)_{jk} = i2c\langle s_j, \partial_x s_k \rangle$ . Using the  
 1118 expansion (6.10) from Theorem 6.4,

1119 (6.36) 
$$\begin{aligned} \langle s_j, \partial_x s_k \rangle &= \sum_{m=1}^n d_{jm} d_{km} \langle \partial_x U_x^m, \partial_x^2 U^m \rangle + \sum_{m \neq \ell} d_{jm} d_{k\ell} \langle \partial_x U^m, \partial_x^2 U^\ell \rangle \\ &\quad + \langle s_j, \partial_x w_k \rangle + \sum_{\ell=1}^n d_{k\ell} \langle w_j, \partial_x^2 U^\ell \rangle. \end{aligned}$$

1120 By translation invariance of the inner product on  $L^2(\mathbb{R})$ ,

1121 
$$\langle \partial_x U^m, \partial_x^2 U^m \rangle = \langle \partial_x U, \partial_x(\partial_x U) \rangle = 0,$$

1122 since the operator  $\partial_x$  is skew-symmetric. For  $m \neq \ell$ ,  $U^m$  and  $U^\ell$  are exponentially  
 1123 separated by at least  $2X_{\min}$ ; thus, by Lemma 6.11,

1124 
$$\langle \partial_x U^m, \partial_x^2 U^\ell \rangle = \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

1125 The last two terms in (6.36) are  $\mathcal{O}(e^{-2\alpha X_{\min}})$  using Hölder's inequality and the bound  
 1126 (6.11) from Theorem 6.4, which applies to  $\partial_x w_k$  as well as  $w_j$ . Combining these  
 1127 estimates we obtain (6.35).  $\square$

1128 Using the expansion (6.10) from Theorem 6.4, the matrix  $\mathbf{K}_2$  in (6.27) becomes,  
 (6.37)

1129 
$$(\mathbf{K}_2)_{jk} = 4c^2 \left\langle \sum_{m=1}^n d_{jm} \partial_x^2 U^m + \partial_x w_j, \sum_{\ell=1}^n d_{k\ell} P_{S^\perp} (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x^2 U^\ell + P_{S^\perp} (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x w_k \right\rangle.$$

1130 Before we can evaluate this expression, we need to look at  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1}$ .

1131 LEMMA 6.14.  $P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp} : S^\perp \rightarrow S^\perp$  is an invertible linear operator with  
 1132 bounded inverse.

1133 *Proof.* By (6.12), the essential spectrum of  $\mathcal{A}_0(U_n)$  is bounded away from 0,  
 1134 thus the operator  $\mathcal{A}_0(U_n)$  is Fredholm with index 0. Since  $\mathcal{A}_0(U_n)$  is self-adjoint,  
 1135  $P_{S^\perp} \mathcal{A}_0(U_n)$  is also self-adjoint, and it is not hard to show that  $P_{S^\perp} \mathcal{A}_0(U_n)$  is Fredholm  
 1136 with index 0 and has kernel  $S$ . Thus the restriction  $P_{S^\perp} \mathcal{A}_0(U_n)|_{S^\perp} = P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp}$   
 1137 is invertible on  $S^\perp$ . By the definition of  $S$  and Theorem 6.4,  $P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp}$  has  
 1138 no eigenvalues of magnitude less than  $\delta$ . Since the essential spectrum of  $\mathcal{A}_0(U_n)$  is  
 1139 bounded away from 0, by the resolvent bound for normal operators, the linear operator  
 1140  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1}$  is bounded on  $S^\perp$ .  $\square$

1141 Before we can evaluate the term  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x^2 U^\ell$  from (6.37), we  
 1142 will need the following lemma which gives an expansion for  $e^{U_n(x)}$ .

1143 LEMMA 6.15. For the  $n$ -pulse,  $U_n(x)$ , and for all  $i = 1, \dots, n$ ,

1144 
$$\exp(U_n(x)) = \exp(U^i(x)) + \sum_{j \neq i} (\exp(U^j(x)) - 1) + \mathcal{O}(e^{-\alpha X_{\min}})$$

1145 *Proof.* Fix  $i$  in the expansion (6.8) and let  $S(x) = \sum_{j \neq i} U_j(x)$ , so that  $U_n =$   
 1146  $U^i + S + \mathcal{O}(e^{-\alpha X_{\min}})$ . Since  $U_n(x)$  is bounded,

$$\begin{aligned} \exp(U_n(x)) &= \exp(U^i(x)) \exp(S(x)) (1 + \mathcal{O}(e^{-\alpha X_{\min}})) \\ &= \exp(U^i(x)) \exp(S(x)) + \mathcal{O}(e^{-\alpha X_{\min}}). \end{aligned}$$

1148 Using the Taylor expansion for the exponential,

$$\begin{aligned} \exp(U^i(x)) \exp(S(x)) &= \sum_{m=0}^{\infty} \frac{U^i(x)^m}{m!} \sum_{n=0}^{\infty} \frac{S(x)^n}{n!} \\ &= \sum_{m=0}^{\infty} \frac{U^i(x)^m}{m!} + \sum_{n=0}^{\infty} \frac{S(x)^n}{n!} - 1 + \sum_{m=1}^{\infty} \frac{U^i(x)^m}{m!} \sum_{n=1}^{\infty} \frac{S(x)^n}{n!} \\ &= \exp(U^i(x)) + \exp(S(x)) - 1 + \sum_{m=1}^{\infty} \frac{U^i(x)^m}{m!} \sum_{n=1}^{\infty} \frac{S(x)^n}{n!} \end{aligned}$$

1150 For the last term on the RHS,

$$\begin{aligned} \left| \sum_{m=1}^{\infty} \frac{U^i(x)^m}{m!} \sum_{n=1}^{\infty} \frac{S(x)^n}{n!} \right| &= |U^i(x)S(x)| \sum_{m=0}^{\infty} \frac{|U^i(x)|^m}{(m+1)!} \sum_{n=0}^{\infty} \frac{|S(x)|^n}{(n+1)!} \\ &\leq |U^i(x)S(x)| e^{|U^i(x)|} e^{|S(x)|} \\ &\leq C e^{-2\alpha X_{\min}}, \end{aligned}$$

1152 where in the last line we used the fact that  $U_n(x)$  is bounded together with the bound  
 1153 (6.30) from Lemma 6.11, since  $U^i$  and each peak in  $S$  are exponentially separated.  
 1154 Combining all of this,

$$\exp(U_n(x)) = \exp(U^i(x)) + \exp(S(x)) - 1 + \mathcal{O}(e^{-\alpha X_{\min}})$$

1156 Repeat this procedure  $n - 2$  more times to get the result.  $\square$

1157 We can now evaluate  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x^2 U^\ell$ .

LEMMA 6.16.

$$(6.38) \quad (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x^2 U^\ell = -\frac{1}{2c} P_{S^\perp} \partial_c U^\ell + \mathcal{O}(e^{-2\alpha X_{\min}}).$$

1159 *Proof.* Let  $y = (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1} P_{S^\perp} \partial_x^2 U^\ell$ , which is well-defined by Lemma  
 1160 6.14. Since  $P_{S^\perp} \partial_x^2 U^\ell$  is smooth and  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1}$  is bounded,  $y$  is smooth as  
 1161 well and is the unique solution to the equation

$$(6.39) \quad (P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp}) y = P_{S^\perp} \partial_x^2 U^\ell.$$

1163 Using Lin's method as in [31], we will look for a solution to (6.39) of the form,

$$(6.40) \quad \tilde{y} = -\frac{1}{2c} P_{S^\perp} \partial_c U^\ell + \tilde{w}.$$

1165 This ansatz is suggested by

$$(6.41) \quad \mathcal{A}_0(U) \partial_c U = -2c \partial_x^2 U,$$

1167 which we obtain by taking  $u = U$  in equation (6.5) and differentiating with respect  
 1168 to  $c$ , which we can do since  $U$  is smooth in  $c$  by Hypothesis 6.1. Substituting (6.40)  
 1169 into (6.39) and simplifying, we obtain the equation for  $\tilde{w}$

$$1170 \quad (6.42) \quad \mathcal{A}_0(U_n)\tilde{w} + h(x) = 0,$$

1171 where  $h(x)$  is a small remainder term with uniform bound  $\|h(x)\|_\infty = \mathcal{O}(e^{-\alpha X_{\min}})$ .  
 1172 Using Lemma 6.15, for  $j = 1, \dots, n$  we can write the operator  $\mathcal{A}_0(U_n)$  as,

$$1173 \quad (6.43) \quad \mathcal{A}_0(U_n) = \mathcal{A}_0(U^j) + \sum_{k \neq j} (e^{U^k(x)} - 1) + \tilde{h}(x),$$

1174 where  $\tilde{h}(x)$  is another small remainder term with uniform bound  $\|\tilde{h}\|_\infty = \mathcal{O}(e^{-\alpha X_{\min}})$ .

1175 We now follow the procedure in [31], which we briefly outline below. Let  $W =$   
 1176  $(\tilde{w}, \partial_x \tilde{w}, \partial_x^2 \tilde{w}, \partial_x^3 \tilde{w})$ . As in [31], we rewrite (6.42) as a first-order system for  $W$ , and  
 1177 we take  $W$  to be a piecewise function consisting of the  $2n$  pieces  $W_j^\pm, j = 1, \dots, n$ ,  
 1178 where

$$1179 \quad W_j^-(x) \in C^0([-X_{j-1}, 0])$$

$$1180 \quad W_j^+(x) \in C^0([0, X_j])$$

1182 with  $X_0 = X_n = \infty$ . Following this procedure, and using the expansions (6.43) for  
 1183  $\mathcal{A}_0(U_n)$  on the  $j$ -th piece, we obtain the system of equations

$$1184 \quad (6.44) \quad \begin{aligned} (W_j^\pm)'(x) &= A(U(x))W_j^\pm(x) + G_j(x)W_j^\pm(x) + BH(x) \\ W_j^+(X_i) - W_{j+1}^-(-X_j) &= 0 \\ W_j^-(0) - W_j^+(0) &= 0 \end{aligned}$$

1185 where

$$1186 \quad A(U(x)) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -e^{U(x)} & 0 & -c^2 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

1187 and

$$1188 \quad G_j(x) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \sum_{k \neq j} (1 - e^{U(x - \rho_{kj})}) & 0 & 0 & 0 \end{pmatrix},$$

1189  $\rho_{kj}$  is the signed distance from peak of  $U^k$  to peak of  $U^j$  in  $U_n$ , and  $\|H\| = \mathcal{O}(e^{-\alpha X_{\min}})$ .  
 1190 For  $k \neq j$ ,  $|\rho_{kj}| \geq 2X_{\min}$ . This implies  $e^{U(x - \rho_{kj})} = \mathcal{O}(e^{-\alpha X_{\min}})$  on the  $j$ -th piece, thus  
 1191 we can use a Taylor expansion to show  $\|G_j\| = \mathcal{O}(e^{-\alpha X_{\min}})$ . Following the procedure  
 1192 in [31], we obtain a unique piecewise solution  $W_j^\pm$  to the first two equations of (6.44).  
 1193 The third equation is generally not satisfied, so what we have constructed is a unique  
 1194 solution  $\tilde{y}$  of the form (6.40) to (6.39) which is continuous except for  $n - 1$  jumps. By  
 1195 uniqueness, we must have  $\tilde{y} = y$ , thus  $y$  is actually of the form (6.40) with  $\tilde{w}$  smooth.  
 1196 Finally, Lin's method gives us the uniform bound  $\|\tilde{w}\|_\infty = \mathcal{O}(e^{-2\alpha X_{\min}})$ , from which  
 1197 (6.38) follows.  $\square$

1198 We prove one more lemma before we evaluate the matrix  $\mathbf{K}_2$  from (6.27).

LEMMA 6.17. For the coefficients  $d_{jk}$  in (6.10) from Theorem 6.4,

$$(6.45) \quad \sum_{m=1}^n d_{jm} d_{km} = \delta_{jk} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

*Proof.* Using the expansion (6.10) from Theorem 6.4,

$$\begin{aligned} \langle s_j, s_k \rangle &= \sum_{m=1}^n d_{jm} d_{km} \langle \partial_x U^m, \partial_x U^m \rangle + \sum_{m \neq \ell} d_{jm} d_{k\ell} \langle \partial_x U^m, \partial_x U^\ell \rangle \\ &\quad + \langle s_j, w_k \rangle + \sum_{\ell=1}^n d_{k\ell} \langle w_j, \partial_x U^\ell \rangle. \end{aligned}$$

As in Lemma 6.13, the second term on the RHS is  $\mathcal{O}(e^{-(3\alpha/2)X_{\min}})$ , and the last two terms on the RHS are  $\mathcal{O}(e^{-2\alpha X_{\min}})$ . By translation invariance,  $\langle \partial_x U^m, \partial_x U^m \rangle = \langle \partial_x U, \partial_x U \rangle = \|\partial_x U\|^2$  for all  $m$ . This reduces to

$$\langle s_j, s_k \rangle = \|\partial_x U\|^2 \sum_{m=1}^n d_{jm} d_{km} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

Dividing by  $\|\partial_x U\|^2$  and using the orthogonality relation (6.26) gives us (6.45).  $\square$

Finally, we can evaluate the matrix  $\mathbf{K}_2$  from (6.27).

LEMMA 6.18. For the matrix  $\mathbf{K}_2$  in (6.27),

$$(6.46) \quad (\mathbf{K}_2)_{jk} = -2c \langle \partial_x^2 U, \partial_c U \rangle \delta_{jk} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

*Proof.* By Lemma 6.14,  $(P_{S^\perp} \mathcal{A}_0(U_n) P_{S^\perp})^{-1}$  is a bounded linear operator. Using the bound (6.11) from Theorem 6.4,

$$P_{S^\perp} (P_{S^\perp} \mathcal{A}_0(U_n)|_{S^\perp})^{-1} P_{S^\perp} \partial_x w_k = \mathcal{O}(e^{-2\alpha X_{\min}}).$$

Using this and (6.38) from Lemma 6.16, (6.37) becomes,

$$\begin{aligned} (\mathbf{K}_2)_{jk} &= 4c^2 \left\langle \sum_{m=1}^n d_{jm} \partial_x^2 U^m + \partial_x w_j, -\frac{1}{2c} \sum_{\ell=1}^n d_{k\ell} P_{S^\perp} \partial_c U^\ell + \mathcal{O}(e^{-2\alpha X_{\min}}) \right\rangle \\ &= -2c \left( \sum_{m=1}^n d_{jm} d_{km} \langle \partial_x^2 U^m, P_{S^\perp} \partial_c U^m \rangle + \sum_{m \neq \ell} d_{jm} d_{k\ell} \langle \partial_x^2 U^m, P_{S^\perp} \partial_c U^\ell \rangle \right. \\ &\quad \left. + \sum_{\ell=1}^n \langle \partial_x w_j, d_{k\ell} \partial_c U^\ell \rangle \right) + \mathcal{O}(e^{-2\alpha X_{\min}}). \end{aligned}$$

For  $m \neq \ell$ ,  $\partial_x^2 U^m$  and  $\partial_c U^\ell$  are exponentially separated, thus the second term on the RHS is  $\mathcal{O}(e^{-(3\alpha/2)X_{\min}})$  by Lemma 6.11. Using Hölder's inequality and the remainder bound (6.11), the third term on the RHS is  $\mathcal{O}(e^{-2\alpha X_{\min}})$ . Thus we are left with

$$(6.47) \quad (\mathbf{K}_2)_{jk} = -2c \sum_{m=1}^n d_{jm} d_{km} \langle \partial_x^2 U^m, P_{S^\perp} \partial_c U^m \rangle + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

1220 To evaluate the inner product, we first evaluate  $P_S \partial_c U^m$ . Recalling the normalization  
 1221 (6.26) and using the expansion (6.10), since the  $s_j$  are orthogonal,

$$\begin{aligned}
 P_S \partial_c U^m &= \frac{1}{\|\partial_x U\|} \sum_{j=1}^n \langle s_j, \partial_c U^m \rangle \\
 &= \frac{1}{\|\partial_x U\|} \sum_{j=1}^n \sum_{k=1}^n \langle d_{jk} \partial_x U^k + w_k, \partial_c U^m \rangle \\
 1222 &= \frac{1}{\|\partial_x U\|} \left( \sum_{j=1}^n d_{jm} \langle \partial_x U^m, \partial_c U^m \rangle + \sum_{j=1}^n \sum_{k \neq m} d_{jk} \langle \partial_x U^k, \partial_c U^m \rangle \right) + \mathcal{O}(e^{-2X_{\min}}) \\
 &= \frac{1}{\|\partial_x U\|} \sum_{j=1}^n d_{jm} \langle \partial_x U, \partial_c U \rangle + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}) \\
 &= \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).
 \end{aligned}$$

1223 The third line follows from Lemma 6.11 since  $\partial_x U^k$  and  $\partial_c U^m$  are exponentially  
 1224 separated for  $k \neq m$ , and in the fourth line we use  $\langle \partial_x U, \partial_c U \rangle = 0$ , since  $\partial_x U$  is an  
 1225 odd function and  $\partial_c U$  is an even function. From this, we have

$$1226 \quad P_{S^\perp} \partial_c U^m = (\mathcal{I} - P_S) \partial_c U^m = \partial_c U^m + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}).$$

1227 Substituting this into equation (6.47) and using Lemma 6.17 and translation invari-  
 1228 ance, this becomes

$$\begin{aligned}
 1229 \quad (\mathbf{K}_2)_{jk} &= -2c \sum_{m=1}^n d_{jm} d_{km} \langle \partial_x^2 U^m, \partial_c U^m \rangle = -2c \langle \partial_x^2 U, \partial_c U \rangle \sum_{m=1}^n d_{jm} d_{km} \\
 &= -2c \langle \partial_x^2 U, \partial_c U \rangle \delta_{jk} + \mathcal{O}(e^{-(3\alpha/2)X_{\min}}),
 \end{aligned}$$

1230 which is (6.46). □

1231 Using (6.35) from Lemma 6.13 and (6.46) from Lemma 6.18, the Krein matrix  
 1232 (6.27) becomes,

$$\begin{aligned}
 1233 \quad -\frac{\mathbf{K}_S(z)}{z} &= \|\partial_x U\|^2 \text{diag}(\nu_1, \dots, \nu_n) - (\|\partial_x U\|^2 - 2c \langle \partial_x^2 U, \partial_c U \rangle) \mathbf{I}_n \bar{z}^2 \\
 &\quad + \mathcal{O}(e^{-(3\alpha/2)X_{\min}} |z| + |z|^3).
 \end{aligned}$$

1234 Integrating by parts,

$$\begin{aligned}
 1235 \quad -\frac{\mathbf{K}_S(z)}{z} &= \|\partial_x U\|^2 \text{diag}(\nu_1, \dots, \nu_n) - (\langle \partial_x U, \partial_x U \rangle + c \langle \partial_c \partial_x U, \partial_x U \rangle) \mathbf{I}_n \bar{z}^2 \\
 &\quad + \mathcal{O}(e^{-(3\alpha/2)X_{\min}} |z| + |z|^3) \\
 &= \|\partial_x U\|^2 \text{diag}(\nu_1, \dots, \nu_n) - \partial_c (c \|\partial_x U\|^2) \mathbf{I}_n \bar{z}^2 + \mathcal{O}(e^{-(3\alpha/2)X_{\min}} |z| + |z|^3) \\
 &= \|\partial_x U\|^2 \text{diag}(\nu_1, \dots, \nu_n) + d''(c) \mathbf{I}_n \bar{z}^2 + \mathcal{O}(e^{-(3\alpha/2)X_{\min}} |z| + |z|^3),
 \end{aligned}$$

1236 which is (6.20) in Theorem 6.8.

1237 REFERENCES

- [1] J. ALEXANDER, R. GARDNER, AND C. JONES, *A topological invariant arising in the stability of travelling waves*, J. Reine Angew. Math., 410 (1990), pp. 167–212.
- [2] J. BRONSKI, M. JOHNSON, AND T. KAPITULA, *An instability index theory for quadratic pencils and applications*, Comm. Math. Phys., 327 (2014), pp. 521–550.
- [3] C. BUZZI AND J. LAMB, *Reversible Hamiltonian Liapunov center theorem*, Disc. Cont. Dyn. Sys. Ser. B, 5 (2005), pp. 51–66.
- [4] J. CHAMARD, J. OTTA, AND D. LLOYD, *Computation of minimum energy paths for quasi-linear problems*, J. Scientific Computing, 49 (2011), pp. 180–194.
- [5] A. CHAMPNEYS AND M. GROVES, *A global investigation of solitary-wave solutions to a two-parameter model for water waves*, J. Fluid Mech., 342 (1997), pp. 199–229.
- [6] Y. CHEN AND P. MCKENNA, *Traveling waves in a nonlinearly suspended beam: Theoretical results and numerical observations*, J. of Diff. Eq., 136 (1997), pp. 325–355.
- [7] B. DECONINCK AND T. KAPITULA, *On the spectral and orbital stability of spatially periodic stationary solutions of generalized Korteweg-de Vries equations*, in Hamiltonian Partial Diff. Eq. Appl., P. G. et al., ed., vol. 75, Springer, 2015, pp. 285–322.
- [8] B. DECONINCK AND O. TRICHTCHENKO, *High-frequency instabilities of small-amplitude solutions of Hamiltonian PDEs*, Disc. Cont. Dyn. Systems, 37 (2015), pp. 1323–1358.
- [9] M. GRILLAKIS, J. SHATAH, AND W. STRAUSS, *Stability theory of solitary waves in the presence of symmetry, I*, Journal of Functional Analysis, 74 (1987), pp. 160–197.
- [10] S. HAMMARLING, C. MUNRO, AND F. TISSEUR, *An algorithm for the complete solution of quadratic eigenvalue problems*, ACM Trans. Math. Software, 39 (2013), pp. 1–19.
- [11] M. HÄRÄGÜŞ AND G. IOOSS, *Local Bifurcations, Center Manifolds, and Normal Forms in Infinite-Dimensional Dynamical Systems*, Springer, 2011.
- [12] M. HÄRÄGÜŞ AND T. KAPITULA, *On the spectra of periodic waves for infinite-dimensional Hamiltonian systems*, Physica D, 237 (2008), pp. 2649–2671.
- [13] I. IPSEN AND R. REHMAN, *Perturbation bounds for determinants and characteristic polynomials*, SIAM J. Matrix Anal. and Appl., 30 (2008), pp. 762–776.
- [14] T. KAPITULA, *The Krein signature, Krein eigenvalues, and the Krein Oscillation Theorem*, Indiana U. Math. J., 59 (2010), pp. 1245–1276.
- [15] T. KAPITULA, E. HIBMA, H.-P. KIM, AND J. TIMKOVICH, *Instability indices for matrix polynomials*, Linear Algebra Appl., 439 (2013), pp. 3412–3434.
- [16] T. KAPITULA, P. KEVREKIDIS, AND B. SANDSTEDTE, *Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems*, Physica D, 195 (2004), pp. 263–282.
- [17] T. KAPITULA, P. KEVREKIDIS, AND B. SANDSTEDTE, *Addendum: Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems*, Physica D, 201 (2005), pp. 199–201.
- [18] T. KAPITULA, P. KEVREKIDIS, AND D. YAN, *The Krein matrix: General theory and concrete applications in atomic Bose-Einstein condensates*, SIAM J. Appl. Math., 73 (2013), pp. 1368–1395.
- [19] T. KAPITULA AND K. PROMISLOW, *Stability indices for constrained self-adjoint operators*, Proc. Amer. Math. Soc., 140 (2012), pp. 865–880.
- [20] T. KAPITULA AND K. PROMISLOW, *Spectral and Dynamical Stability of Nonlinear Waves*, Springer-Verlag, 2013.
- [21] T. KAPITULA AND A. STEFANOV, *A Hamiltonian-Krein (instability) index theory for KdV-like eigenvalue problems*, Stud. Appl. Math., 132 (2014), pp. 183–211.
- [22] T. KATO, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, 1980.
- [23] R. KOLLÁR, B. DECONINCK, AND O. TRICHTCHENKO, *Direct characterization of spectral stability of small amplitude periodic waves in scalar Hamiltonian problems via dispersion relation*, SIAM J. Math. Anal. (to appear).
- [24] R. KOLLÁR AND P. MILLER, *Graphical Krein signature theory and Evans-Krein functions*, SIAM Review, 56 (2014), pp. 73–123.
- [25] Y. LI AND K. PROMISLOW, *Structural stability of non-ground state traveling waves of coupled nonlinear Schrödinger equations*, Physica D, 124 (1998), pp. 137–165.
- [26] P. MCKENNA AND W. WALTER, *Travelling waves in a suspension bridge*, SIAM J. Appl. Math., 50 (1990), pp. 703–715.
- [27] D. PELINOVSKY, *Inertia law for spectral stability of solitary waves in coupled nonlinear Schrödinger equations*, Proc. Royal Soc. London A, 461 (2005), pp. 783–812.
- [28] D. PELINOVSKY, *Spectral stability of nonlinear waves in KdV-type evolution equations*, in Nonlinear Physical Systems: Spectral Analysis, Stability, and Bifurcations, O. Kirillov and D. Pelinovsky, eds., Wiley-ISTE, 2014, pp. 377–400.
- [29] B. SANDSTEDTE, *Verzweigungstheorie homokliner Verdopplungen*, PhD thesis, University of Stuttgart, 1993.



- [30] B. SANDSTEDE, *Instability of localized buckling modes in a one-dimensional strut model*, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, 355 (1997), pp. 2083–2097.
- [31] B. SANDSTEDE, *Stability of multiple-pulse solutions*, Trans. Amer. Math. Soc., 350 (1998), pp. 429–472.
- [32] B. SANDSTEDE, *Homoclinic flip bifurcations in conservative reversible systems*. preprint, 2013.
- [33] B. SANDSTEDE AND A. SCHEEL, *Relative Morse indices, Fredholm indices, and group velocities*, Disc. Cont. Dyn. Syst., 20 (2008), pp. 139–158.
- [34] D. SMETS AND J. VAN DEN BERG, *Homoclinic solutions for Swift-Hohenberg and suspension bridge type equations*, J. Diff. Eq., 184 (2002), pp. 78–96.
- [35] O. TRICHTCHENKO, B. DECONINCK, AND R. KOLLÁR, *Stability of periodic traveling waves solutions to the Kawahara equation*, SIAM J. Appl. Dyn., 17 (2018), pp. 2761–2783.
- [36] J. VAN DEN BERG, M. BREDEN, J.-P. LESSARD, AND M. MURRAY, *Continuation of homoclinic orbits in the suspension bridge equation: A computer-assisted proof*, J. Diff. Eq., 264 (2018), pp. 3086–3130.
- [37] V. VOUGALTER AND D. PELINOVSKY, *Eigenvalues of zero energy in the linearized NLS problem*, J. Math. Phys., 47 (2006), p. 062701.
- [38] A. WEINSTEIN, *Normal modes for nonlinear Hamiltonian systems*, Inventiones math., 20 (1973), pp. 47–57.