


TEXTBOOKS IN MATHEMATICS

# GRAPHS & DIGRAPHS

SIXTH EDITION



GARY CHARTRAND  
LINDA LESNIAK  
PING ZHANG



CRC Press  
Taylor & Francis Group

A CHAPMAN & HALL BOOK

TEXTBOOKS in MATHEMATICS

# GRAPHS & DIGRAPHS

**SIXTH EDITION**

GARY CHARTRAND

Western Michigan University  
Kalamazoo, USA

LINDA LESNIAK

Western Michigan University  
Kalamazoo, USA

PING ZHANG

Western Michigan University  
Kalamazoo, USA



**CRC Press**

Taylor & Francis Group

Boca Raton London New York

---

CRC Press is an imprint of the  
Taylor & Francis Group an **informa** business

A CHAPMAN & HALL BOOK

# TEXTBOOKS in MATHEMATICS

---

Series Editors: Al Boggess and Ken Rosen

---

## PUBLISHED TITLES

### ABSTRACT ALGEBRA: AN INQUIRY-BASED APPROACH

Jonathan K. Hodge, Steven Schlicker, and Ted Sundstrom

### ADVANCED LINEAR ALGEBRA

Nicholas Loehr

### ADVANCED LINEAR ALGEBRA

Hugo Woerdeman

### ADVANCED LINEAR ALGEBRA, SECOND EDITION

Bruce N. Cooperstein

### APPLIED ABSTRACT ALGEBRA WITH MAPLE™ AND MATLAB®, THIRD EDITION

Richard Klima, Neil Sigmon, and Ernest Stitzinger

### APPLIED DIFFERENTIAL EQUATIONS: THE PRIMARY COURSE

Vladimir Dobrushkin

### COMPUTATIONAL MATHEMATICS: MODELS, METHODS, AND ANALYSIS WITH MATLAB® AND MPI, SECOND EDITION

Robert E. White

### DIFFERENTIAL EQUATIONS: THEORY, TECHNIQUE, AND PRACTICE, SECOND EDITION

Steven G. Krantz

### DIFFERENTIAL EQUATIONS: THEORY, TECHNIQUE, AND PRACTICE WITH BOUNDARY VALUE PROBLEMS

Steven G. Krantz

### DIFFERENTIAL EQUATIONS WITH MATLAB®: EXPLORATION, APPLICATIONS, AND THEORY

Mark A. McKibben and Micah D. Webster

### ELEMENTARY NUMBER THEORY

James S. Kraft and Lawrence C. Washington

### EXPLORING LINEAR ALGEBRA: LABS AND PROJECTS WITH MATHEMATICA®

Crista Arangala

### GRAPHS & DIGRAPHS, SIXTH EDITION

Gary Chartrand, Linda Lesniak, and Ping Zhang

## PUBLISHED TITLES CONTINUED

### INTRODUCTION TO ABSTRACT ALGEBRA, SECOND EDITION

Jonathan D. H. Smith

### INTRODUCTION TO MATHEMATICAL PROOFS: A TRANSITION TO ADVANCED MATHEMATICS, SECOND EDITION

Charles E. Roberts, Jr.

### INTRODUCTION TO NUMBER THEORY, SECOND EDITION

Marty Erickson, Anthony Vazzana, and David Garth

### LINEAR ALGEBRA, GEOMETRY AND TRANSFORMATION

Bruce Solomon

### MATHEMATICAL MODELLING WITH CASE STUDIES: USING MAPLE™ AND MATLAB®, THIRD EDITION

B. Barnes and G. R. Fulford

### MATHEMATICS IN GAMES, SPORTS, AND GAMBLING—THE GAMES PEOPLE PLAY, SECOND EDITION

Ronald J. Gould

### THE MATHEMATICS OF GAMES: AN INTRODUCTION TO PROBABILITY

David G. Taylor

### MEASURE THEORY AND FINE PROPERTIES OF FUNCTIONS, REVISED EDITION

Lawrence C. Evans and Ronald F. Gariepy

### NUMERICAL ANALYSIS FOR ENGINEERS: METHODS AND APPLICATIONS, SECOND EDITION

Bilal Ayyub and Richard H. McCuen

### ORDINARY DIFFERENTIAL EQUATIONS: AN INTRODUCTION TO THE FUNDAMENTALS

Kenneth B. Howell

### RISK ANALYSIS IN ENGINEERING AND ECONOMICS, SECOND EDITION

Bilal M. Ayyub

### TRANSFORMATIONAL PLANE GEOMETRY

Ronald N. Umble and Zhigang Han

CRC Press  
Taylor & Francis Group  
6000 Broken Sound Parkway NW, Suite 300  
Boca Raton, FL 33487-2742

© 2016 by Taylor & Francis Group, LLC  
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works  
Version Date: 20151109

International Standard Book Number-13: 978-1-4987-3580-3 (eBook - PDF)

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access [www.copyright.com](http://www.copyright.com) (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

**Trademark Notice:** Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

**Visit the Taylor & Francis Web site at**  
**<http://www.taylorandfrancis.com>**

**and the CRC Press Web site at**  
**<http://www.crcpress.com>**

**To**

the memory of my mother and father.    G. C.

my mother and the memory of my father Stanley.    L. L.

my mother and the memory of my father.    P. Z.



# Table of Contents

Preface to the Sixth Edition	xi
<b>1. Introduction</b>	<b>1</b>
1.1 Graphs	1
1.2 The Degree of a Vertex	5
1.3 Isomorphic Graphs	7
1.4 Regular Graphs	12
1.5 Bipartite Graphs	13
1.6 Operations on Graphs	16
1.7 Degree Sequences	18
1.8 Multigraphs	25
• Exercises for Chapter 1	28
<b>2. Connected Graphs and Distance</b>	<b>37</b>
2.1 Connected Graphs	37
2.2 Distance in Graphs	44
• Exercises for Chapter 2	51
<b>3. Trees</b>	<b>57</b>
3.1 Nonseparable Graphs	57
3.2 Introduction to Trees	62
3.3 Spanning Trees	69
3.4 The Minimum Spanning Tree Problem	81
• Exercises for Chapter 3	86
<b>4. Connectivity</b>	<b>95</b>
4.1 Connectivity and Edge-Connectivity	95
4.2 Theorems of Menger and Whitney	102
• Exercises for Chapter 4	110
<b>5. Eulerian Graphs</b>	<b>115</b>
5.1 The Königsberg Bridge Problem	115
5.2 Eulerian Circuits and Trails	117
• Exercises for Chapter 5	123



<b>6. Hamiltonian Graphs</b>	<b>125</b>
6.1 Hamilton's Icosian Game	125
6.2 Sufficient Conditions for Hamiltonicity	128
6.3 Toughness of Graphs	134
6.4 Highly Hamiltonian Graphs	140
6.5 Powers of Graphs and Line Graphs	145
• Exercises for Chapter 6	154
<b>7. Digraphs</b>	<b>161</b>
7.1 Introduction to Digraphs	161
7.2 Strong Digraphs	166
7.3 Eulerian and Hamiltonian Digraphs	167
7.4 Tournaments	170
7.5 Kings in Tournaments	179
7.6 Hamiltonian Tournaments	180
• Exercises for Chapter 7	184
<b>8. Flows in Networks</b>	<b>191</b>
8.1 Networks	191
8.2 The Max-Flow Min-Cut Theorem	199
8.3 Menger Theorems for Digraphs	207
• Exercises for Chapter 8	212
<b>9. Automorphisms and Reconstruction</b>	<b>217</b>
9.1 The Automorphism Group of a Graph	217
9.2 Cayley Color Graphs	223
9.3 The Reconstruction Problem	228
• Exercises for Chapter 9	235
<b>10. Planar Graphs</b>	<b>239</b>
10.1 The Euler Identity	239
10.2 Maximal Planar Graphs	248
10.3 Characterizations of Planar Graphs	252
10.4 Hamiltonian Planar Graphs	264
• Exercises for Chapter 10	268

<b>11. Nonplanar Graphs</b>	<b>275</b>
11.1 The Crossing Number of a Graph	275
11.2 The Genus of a Graph	286
11.3 The Graph Minor Theorem	300
• Exercises for Chapter 11	302
<b>12. Matchings, Independence and Domination</b>	<b>305</b>
12.1 Matchings	305
12.2 1-Factors	310
12.3 Independence and Covers	317
12.4 Domination	321
• Exercises for Chapter 12	329
<b>13. Factorization and Decomposition</b>	<b>335</b>
13.1 Factorization	335
13.2 Decomposition	343
13.3 Cycle Decomposition	345
13.4 Graceful Graphs	351
• Exercises for Chapter 13	358
<b>14. Vertex Colorings</b>	<b>363</b>
14.1 The Chromatic Number of a Graph	363
14.2 Color-Critical Graphs	371
14.3 Bounds for the Chromatic Number	374
• Exercises for Chapter 14	385
<b>15. Perfect Graphs and List Colorings</b>	<b>393</b>
15.1 Perfect Graphs	393
15.2 The Perfect and Strong Perfect Graph Theorems	402
15.3 List Colorings	405
• Exercises for Chapter 15	410
<b>16. Map Colorings</b>	<b>415</b>
16.1 The Four Color Problem	415
16.2 Colorings of Planar Graphs	426
16.3 List Colorings of Planar Graphs	428
16.4 The Conjectures of Hajós and Hadwiger	434

16.5 Chromatic Polynomials	438
16.6 The Heawood Map-Coloring Problem	444
• Exercises for Chapter 16	448
<b>17. Edge Colorings</b>	<b>453</b>
17.1 The Chromatic Index of a Graph	453
17.2 Class One and Class Two Graphs	460
17.3 Tait Colorings	467
• Exercises for Chapter 17	476
<b>18. Nowhere-Zero Flows, List Edge Colorings</b>	<b>481</b>
18.1 Nowhere-Zero Flows	481
18.2 List Edge Colorings	491
18.3 Total Colorings	495
• Exercises for Chapter 18	500
<b>19. Extremal Graph Theory</b>	<b>503</b>
19.1 Turán's Theorem	503
19.2 Extremal Subgraphs	505
19.3 Cages	509
• Exercises for Chapter 19	520
<b>20. Ramsey Theory</b>	<b>523</b>
20.1 Classical Ramsey Numbers	523
20.2 More General Ramsey Numbers	532
• Exercises for Chapter 20	538
<b>21. The Probabilistic Method</b>	<b>543</b>
21.1 The Probabilistic Method	543
21.2 Random Graphs	554
• Exercises for Chapter 21	561
<b>Hints and Solutions to Odd-Numbered Exercises</b>	<b>563</b>
<b>References</b>	<b>589</b>
<b>Supplemental References</b>	<b>606</b>
<b>Index of Names</b>	<b>607</b>
<b>Index of Mathematical Terms</b>	<b>613</b>
<b>Index of Symbols</b>	<b>625</b>

## Preface to the Sixth Edition

Graph theory is an area of mathematics whose origin dates back to 1736 with the solution of the famous Königsberg Bridge Problem by the eminent Swiss mathematician Leonhard Euler. During the next several decades, topics in graph theory arose primarily through recreational mathematics. The development of graph theory received a substantial boost in 1852 when the young British mathematician Francis Guthrie introduced one of the best known problems in all of mathematics: the Four Color Problem. It wasn't until late in the 19th century, however, when graph theory became a theoretical area of mathematics through the research of the Danish mathematician Julius Petersen. Major progress in graph theory, however, didn't occur until World War II ended. Since then, though, the subject has developed into an area with a fascinating history, numerous interesting problems and applications in many diverse fields. It is the beauty of the subject, however, that has attracted so many to this field.

The goal of this sixth edition is, as with the previous editions, to describe much of the story that is graph theory – through its concepts, its theorems, its applications and its history. The audience for the sixth edition is beginning graduate students and advanced undergraduate students. The primary prerequisite required of students using this book is a knowledge of mathematical proofs. For some topics, an elementary knowledge of linear algebra and group theory is useful. For Chapter 21, an elementary knowledge of probability is needed. Proofs of some of the results that appear in this book have not been supplied because the techniques are beyond the scope of the book or are inordinately lengthy. Nevertheless, these results have been included due to their interest and since they provide a more complete description of what is known on a particular topic.

A one-semester course in graph theory using this textbook can be designed by selecting topics of greatest interest to the instructor and students. There is more than ample material available for a two-semester sequence in graph theory. Our goal has been to prepare a book that is interesting, carefully written, student-friendly and consisting of clear proofs. The sixth edition has been divided into shorter chapters as well as more sections and subsections to make reading and locating material easier for instructors and students. The following major additions have been made to the sixth edition:

- more than 160 new exercises
- several conjectures and open problems
- many new theorems and examples
- new material on graph decompositions
- a proof of the Perfect Graph Theorem

- material on Hamiltonian extension
- a new chapter on the probabilistic method in graph theory and random graphs.

At the end of the book is an index of mathematicians, an index of mathematical terms and an index of symbols. The references list research papers referred to in the book (indicating the page number(s) where the reference occurs) and some useful supplemental references. There is also a section giving hints and solutions to all odd-numbered exercises.

Over the years, there have been some changes in notation that a number of mathematicians now use. When certain notation appears to have been adopted by sufficiently many mathematicians working in graph theory so that this has become the norm, we have adhered to these changes. As with the fifth edition, the following notation is used in the sixth edition:

- a path is now expressed as  $P = (v_1, v_2, \dots, v_k)$  and a cycle as  $C = (v_1, v_2, \dots, v_k, v_1)$ ;
- the Cartesian product of two graphs  $G$  and  $H$  is expressed as  $G \square H$ , rather than the previous  $G \times H$ ;
- the union of  $G$  and  $H$  is expressed by  $G + H$ , rather than  $G \cup H$ ;
- the join of two graphs  $G$  and  $H$  is expressed as  $G \vee H$ , rather than  $G + H$ .

We are most grateful to Bob Ross, senior editor of CRC Press, who has been a constant source of support and assistance throughout the entire writing process.

Gary Chartrand, Linda Lesniak and Ping Zhang

# Chapter 1

## Introduction

*The theory of graphs is one of the few fields of mathematics with a definite birth date.*

It is the subject of graph theory of course that we are about to describe. The statement above was made in 1963 by the mathematician Oystein Ore who will be encountered in Chapter 6. While graph theory was probably Ore's major mathematical area of interest during the latter part of his career, he is also known for his work and interest in number theory (the study of integers) and the history of mathematics.

Although awareness of integers can be traced back for many centuries, geometry has an even longer history. Early geometry concerned distance, lengths, angles, areas and volumes, which were used for surveying, construction and astronomy. While geometry dealt with magnitudes, the German mathematician Gottfried Leibniz introduced another branch of geometry called the geometry of position. This branch of geometry did not deal with measurements and calculations, but rather with the determination of position and its properties. The famous mathematician Leonhard Euler said that it hadn't been determined what kinds of problems could be studied with the aid of the geometry of position but in 1736 he believed that he had found one, which led to the origin of graph theory. It is this event to which Oystein Ore was referring in his quote above. We will visit Euler again, in Chapter 5 as well as in Chapters 10 and 11.

### 1.1 Graphs

Graphs arise in many different settings. Let's look at three of these.

**Example 1.1** Eight students  $s_1, s_2, \dots, s_8$  have been invited to a dinner. Each student knows only some of the other students. The students that each student knows are listed below.

$s_1 : s_4, s_5, s_8$	$s_2 : s_3, s_4, s_6, s_8$
$s_3 : s_2, s_5, s_6, s_7, s_8$	$s_4 : s_1, s_2, s_5, s_6, s_7$
$s_5 : s_1, s_3, s_4, s_8$	$s_6 : s_2, s_3, s_4, s_7$
$s_7 : s_3, s_4, s_6$	$s_8 : s_1, s_2, s_3, s_5$

In order to determine if these eight students can be seated at a round table where each student sits next to two students he or she knows, it is useful to represent this situation by the diagram shown in Figure 1.1. Each point or small circle in the diagram represents a student and two points are joined by a line segment if the two students know each other. This diagram is referred to as a *graph*.




Figure 1.1: The diagram in Example 1.1

A related question is whether the students could be seated at a round table so that each student sits next to two students he or she does not know. ♦

**Example 1.2** There are six special locations in a neighborhood park. Twelve trails are to be built between certain pairs of these locations, namely all pairs of locations except  $\{a_1, a_2\}$ ,  $\{b_1, b_2\}$ ,  $\{c_1, c_2\}$  (see Figure 1.2(a)). A trail can be straight or curved. Can this be done without any trails crossing? This situation can be represented by the diagram with six points (each point representing a location), where two points are joined by a line segment or a curve if the two points represent locations to be joined by a trail (see Figure 1.2(b)). Once again, this diagram is a graph. ♦

**Example 1.3** A chemical company is to ship eight chemicals (denoted by  $c_1, c_2, \dots, c_8$ ) to a chemistry department in a university. Because some pairs of chemicals should not be shipped in the same container, more than one container needs to be used for this shipment. Each chemical is listed below together with the chemicals that should not be placed in the same container as this chemical.

$c_1 : c_2$	$c_2 : c_1, c_8$	$c_3 : c_5, c_6, c_7$	$c_4 : c_5, c_7$
$c_5 : c_3, c_4, c_8$	$c_6 : c_3, c_7$	$c_7 : c_3, c_4, c_6$	$c_8 : c_2, c_5$

It would be useful to know the minimum number of containers needed to ship these eight chemicals. This situation can be represented by the diagram




Figure 1.2: Constructing a graph in Example 1.2

in Figure 1.3, whose eight points represent the eight chemicals and where two points are joined by a line segment or curve if these chemicals cannot be shipped in the same container. Here too, this diagram is a graph. ♦




Figure 1.3: The graph in Example 1.3

We now give a formal definition of the term *graph*. A **graph**  $G$  is a finite nonempty set  $V$  of objects called **vertices** (the singular is **vertex**) together with a possibly empty set  $E$  of 2-element subsets of  $V$  called **edges**. Vertices are sometimes referred to as **points** or **nodes**, while edges are sometimes called **lines** or **links**. In fact, historically, graphs were referred to as *linkages* by some. Calling these structures graphs was evidently the idea of James Joseph Sylvester (1814–1897), a well-known British mathematician who became the first mathematics professor at Johns Hopkins University in Baltimore and who founded and became editor-in-chief of the first mathematics journal in the United States (the *American Journal of Mathematics*).

To indicate that a graph  $G$  has **vertex set**  $V$  and **edge set**  $E$ , we write  $G = (V, E)$ . To emphasize that  $V$  and  $E$  are the vertex set and edge set of a graph  $G$ , we often write  $V$  as  $V(G)$  and  $E$  as  $E(G)$ . Each edge  $\{u, v\}$  of  $G$  is usually denoted by  $uv$  or  $vu$ . If  $e = uv$  is an edge of  $G$ , then  $e$  is said to **join**  $u$  and  $v$ .



As the examples described above indicate, a graph  $G$  can be represented by a diagram, where each vertex of  $G$  is represented by a point or small circle and an edge joining two vertices is represented by a line segment or curve joining the corresponding points in the diagram. It is customary to refer to such a diagram as the graph  $G$  itself. In addition, the points in the diagram are referred to as the vertices of  $G$  and the line segments are referred to as the edges of  $G$ . For example, the graph  $G$  with vertex set  $V(G) = \{u, v, w, x, y\}$  and edge set  $E(G) = \{uv, uy, vx, vy, wy, xy\}$  is shown in Figure 1.4. Even though the edges  $vx$  and  $wy$  cross in Figure 1.4, their point of intersection is not a vertex of  $G$ .




Figure 1.4: A graph

If  $uv$  is an edge of  $G$ , then  $u$  and  $v$  are **adjacent vertices**. Two adjacent vertices are referred to as **neighbors** of each other. The set of neighbors of a vertex  $v$  is called the **open neighborhood** of  $v$  (or simply the **neighborhood** of  $v$ ) and is denoted by  $N_G(v)$ , or  $N(v)$  if the graph  $G$  is understood. The set  $N[v] = N(v) \cup \{v\}$  is called the **closed neighborhood** of  $v$ . If  $uv$  and  $vw$  are distinct edges in  $G$ , then  $uv$  and  $vw$  are **adjacent edges**. The vertex  $u$  and the edge  $uv$  are said to be **incident** with each other. Similarly,  $v$  and  $uv$  are incident.

For the graph  $G$  of Figure 1.4, the vertices  $u$  and  $v$  are therefore adjacent in  $G$ , while the vertices  $u$  and  $x$  are not adjacent. The edges  $uv$  and  $vx$  are adjacent in  $G$ , while the edges  $vx$  and  $wy$  are not adjacent. The vertex  $v$  is incident with the edge  $uv$  but is not incident with the edge  $wy$ .

The number of vertices in a graph  $G$  is the **order** of  $G$  and the number of edges is the **size** of  $G$ . The order of the graph  $G$  of Figure 1.4 is 5 and its size is 6. We typically use  $n$  and  $m$  for the order and size, respectively, of a graph. A graph of order 1 is called a **trivial graph**. A **nontrivial graph** therefore has two or more vertices. A graph of size 0 is called an **empty graph**. A **nonempty graph** then has one or more edges. In any empty graph, no two vertices are adjacent. At the other extreme is a **complete graph** in which every two distinct vertices are adjacent. The size of a complete graph of order  $n$  is  $\binom{n}{2} = n(n-1)/2$ . Therefore, for every graph  $G$  of order  $n$  and size  $m$ , it follows that  $0 \leq m \leq \binom{n}{2}$ . The complete graph of order  $n$  is denoted by  $K_n$ . The complete graphs  $K_n$  for  $1 \leq n \leq 5$  are shown in Figure 1.5.




Figure 1.5: Some complete graphs

Two other classes of graphs that are often encountered are the paths and cycles. For an integer  $n \geq 1$ , the **path**  $P_n$  is a graph of order  $n$  and size  $n - 1$  whose vertices can be labeled by  $v_1, v_2, \dots, v_n$  and whose edges are  $v_i v_{i+1}$  for  $i = 1, 2, \dots, n - 1$ . For an integer  $n \geq 3$ , the **cycle**  $C_n$  is a graph of order  $n$  and size  $n$  whose vertices can be labeled by  $v_1, v_2, \dots, v_n$  and whose edges are  $v_1 v_n$  and  $v_i v_{i+1}$  for  $i = 1, 2, \dots, n - 1$ . The cycle  $C_n$  is also referred to as an  **$n$ -cycle** and the **3-cycle** is also called a **triangle**. The paths and cycles of order 5 or less are shown in Figure 1.6. Observe that  $P_1 = K_1$ ,  $P_2 = K_2$  and  $C_3 = K_3$ .




Figure 1.6: Paths and cycles of order 5 or less

## 1.2 The Degree of a Vertex

The **degree of a vertex**  $v$  in a graph  $G$  is the number of vertices in  $G$  that are adjacent to  $v$ . Thus, the degree of  $v$  is the number of vertices in its neighborhood  $N(v)$ . Equivalently, the degree of  $v$  is the number of edges incident with  $v$ . The degree of a vertex  $v$  is denoted by  $\deg_G v$  or, more simply, by  $\deg v$  if the graph  $G$  under discussion is clear. Hence,  $\deg v = |N(v)|$ . A vertex of degree 0 is referred to as an **isolated vertex** and a vertex of degree 1 is an **end-vertex** or a **leaf**. An edge incident with an end-vertex is called a **pendant edge**. The largest degree among the vertices of  $G$  is called the **maximum degree** of  $G$  and is denoted by  $\Delta(G)$ . The **minimum degree** of  $G$  is denoted by  $\delta(G)$ . (The symbols  $\Delta$  and  $\delta$  are the upper case and lower case Greek letter delta, respectively.) Thus, if  $v$  is a vertex of a graph  $G$  of order  $n$ , then

$$0 \leq \delta(G) \leq \deg v \leq \Delta(G) \leq n - 1.$$

For the graph  $G$  of Figure 1.4,

$$\deg w = 1, \deg u = \deg x = 2, \deg v = 3 \text{ and } \deg y = 4.$$

Thus,  $\delta(G) = 1$  and  $\Delta(G) = 4$ .

## The First Theorem of Graph Theory

A well-known theorem in graph theory dealing with the sum of the degrees of the vertices of a graph was observed (indirectly) by Leonhard Euler in a 1736 paper [86] of his that is now considered the first paper ever written on graph theory – even though graphs were never mentioned in the paper. This observation is often referred to as the **First Theorem of Graph Theory**. Some have called Theorem 1.4 the **Handshaking Lemma**, although Euler never used this name.

**Theorem 1.4 (The First Theorem of Graph Theory)** *If  $G$  is a graph of size  $m$ , then*

$$\sum_{v \in V(G)} \deg v = 2m.$$

**Proof.** When summing the degrees of the vertices of  $G$ , each edge of  $G$  is counted twice, once for each of its two incident vertices. ■

The sum of the degrees of the vertices of the graph  $G$  of Figure 1.4 is 12, which is twice the size 6 of  $G$ , as expected from Theorem 1.4. The **average degree** of a graph  $G$  of order  $n$  and size  $m$  is

$$\frac{\sum_{v \in V(G)} \deg v}{n} = \frac{2m}{n}.$$

For example, the average degree of the graph  $G$  of Figure 1.4 (having order  $n = 5$  and size  $m = 6$ ) is  $2m/n = 12/5$ . Since the average degree of this graph is strictly between 2 and 3, it follows that  $G$  must have a vertex of degree 3 or more and a vertex of degree 2 or less. This graph actually has vertices of degrees 3 and 4 as well as vertices of degrees 1 and 2.

## Even and Odd Vertices

A vertex in a graph  $G$  is **even** or **odd**, according to whether its degree in  $G$  is even or odd. Thus, the graph  $G$  of Figure 1.4 has three even vertices and two odd vertices. While a graph can have either an even or an odd number of even vertices, this is not the case for odd vertices.

**Corollary 1.5** *Every graph has an even number of odd vertices.*

**Proof.** Suppose that  $G$  is a graph of size  $m$ . By Theorem 1.4,

$$\sum_{v \in V(G)} \deg v = 2m,$$

which, of course, is an even number. Since the sum of the degrees of the even vertices of  $G$  is even, the sum of the degrees of the odd vertices of  $G$  must be even as well, implying that  $G$  has an even number of odd vertices. ■

## 1.3 Isomorphic Graphs

There is only one graph of order 1, two graphs of order 2, four graphs of order 3 and eleven graphs of order 4. All 18 of these graphs are shown in Figure 1.7. This brings up the question of why every two graphs in Figure 1.7 are considered different. In fact, there is the related question of what it means for two graphs to be considered the same. The technical term for this is *isomorphic graphs* (graphs having the same structure).




Figure 1.7: The (non-isomorphic) graphs of order 4 or less

Two graphs  $G$  and  $H$  are **isomorphic** if there exists a bijective function  $\phi : V(G) \rightarrow V(H)$  such that two vertices  $u$  and  $v$  are adjacent in  $G$  if and only if  $\phi(u)$  and  $\phi(v)$  are adjacent in  $H$ . The function  $\phi$  is called an **isomorphism** from  $G$  to  $H$ . If  $\phi : V(G) \rightarrow V(H)$  is an isomorphism, then the inverse function  $\phi^{-1} : V(H) \rightarrow V(G)$  is an isomorphism from  $H$  to  $G$ . If  $G$  and  $H$  are isomorphic, we write  $G \cong H$ . If there is no such function  $\phi$  as described above, then  $G$  and  $H$  are **non-isomorphic graphs** and we write  $G \not\cong H$ .

The graphs  $G$  and  $H$  of Figure 1.8 (both of order 7 and size 8) are isomorphic and the function  $\phi : V(G) \rightarrow V(H)$  defined by

$$\phi(u_1) = v_4, \phi(u_2) = v_5, \phi(u_3) = v_1,$$

$$\phi(u_4) = v_6, \phi(u_5) = v_2, \phi(u_6) = v_3, \phi(u_7) = v_7$$

is an isomorphism, although there are three other isomorphisms. The graphs  $F_1$  and  $F_2$  of Figure 1.8 (both of order 7 and size 10) are not isomorphic however. An explanation of this will be given shortly.

The graphs  $G_1$  and  $G_2$  in Figure 1.9 are isomorphic, while  $G_1$  and  $G_3$  are not isomorphic. For example, the function  $\phi : V(G_1) \rightarrow V(G_2)$  defined by

$$\phi(u_1) = v_1, \phi(u_2) = v_3, \phi(u_3) = v_5,$$

$$\phi(u_4) = v_2, \phi(u_5) = v_4, \phi(u_6) = v_6$$

is an isomorphism. The graph  $G_3$  of Figure 1.9 contains three mutually adjacent vertices  $w_1, w_2, w_6$ . If  $G_1$  and  $G_3$  were isomorphic, then for an isomorphism




Figure 1.8: Isomorphic and non-isomorphic graphs

$\alpha : V(G_3) \rightarrow V(G_1)$ , the vertices  $\alpha(w_1), \alpha(w_2), \alpha(w_6)$  must also be mutually adjacent in  $G_1$ . Since  $G_1$  does not contain three mutually adjacent vertices, there is no isomorphism from  $G_3$  to  $G_1$  and so  $G_1 \not\cong G_3$ . Furthermore,  $G_2 \not\cong G_3$  as well.




Figure 1.9: Isomorphic and non-isomorphic graphs

Suppose that two graphs  $G$  and  $H$  are isomorphic. Then there exists an isomorphism  $\phi : V(G) \rightarrow V(H)$ . Since  $\phi$  is a bijective function,  $|V(G)| = |V(H)|$ . Furthermore, since two vertices  $u$  and  $v$  are adjacent in  $G$  if and only if  $\phi(u)$  and  $\phi(v)$  are adjacent in  $H$ , it follows that  $|E(G)| = |E(H)|$ . These facts are summarized below, together with another necessary condition for two graphs to be isomorphic.

**Theorem 1.6** *If two graphs  $G$  and  $H$  are isomorphic, then they have the same order and the same size, and the degrees of the vertices of  $G$  are the same as the degrees of the vertices of  $H$ .*

**Proof.** We have already observed that isomorphic graphs have the same order and the same size. Let  $v$  be a vertex of  $G$  and suppose that  $\deg v = k$ . Then  $v$  is adjacent to  $k$  vertices, say  $v_1, v_2, \dots, v_k$ . Suppose that  $v$  is not adjacent to  $u_1, u_2, \dots, u_\ell$ . If  $\phi : V(G) \rightarrow V(H)$  is an isomorphism, then  $\phi(v)$  is adjacent to  $\phi(v_1), \phi(v_2), \dots, \phi(v_k)$  while  $\phi(v)$  is not adjacent to  $\phi(u_1), \phi(u_2), \dots, \phi(u_\ell)$ . Hence,  $\phi(v)$  has degree  $k$  in  $H$ . ■

The proof of Theorem 1.6 also shows that every isomorphism from a graph  $G$  to a graph  $H$  maps every vertex of  $G$  to a vertex of the same degree in  $H$ .

It is therefore a consequence of Theorem 1.6 that if  $G$  and  $H$  are two graphs such that (1) the orders of  $G$  and  $H$  are different, or (2) the sizes of  $G$  and  $H$  are different or (3) the degrees of the vertices of  $G$  and those of the vertices of  $H$  are different, then  $G$  and  $H$  are not isomorphic. Since the graph  $F_2$  in Figure 1.8 contains a vertex of degree 5 and no vertex of  $F_1$  has degree 5, it follows by Theorem 1.6 that  $F_1 \not\cong F_2$ .

The conditions described in Theorem 1.6 are strictly necessary for two graphs to be isomorphic – they are not sufficient. Indeed, the graphs  $G_1$  and  $G_3$  of Figure 1.9 have the same order, the same size and the degrees of the vertices of  $G_1$  and  $G_3$  are the same; yet  $G_1$  and  $G_3$  are not isomorphic.

Next, consider the four graphs  $H_1, H_2, H_3$  and  $H_4$  shown in Figure 1.10. The graphs  $H_1$  and  $H_2$  have order 7, size 7 and the degrees of the vertices of these two graphs are the same. Furthermore, each graph contains a single triangle. Nevertheless,  $H_1 \not\cong H_2$ , for suppose that there is an isomorphism  $\phi$  from  $H_1$  to  $H_2$ . Since each graph has only one vertex of degree 3 and one vertex of degree 4,  $\phi(u_1) = v_2$  and  $\phi(y_1) = x_2$ . Since  $v_1$  is adjacent to  $u_1$  and  $y_1$ , it follows that  $\phi(v_1)$  is adjacent to  $v_2$  and  $x_2$ . However,  $H_2$  contains no vertex adjacent to  $v_2$  and  $x_2$ . Thus,  $H_1 \not\cong H_2$ .




Figure 1.10: Non-isomorphic graphs

The graphs  $H_3$  and  $H_4$  of Figure 1.10 have order 9, size 8 and the degrees of the vertices of these two graphs are the same; yet these two graphs as well are *not* isomorphic. Suppose that there is an isomorphism  $\phi$  from  $H_3$  to  $H_4$ . Consider the vertex  $v_3$ . Since  $\deg_{H_3} v_3 = 2$ , it follows that  $\phi(v_3)$  has degree 2 in  $H_4$ . Since  $v_3$  is adjacent to  $u_3$  and  $w_3$ , we must have  $\phi(v_3)$  adjacent to  $\phi(u_3)$  and  $\phi(w_3)$ . Since  $\deg_{H_3} u_3 = \deg_{H_3} w_3 = 2$ , it follows that  $\phi(u_3)$  and  $\phi(w_3)$  have degree 2 in  $H_4$ . But no vertex of degree 2 in  $H_4$  is adjacent to two vertices of degree 2. Thus,  $H_3 \not\cong H_4$ .

### Subgraphs

A graph  $H$  is a **subgraph** of a graph  $G$  if  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ , in which case we write  $H \subseteq G$ . If  $H$  is a subgraph of  $G$ , then  $G$  is a **supergraph** of  $H$ . If  $V(H) = V(G)$ , then  $H$  is a **spanning subgraph** of  $G$ . If  $H$  is a subgraph of a graph  $G$  where  $H \not\cong G$ , then  $H$  is a **proper subgraph** of  $G$ . Therefore, if  $H$  is a proper subgraph of  $G$ , then either  $V(H)$  is a proper subset of  $V(G)$  or  $E(H)$  is a proper subset of  $E(G)$ .

Figure 1.11 shows six graphs, namely  $G$  and the graphs  $G_i$  for  $i = 1, 2, \dots, 5$ . All six of these graphs are proper subgraphs of  $G$ , except  $G$  itself and  $G_1$ . Although  $G$  is a subgraph of itself, it is not a proper subgraph of  $G$ . The graph  $G_1$  contains the edge  $uz$ , which is not an edge of  $G$  and so  $G_1$  is not even a subgraph of  $G$ . The graph  $G_3$  is a spanning subgraph of  $G$  since  $V(G_3) = V(G)$ .




Figure 1.11: Graphs and subgraphs

### Induced Subgraphs

For a nonempty subset  $S$  of  $V(G)$ , the **subgraph  $G[S]$  of  $G$  induced by  $S$**

has  $S$  as its vertex set and two vertices  $u$  and  $v$  are adjacent in  $G[S]$  if and only if  $u$  and  $v$  are adjacent in  $G$ . A subgraph  $H$  of a graph  $G$  is called an **induced subgraph** if there is a nonempty subset  $S$  of  $V(G)$  such that  $H = G[S]$ . Thus  $G[V(G)] = G$ . For a nonempty set  $X$  of edges of a graph  $G$ , the **subgraph  $G[X]$  induced by  $X$**  has  $X$  as its edge set and a vertex  $v$  belongs to  $G[X]$  if  $v$  is incident with at least one edge in  $X$ . A subgraph  $H$  of  $G$  is **edge-induced** if there is a nonempty subset  $X$  of  $E(G)$  such that  $H = G[X]$ . Thus,  $G[E(G)] = G$  if and only if  $G$  has no isolated vertices.

Once again, consider the graphs shown in Figure 1.11. Since  $xy \in E(G)$  but  $xy \notin E(G_4)$ , the subgraph  $G_4$  is not an induced subgraph of  $G$ . On the other hand, the subgraphs  $G_2$  and  $G_5$  are both induced subgraphs of  $G$ . Indeed, for  $S_1 = \{v, x, y, z\}$  and  $S_2 = \{u, v, y, z\}$ ,  $G_2 = G[S_1]$  and  $G_5 = G[S_2]$ . The subgraph  $G_4$  of  $G$  is edge-induced; in fact,  $G_4 = G[X]$ , where  $X = \{uw, wx, wy, xz, yz\}$ .

For a vertex  $v$  and an edge  $e$  in a nonempty graph  $G = (V, E)$ , the subgraph  $G - v$ , obtained by deleting  $v$  from  $G$ , is the induced subgraph  $G[V - \{v\}]$  of  $G$  and the subgraph  $G - e$ , obtained by deleting  $e$  from  $G$ , is the spanning subgraph of  $G$  with edge set  $E - \{e\}$ . More generally, for a proper subset  $U$  of  $V$ , the graph  $G - U$  is the induced subgraph  $G[V - U]$  of  $G$ . For a subset  $X$  of  $E$ , the graph  $G - X$  is the spanning subgraph of  $G$  with edge set  $E - X$ . If  $u$  and  $v$  are distinct nonadjacent vertices of  $G$ , then  $G + uv$  is the graph with  $V(G + uv) = V(G)$  and  $E(G + uv) = E(G) \cup \{uv\}$ . Thus,  $G$  is a spanning subgraph of  $G + uv$ . For the graph  $G$  of Figure 1.12, the set  $U = \{t, x\}$  of vertices and the set  $X = \{tw, ux, vx\}$  of edges, the subgraphs  $G - u$ ,  $G - wx$ ,  $G - U$  and  $G - X$  of  $G$  are also shown in that figure, as is the graph  $G + uv$ .




Figure 1.12: Deleting vertices and edges from and adding edges to a graph



If every two edges  $e_1$  and  $e_2$  of a graph  $G$  have the property that  $G - e_1 \cong G - e_2$ , then we write  $G - e$  for the deletion of any edge from  $G$ . Hence, for  $n \geq 2$ ,  $K_n - e$  denotes the graph obtained by deleting any edge from  $K_n$ . If  $G + uv \cong G + xy$  for any two pairs  $\{u, v\}$  and  $\{x, y\}$  of nonadjacent vertices of  $G$ , then we write  $G + e$  for the addition of any edge to  $G$ . In particular,  $C_4 + e = K_4 - e$ .

## 1.4 Regular Graphs

There are certain classes of graphs that occur so often that they deserve special mention and, in some cases, special notation. We describe some of the most prominent of these now.

A graph  $G$  is **regular** if the vertices of  $G$  have the same degree and is **regular of degree  $r$**  if this degree is  $r$ . Such graphs are also called  **$r$ -regular**. The complete graph of order  $n$  is therefore a regular graph of degree  $n - 1$  and every cycle is 2-regular. In Figure 1.13 are shown all (non-isomorphic) regular graphs of orders 4 and 5, including the cycles  $C_4$  and  $C_5$  and the complete graphs  $K_4$  and  $K_5$ . Since no graph has an odd number of odd vertices, there is no 1-regular or 3-regular graph of order 5. Indeed, the pairs  $r, n$  of integers for which there exist  $r$ -regular graphs of order  $n$  are predictable.




Figure 1.13: The regular graphs of orders 4 and 5

**Theorem 1.7** *For integers  $r$  and  $n$ , there exists an  $r$ -regular graph of order  $n$  if and only if  $0 \leq r \leq n - 1$  and  $r$  and  $n$  are not both odd.*

**Proof.** That the conditions are necessary is an immediate consequence of Corollary 1.5 and the fact that  $0 \leq \deg v \leq n - 1$  for every vertex  $v$  in a graph of order  $n$ . For the converse, suppose that  $r$  and  $n$  are integers where  $0 \leq r \leq n - 1$  and  $r$  and  $n$  are not both odd. Assume first that  $r$  is even. If  $r = 0$ , then the graph of order  $n$  consisting of  $n$  isolated vertices is  $r$ -regular. So we may assume that  $r = 2k$  for some integer  $k \geq 1$  and  $n \geq 2k + 1$ . Let  $G$  be the graph with  $V(G) = \{v_1, v_2, \dots, v_n\}$  such that  $v_i$  ( $1 \leq i \leq n$ ) is adjacent

to  $v_{i\pm 1}, v_{i\pm 2}, \dots, v_{i\pm k}$  (subscripts expressed modulo  $n$ ). The resulting graph  $G$  is then an  $r$ -regular graph of order  $n$ . If  $r$  is odd, then  $n = 2\ell$  is even. Hence  $r = 2k + 1$  for some integer  $k$  with  $0 \leq k \leq \ell - 1$ . For the graph  $G$  in this case, the vertex  $v_i$  is adjacent to the  $2k$  vertices described above and adjacent as well to  $v_{i+\ell}$ . Again,  $G$  is an  $r$ -regular graph of order  $n$ . ■

The 4-regular and 5-regular graphs of order 10 constructed in the proof of Theorem 1.7 are shown in Figure 1.14.




Figure 1.14: 4-regular and 5-regular graphs of order 10

### The Petersen Graph

A 3-regular graph is also called a **cubic graph**. The graphs of Figure 1.9 are cubic as is the complete graph  $K_4$ . One of the best known cubic graphs is the **Petersen graph**, named for the Danish mathematician Julius Petersen whose 1891 research on regular graphs [186] is often credited as the beginning of the study of graphs as a theoretical subject. In fact, the Petersen graph is one of the best known graphs. Three different drawings of the Petersen graph are shown in Figure 1.15. We will have many occasions to encounter this graph.




Figure 1.15: Three drawings of the Petersen graph

## 1.5 Bipartite Graphs

Another class of graphs that we often encounter are the bipartite graphs. A graph  $G$  is **bipartite** if  $V(G)$  can be partitioned into two sets  $U$  and  $W$  (called

**partite sets**) so that every edge of  $G$  joins a vertex of  $U$  and a vertex of  $W$ . The graph  $G$  in Figure 1.16(a) is bipartite with partite sets  $\{v_1, v_3, v_5, v_7\}$  and  $\{v_2, v_4, v_6\}$ . This graph is redrawn in Figure 1.16(b) to see more clearly that it is bipartite. If  $G$  is an  $r$ -regular bipartite graph,  $r \geq 1$ , with partite sets  $U$  and  $W$ , then  $|U| = |W|$ . This follows since the size of  $G$  is  $r|U| = r|W|$ . For two nonempty sets  $X$  and  $Y$  of vertices in a graph  $G$ , the set

$$[X, Y] = \{xy : x \in X, y \in Y\}$$

consists of those edges joining a vertex of  $X$  and a vertex of  $Y$ . Thus, if  $G$  is a bipartite graph with partite sets  $U$  and  $W$ , then  $[U, W] = E(G)$ .




Figure 1.16: A bipartite graph

A graph  $G$  is a **complete bipartite graph** if  $V(G)$  can be partitioned into two sets  $U$  and  $W$  (called **partite sets** again) so that  $uw$  is an edge of  $G$  if and only if  $u \in U$  and  $w \in W$ . If  $|U| = s$  and  $|W| = t$ , then this complete bipartite graph has order  $s+t$  and size  $st$  and is denoted by  $K_{s,t}$  (or  $K_{t,s}$ ). The complete bipartite graph  $K_{1,t}$  is called a **star**. The complete bipartite graphs  $K_{1,3}$ ,  $K_{2,2}$ ,  $K_{2,3}$  and  $K_{3,3}$  are shown in Figure 1.17. Observe that  $K_{2,2} = C_4$ . The star  $K_{1,3}$  is sometimes referred to as a **claw**.




Figure 1.17: Complete bipartite graphs

Since the size of the complete bipartite graph  $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$  is  $\lfloor n/2 \rfloor \cdot \lceil n/2 \rceil = \lfloor n^2/4 \rfloor$ , there are bipartite graphs of order  $n$  and size  $\lfloor n^2/4 \rfloor$ . No bipartite graph of order  $n$  can have a larger size however.

**Theorem 1.8** *The size of every bipartite graph of order  $n$  is at most  $\lfloor n^2/4 \rfloor$ .*

**Proof.** Let  $G$  be bipartite graph of order  $n$  with partite sets  $U$  and  $W$ . Then  $|U| = x$  and  $|W| = n - x$  for some integer  $x$  with  $1 \leq x \leq n - 1$ . Hence the size of  $G$  is at most  $x(n - x)$ .

Since  $(n - 2x)^2 \geq 0$ , it follows that

$$n^2 \geq 4nx - 4x^2 = 4x(n - x)$$

and so  $x(n - x) \leq n^2/4$ . Since  $x(n - x)$  is an integer,  $x(n - x) \leq \lfloor n^2/4 \rfloor$ . ■

No bipartite graph  $G$  can contain a triangle  $H$ , for otherwise, at least two vertices  $u$  and  $v$  of  $H$  must belong to the same partite set and so  $u$  and  $v$  are not adjacent. Hence if  $G$  is a graph of order  $n \geq 3$  and size  $m \leq \lfloor n^2/4 \rfloor$ , then  $G$  need not contain a triangle. However, in 1907 the Dutch mathematician Willem Mantel [161] showed that any graph of order  $n$  with a larger size must contain a triangle.

**Theorem 1.9** *Every graph of order  $n \geq 3$  and size  $m > \lfloor n^2/4 \rfloor$  contains a triangle.*

**Proof.** First, observe that the result is true for  $n = 3$  and  $n = 4$ . Suppose, however, that the statement is false. Then there is a smallest integer  $n \geq 5$  and a graph  $G$  of order  $n$  and size  $m > \lfloor n^2/4 \rfloor$  not containing a triangle. Let  $uv$  be an edge of  $G$ . Since  $G$  contains no triangle, there is no vertex in  $G$  adjacent to both  $u$  and  $v$ . Hence,  $(\deg u - 1) + (\deg v - 1) \leq n - 2$  and so  $\deg u + \deg v \leq n$ . Let  $G' = G - u - v$ . Since  $G'$  is a subgraph of  $G$ , it follows that  $G'$  also does not contain a triangle. Furthermore,  $G'$  has order  $n - 2$  and size

$$m' = m - (\deg u + \deg v) + 1 > \lfloor n^2/4 \rfloor - n + 1.$$

Thus,

$$m' > \frac{n^2}{4} - n + 1 = \frac{n^2 - 4n + 4}{4} = \frac{(n - 2)^2}{4}.$$

From the defining property of the graph  $G$ , it follows that  $G'$  contains a triangle, producing a contradiction. ■

Therefore, from Theorems 1.8 and 1.9, it follows that every graph of order  $n \geq 3$  and size  $m > \lfloor n^2/4 \rfloor$  not only fails to be bipartite, it must, in fact, contain a triangle.

### Complete Multipartite Graphs

Bipartite graphs belong to a more general class of graphs. For an integer  $k \geq 1$ , a graph  $G$  is a  **$k$ -partite graph** if  $V(G)$  can be partitioned into  $k$  subsets  $V_1, V_2, \dots, V_k$  (again called **partite sets**) such that every edge of  $G$  joins vertices in two different partite sets. A 1-partite graph is then an empty graph and a 2-partite graph is bipartite. A **complete  $k$ -partite graph**  $G$  is a  $k$ -partite graph with the property that two vertices are adjacent in  $G$  if and only if the vertices belong to different partite sets. If  $|V_i| = n_i$  for  $1 \leq i \leq k$ , then  $G$  is denoted by  $K_{n_1, n_2, \dots, n_k}$  (the order in which the numbers  $n_1, n_2, \dots, n_k$

are written is not important). If  $n_i = 1$  for all  $i$  ( $1 \leq i \leq k$ ), then  $G$  is the complete graph  $K_k$ . A **complete multipartite graph** is a complete  $k$ -partite graph for some integer  $k \geq 2$ . Some complete multipartite graphs are shown in Figure 1.18




Figure 1.18: Some complete multipartite graphs

## 1.6 Operations on Graphs

There are many ways of producing a new graph from one or more given graphs. The most common of these is the complement of a graph.

### The Complement of a Graph

The **complement**  $\overline{G}$  of a graph  $G$  is that graph with vertex set  $V(G)$  such that two vertices are adjacent in  $\overline{G}$  if and only if these vertices are not adjacent in  $G$ . Any isomorphism from a graph  $G$  to a graph  $H$  is also an isomorphism from  $\overline{G}$  to  $\overline{H}$ . Consequently,  $\overline{G} \cong \overline{H}$  if and only if  $G \cong H$ . If  $G$  is a graph of order  $n$  and size  $m$ , then  $\overline{G}$  is a graph of order  $n$  and size  $\binom{n}{2} - m$ . A graph  $G$  and its complement are shown in Figure 1.19. The complement  $\overline{K_n}$  of the complete graph  $K_n$  is the empty graph of order  $n$ .




Figure 1.19: A graph and its complement

A graph  $G$  is **self-complementary** if  $G$  is isomorphic to  $\overline{G}$ . Certainly, if  $G$  is a self-complementary graph of order  $n$ , then its size is  $m = \binom{n}{2}/2 = n(n-1)/4$ . Since only one of  $n$  and  $n-1$  is even, either  $4 \mid n$  or  $4 \mid (n-1)$ ; that is, if  $G$  is a self-complementary graph of order  $n$ , then either  $n \equiv 0 \pmod{4}$  or

$n \equiv 1 \pmod{4}$ . (See Exercise 29.) The self-complementary graphs of order 5 or less are shown in Figure 1.20.




Figure 1.20: The self-complementary graphs of order 5 or less

### The Union and Join of Graphs

We next describe some common binary operations defined on graphs. This discussion introduces notation that will be especially useful in giving examples. Over the years, different authors have used different notation for the operations we are about to describe. In the following definitions, we assume that  $G_1$  and  $G_2$  are two graphs with disjoint vertex sets.

The **union**  $G = G_1 + G_2$  of  $G_1$  and  $G_2$  has vertex set  $V(G) = V(G_1) \cup V(G_2)$  and edge set  $E(G) = E(G_1) \cup E(G_2)$ . The union  $G + G$  of two disjoint copies of  $G$  is denoted by  $2G$ . Indeed, if a graph  $G$  consists of  $k$  ( $\geq 2$ ) disjoint copies of a graph  $H$ , then we write  $G = kH$ . The graph  $2K_1 + 3K_2 + K_{1,3}$  is shown in Figure 1.21(a). The **join**  $G = G_1 \vee G_2$  of  $G_1$  and  $G_2$  has vertex set  $V(G) = V(G_1) \cup V(G_2)$  and edge set

$$E(G) = E(G_1) \cup E(G_2) \cup \{uv : u \in V(G_1), v \in V(G_2)\}.$$

Using the join operation, we see that  $\overline{K_s} \vee \overline{K_t} = K_{s,t}$ . Another illustration is given in Figure 1.21(b).




Figure 1.21: The union and join of graphs

## The Cartesian Product of Graphs

The **Cartesian product**  $G$  of two graphs  $G_1$  and  $G_2$ , commonly denoted by  $G_1 \square G_2$  or  $G_1 \times G_2$ , has vertex set

$$V(G) = V(G_1) \times V(G_2),$$

where two distinct vertices  $(u, v)$  and  $(x, y)$  of  $G_1 \square G_2$  are adjacent if either

- (1)  $u = x$  and  $vy \in E(G_2)$  or (2)  $v = y$  and  $ux \in E(G_1)$ .

A convenient way of drawing  $G_1 \square G_2$  is to first place a copy of  $G_2$  at each vertex of  $G_1$  (see Figure 1.22(b)) and then join corresponding vertices of  $G_2$  in those copies of  $G_2$  placed at adjacent vertices of  $G_1$  (see Figure 1.22(c)). Equivalently,  $G_1 \square G_2$  can be constructed by placing a copy of  $G_1$  at each vertex of  $G_2$  and adding the appropriate edges. As expected,  $G_1 \square G_2 \cong G_2 \square G_1$  for all graphs  $G_1$  and  $G_2$ .





Figure 1.22: The Cartesian product of two graphs

## Hypercubes

An important class of graphs is defined in terms of Cartesian products. The  $n$ -**cube**  $Q_n$  is  $K_2$  if  $n = 1$ , while for  $n \geq 2$ ,  $Q_n$  is defined recursively as the Cartesian product  $Q_{n-1} \square K_2$  of  $Q_{n-1}$  and  $K_2$ . The  $n$ -cube can also be defined as that graph whose vertex set is the set of ordered  $n$ -tuples  $(a_1, a_2, \dots, a_n)$  or  $a_1 a_2 \dots a_n$  where  $a_i$  is 0 or 1 for  $1 \leq i \leq n$  (commonly called  $n$ -**bit strings**), such that two vertices are adjacent if and only if the corresponding ordered  $n$ -tuples differ at precisely one coordinate. The graph  $Q_n$  is an  $n$ -regular graph of order  $2^n$ . The  $n$ -cubes for  $n = 1, 2, 3$  are shown in Figure 1.23, where their vertices are labeled by  $n$ -bit strings. The graphs  $Q_n$  are also called **hypercubes**.

## 1.7 Degree Sequences

We saw in the First Theorem of Graph Theory (Theorem 1.4) that the sum of the degrees of the vertices of a graph  $G$  is twice the size of  $G$  and in Corollary 1.5

Figure 1.23: The  $n$ -cubes for  $n = 1, 2, 3$ 

that  $G$  must have an even number of odd vertices. We have also described conditions under which a regular graph of order  $n$  can exist. We now consider the degrees of the vertices of a graph in more detail.

A sequence  $d_1, d_2, \dots, d_n$  of nonnegative integers is called a **degree sequence** of a graph  $G$  of order  $n$  if the vertices of  $G$  can be labeled  $v_1, v_2, \dots, v_n$  so that  $\deg v_i = d_i$  for  $1 \leq i \leq n$ . For example, a degree sequence of the graph  $G$  of Figure 1.24 is 4, 3, 2, 2, 1 (or 1, 2, 2, 3, 4 or 2, 1, 4, 2, 3, etc.). We commonly write the degree sequence of a graph as a nonincreasing sequence.




Figure 1.24: A degree sequence of a graph

A finite sequence  $s$  of nonnegative integers is a **graphical sequence** if  $s$  is a degree sequence of some graph. Thus, 4, 3, 2, 2, 1 is graphical. There are some obvious necessary conditions for a sequence  $s : d_1, d_2, \dots, d_n$  of  $n$  nonnegative integers to be graphical. While the conditions that  $d_i \leq n - 1$  for all  $i$  ( $1 \leq i \leq n$ ) and  $\sum_{i=1}^n \deg v_i$  is even are necessary for  $s$  to be graphical, they are not sufficient. For example, the sequence 3, 3, 3, 1 satisfies both conditions but it is not graphical, for if three vertices of a graph of order 4 have degree 3 then the remaining vertex must have degree 3 as well.

It is not all that unusual for a graphical sequence to be the degree sequence of more than one graph. For example, the graphical sequence 3, 2, 2, 2, 1 is the degree sequence of the two (non-isomorphic) graphs in Figure 1.25. On the other hand, each of the 18 graphs in Figure 1.7 has a degree sequence possessed by no other graph.




Figure 1.25: Two graphs with the same degree sequence



## 2-Switches

While two graphs with the same degree sequence need not be isomorphic, each can be obtained from the other by a sequence of edge shifts where at each step, two nonadjacent edges in some graph  $F$  are deleted and two nonadjacent edges in  $\overline{F}$  are added to  $F$  such that the four edges involved are incident with the same four vertices.

Let  $H$  be a graph containing four distinct vertices  $u, v, w$  and  $x$  such that  $uv, wx \in E(H)$  and  $uw, vx \notin E(H)$ . The process of deleting the edges  $uv$  and  $wx$  from  $H$  and adding  $uw$  and  $vx$  to  $H$  is referred to as a **2-switch** in  $H$  (see Figure 1.26, where a dashed line means no edge). This produces a new graph  $G$  having the same degree sequence as  $H$ . Of course, if  $G$  can be produced from  $H$  by a 2-switch, then  $H$  can be obtained from  $G$  by a 2-switch.




Figure 1.26: A 2-switch in a graph

**Theorem 1.10** *Let  $s : d_1, d_2, \dots, d_n$  be a graphical sequence with  $\Delta = d_1 \geq d_2 \geq \dots \geq d_n$  and let  $\mathcal{G}_s$  be the set of all graphs  $F$  with degree sequence  $s$  such that  $V(F) = \{v_1, v_2, \dots, v_n\}$  where  $\deg v_i = d_i$  for  $1 \leq i \leq n$ . Then every graph  $H \in \mathcal{G}_s$  can be transformed into a graph  $G \in \mathcal{G}_s$  by a sequence of 2-switches such that  $N_G(v_1) = \{v_2, v_3, \dots, v_{\Delta+1}\}$ .*

**Proof.** Suppose that this statement is false. Let  $W = \{v_2, v_3, \dots, v_{\Delta+1}\}$ . Among all graphs into which  $H$  can be transformed, let  $G$  be one for which the sum of the subscripts of the vertices in  $N_G(v_1)$  is minimum. Since  $N_G(v_1) \neq W$ , the vertex  $v_1$  is adjacent to a vertex  $v_k$  and is not adjacent to a vertex  $v_j$  with  $j < k$  and so  $d_j \geq d_k$ . Consequently, there is a vertex  $v_\ell$  such that  $v_j v_\ell \in E(G)$  and  $v_k v_\ell \notin E(G)$ . Replacing the edges  $v_1 v_k$  and  $v_j v_\ell$  by  $v_1 v_j$  and  $v_k v_\ell$  is a 2-switch in  $G$  that produces a graph  $G_1 \in \mathcal{G}_s$  for which the sum of the subscripts of the vertices in  $N_{G_1}(v_1)$  is less than that of  $N_G(v_1)$ . Consequently,  $H$  can be transformed into  $G_1$  and so this is a contradiction. ■

One consequence of Theorem 1.10 is that every two graphs with the same degree sequence are related in terms of 2-switches. The following theorem appeared in the book *Graphs and Hypergraphs* by Claude Berge [25].

**Theorem 1.11** *If  $G$  and  $H$  are two graphs with the same degree sequence, then  $H$  can be transformed into  $G$  by a (possibly empty) sequence of 2-switches.*

**Proof.** We proceed by induction on the order  $n$  of  $G$  and  $H$ . If  $n \leq 4$ , then the result is immediate. For a given integer  $n \geq 5$ , assume that every two graphs of order  $n - 1$  with the same degree sequence can be transformed into each other by a sequence of 2-switches. Let  $s : d_1, d_2, \dots, d_n$  be a graphical sequence with  $\Delta = d_1 \geq d_2 \geq \dots \geq d_n$  and let  $\mathcal{G}_s$  be the set of all graphs  $F$  with degree sequence  $s$  such that  $V(F) = \{v_1, v_2, \dots, v_n\}$  where  $\deg v_i = d_i$  for  $1 \leq i \leq n$ . Let  $W = \{v_2, v_3, \dots, v_{\Delta+1}\}$ . Let  $G$  and  $H$  be two graphs in  $\mathcal{G}_s$ . By Theorem 1.10,  $G$  can be transformed into a graph  $G_1 \in \mathcal{G}_s$  such that  $N_{G_1}(v_1) = W$  and  $H$  can be transformed into a graph  $H_1 \in \mathcal{G}_s$  such that  $N_{H_1}(v_1) = W$ . Since  $G_1 - v_1$  and  $H_1 - v_1$  are two graphs of order  $n - 1$  with the same degree sequence, it follows by the induction hypothesis that  $G_1 - v_1$  can be transformed into  $H_1 - v_1$  by a sequence of 2-switches. Hence  $G_1$  can be transformed into  $H_1$  by a sequence of 2-switches and so  $G$  can be transformed into  $H$  by a sequence of 2-switches. ■

### The Havel–Hakimi Theorem

There are necessary and sufficient conditions for a finite sequence of non-negative integers to be graphical. One of these is due to Václav Havel [124] and S. Louis Hakimi [117] and is a consequence of Theorem 1.10. This result is often referred to as the Havel-Hakimi Theorem, despite the fact that Havel and Hakimi gave independent proofs and wrote separate papers containing this theorem.

**Theorem 1.12 (Havel–Hakimi Theorem)** *A sequence  $s : d_1, d_2, \dots, d_n$  of nonnegative integers with  $\Delta = d_1 \geq d_2 \geq \dots \geq d_n$  and  $\Delta \geq 1$  is graphical if and only if the sequence*

$$s_1 : d_2 - 1, d_3 - 1, \dots, d_{\Delta+1} - 1, d_{\Delta+2}, \dots, d_n$$

*is graphical.*

**Proof.** First, assume that  $s_1$  is graphical. Then there exists a graph  $G_1$  of order  $n - 1$  such that  $s_1$  is a degree sequence of  $G_1$ . Thus, the vertices of  $G_1$  can be labeled as  $v_2, v_3, \dots, v_n$  so that

$$\deg_{G_1} v_i = \begin{cases} d_i - 1 & \text{if } 2 \leq i \leq \Delta + 1 \\ d_i & \text{if } \Delta + 2 \leq i \leq n. \end{cases}$$

A new graph  $G$  can now be constructed by adding a new vertex  $v_1$  to  $G_1$  together with the  $\Delta$  edges  $v_1 v_i$  for  $2 \leq i \leq \Delta + 1$ . Since  $\deg_G v_i = d_i$  for  $1 \leq i \leq n$ , it follows that  $s : \Delta = d_1, d_2, \dots, d_n$  is a degree sequence of  $G$  and so  $s$  is graphical.

Conversely, let  $s$  be a graphical sequence. By Theorem 1.10, there exists a graph  $G$  of order  $n$  having degree sequence  $s$  with  $V(G) = \{v_1, v_2, \dots, v_n\}$  such that  $\deg v_i = d_i$  for all  $i$  ( $1 \leq i \leq n$ ) where  $N_G(v_1) = \{v_2, v_3, \dots, v_{\Delta+1}\}$ . Then  $G - v_1$  has degree sequence  $s_1$ . ■

Theorem 1.12 actually provides us with an algorithm for determining whether a given finite sequence of nonnegative integers is graphical. If, upon repeated application of Theorem 1.12, we arrive at a sequence every term of which is 0, then the original sequence is graphical. On the other hand, if we arrive at a sequence containing a negative integer, then the given sequence is not graphical.

We now illustrate Theorem 1.12 with the sequence

$$s : 5, 3, 3, 3, 3, 2, 2, 1, 1, 1.$$

After one application of Theorem 1.12 (deleting 5 from  $s$  and subtracting 1 from the next five terms), we obtain

$$s'_1 : 2, 2, 2, 2, 1, 2, 2, 1, 1, 1.$$


Reordering this sequence, we have

$$s_1 : 2, 2, 2, 2, 2, 1, 1, 1, 1, 1.$$

Continuing in this manner, we get

$$\begin{aligned} s'_2 &: 1, 1, 2, 2, 2, 1, 1, 1, 1 \\ s_2 &: 2, 2, 2, 1, 1, 1, 1, 1, 1 \\ s'_3 = s_3 &: 1, 1, 1, 1, 1, 1, 1, 1 \\ s'_4 &: 0, 1, 1, 1, 1, 1, 1 \\ s_4 &: 1, 1, 1, 1, 1, 1, 0 \\ s'_5 &: 0, 1, 1, 1, 1, 0 \\ s_5 &: 1, 1, 1, 1, 0, 0 \\ s'_6 &: 0, 1, 1, 0, 0 \\ s_6 &: 1, 1, 0, 0, 0 \\ s'_7 = s_7 &: 0, 0, 0, 0. \end{aligned}$$

Therefore,  $s$  is graphical. Of course, if we observe that some sequence prior to  $s_7$  is graphical, then we can conclude by Theorem 1.12 that  $s$  is graphical. For example, the sequence  $s_3$  is clearly graphical since it is the degree sequence of the graph  $G_3 = 4K_2$  in Figure 1.27. By Theorem 1.12, each of the sequences  $s_2, s_1$  and  $s$  is also graphical. To construct a graph with degree sequence  $s_2$ , we proceed in reverse from  $s'_3 = s_3$  to  $s_2$ , observing that a vertex should be added to  $G_3$  so that it is adjacent to two vertices of degree 1. We thus obtain a graph  $G_2$  with degree sequence  $s_2$  (or  $s'_2$ ). Proceeding from  $s'_2$  to  $s_1$ , we again add a new vertex joining it to two vertices of degree 1 in  $G_2$ . This gives a graph  $G_1$  with degree sequence  $s_1$  (or  $s'_1$ ). Finally, we obtain a graph  $G$  with degree sequence  $s$  by considering  $s'_1$ ; that is, we add a new vertex to  $G_1$  joining it to vertices of degrees 2, 2, 2, 2, 1. This procedure is illustrated in Figure 1.27.

Figure 1.27: Construction of a graph  $G$  with a given degree sequence

It should be pointed out that the graph  $G$  in Figure 1.27 is not the only graph with degree sequence  $s$ . However, there are graphs that cannot be produced by the method used to construct the graph  $G$  in Figure 1.27. For example, the graph  $H$  of Figure 1.28 is such a graph.




Figure 1.28: A graph that cannot be constructed by the method following Theorem 1.12

### The Erdős–Gallai Theorem

Suppose that  $s : d_1, d_2, \dots, d_n$  is a graphical sequence with  $d_1 \geq d_2 \geq \dots \geq d_n$ . Then there exists a graph  $G$  of order  $n$  with  $V(G) = \{v_1, v_2, \dots, v_n\}$  such that  $\deg v_i = d_i$  for all  $i$  ( $1 \leq i \leq n$ ). Of course, the sum  $\sum_{i=1}^n d_i$  is even. Let  $k$  be an integer with  $1 \leq k \leq n - 1$ . Suppose that  $V_1 = \{v_1, v_2, \dots, v_k\}$  and  $V_2 = \{v_{k+1}, v_{k+2}, \dots, v_n\}$ . Now the sum  $\sum_{i=1}^k d_i$  counts every edge in  $G[V_1]$  twice and counts each edge in  $[V_1, V_2]$  once. The size of  $G[V_1]$  is at most  $\binom{k}{2} = k(k-1)/2$ , while for each  $i$  ( $k+1 \leq i \leq n$ ) the number of edges joining  $v_i$  and  $V_1$  is at most  $\min\{k, d_i\}$ . Thus,

$$\sum_{i=1}^k d_i \leq 2 \binom{k}{2} + \sum_{i=k+1}^n \min\{k, d_i\} = k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\}. \quad (1.1)$$

Hence, for every graphical sequence  $s$ , we must have both that  $\sum_{i=1}^n d_i$  is even and (1.1) is satisfied for every integer  $k$  with  $1 \leq k \leq n - 1$ . These conditions are not only necessary for a sequence of nonnegative integers to be graphical, they are sufficient as well. Since the proof of the sufficiency is technical, we omit this. This result was established by Paul Erdős and Tibor Gallai [81], who were introduced to graph theory as youngsters by Dénes König, author of the first book on graph theory. In fact, Gallai was König's only doctoral student.

**Theorem 1.13 (Erdős–Gallai Theorem)** *A sequence  $s : d_1, d_2, \dots, d_n$  ( $n \geq 2$ ) of nonnegative integers with  $d_1 \geq d_2 \geq \dots \geq d_n$  is graphical if and only if  $\sum_{i=1}^n d_i$  is even and for each integer  $k$  with  $1 \leq k \leq n-1$ ,*

$$\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\}.$$

### Irregular Graphs

According to Theorem 1.7, there is an  $r$ -regular graph of order  $n$  if and only if  $0 \leq r \leq n-1$  and  $rn$  is even. At the other extreme are nontrivial graphs, no two vertices of which have the same degree. A nontrivial graph  $G$  is **irregular** if  $\deg u \neq \deg v$  for every two vertices  $u$  and  $v$  of  $G$ . Actually, no graph has this property.

**Theorem 1.14** *No graph is irregular.*

**Proof.** Assume, to the contrary, that there exists an irregular graph  $G$  of order  $n \geq 2$ . Since the degree of every vertex of  $G$  is one of the  $n$  integers  $0, 1, \dots, n-1$ , each of these integers is the degree of exactly one vertex. Thus, we may assume that  $V(G) = \{v_1, v_2, \dots, v_n\}$  where  $\deg v_i = i-1$  for  $1 \leq i \leq n$ . Since  $\deg v_1 = 0$ , the vertex  $v_1$  is isolated in  $G$  and since  $\deg v_n = n-1$ , it follows that  $v_n$  is adjacent to  $v_1$ . This is a contradiction. ■

By Theorem 1.14, for each integer  $n \geq 2$ , there is no graph of order  $n$  whose  $n$  vertices have distinct degrees. Since the degrees of the vertices of the graph  $G$  of Figure 1.24 are 4, 3, 2, 2, 1, it is possible for  $n-1$  vertices of a graph of order  $n$  to have distinct degrees. A graph  $G$  of order  $n \geq 2$  is **nearly irregular** if exactly two vertices of  $G$  have the same degree.

**Theorem 1.15** *For every integer  $n \geq 2$ , there are exactly two nearly irregular graphs of order  $n$ .*

**Proof.** First, observe that if  $G$  is a nearly irregular graph of order  $n$ , then  $G$  cannot contain a vertex of degree 0 and a vertex of degree  $n-1$ . Thus, either each vertex of  $G$  has one of the degrees  $1, 2, \dots, n-1$  or each vertex of  $G$  has one of the degrees  $0, 1, \dots, n-2$ . Furthermore, if  $G$  is nearly irregular, then so is  $\overline{G}$ .

We show by induction that for each integer  $n \geq 2$  there are exactly two nearly irregular graphs of order  $n$ . Since  $K_2$  and  $\overline{K}_2$  are nearly irregular, the result holds for  $n = 2$ . Assume for an integer  $n \geq 3$  that there are exactly two nearly irregular graphs of order  $n-1$ . Necessarily, one of these graphs is a graph  $F$  with  $\Delta(F) = n-2$  and  $\delta(F) = 1$  while the other is  $\overline{F}$  where  $\Delta(\overline{F}) = n-3$  and  $\delta(\overline{F}) = 0$ . Then  $H = F + K_1$  and  $\overline{H}$  are nearly irregular graphs of order

$n$ , where  $\Delta(H) = n - 2$  and  $\delta(H) = 0$ . We claim that these are the only nearly irregular graphs of order  $n$ . Assume, to the contrary, that there is a third graph  $G$  of order  $n$  that is nearly irregular. Then either  $\Delta(G) = n - 1$  or  $\delta(G) = 0$ , say the latter. Then  $G = G_1 + K_1$ , where  $G_1$  is the nearly irregular graph of order  $n - 1$  with  $\Delta(G_1) = n - 2$ . However then,  $G_1 \cong H$ , which is a contradiction. ■

It follows by Theorem 1.15 then that for each integer  $n \geq 2$ , there exist exactly two distinct graphical sequences of length  $n$  having exactly two equal terms.

## 1.8 Multigraphs

There are occasions when a graph is not the appropriate structure to model a particular situation. For example, suppose that we are considering various locations in a certain community and there are roads between some pairs of locations that do not pass through any other location. Although this situation may be represented by a graph, there may be some characteristics in this network of roads that are not captured by a graph. For example, suppose that there are pairs of locations connected by two or more roads (not passing through any other location) and this information is important to us.

In the definition of a graph  $G$ , every two distinct vertices are joined by either one edge or no edge of  $G$ . There will be occasions when we will want to permit more than one edge to join two vertices. A **multigraph** is a nonempty set of vertices, every two of which are joined by a finite number of edges. Hence a multigraph  $H$  may be expressed as  $H = (V, E)$ , where  $E$  is a multiset of 2-element subsets of  $V$ . Two or more edges that join the same pair of distinct vertices are called **parallel edges**. The **underlying graph** of a multigraph  $H$  is that graph  $G$  for which  $V(G) = V(H)$  and  $uv \in E(G)$  if  $u$  and  $v$  are joined by at least one edge in  $H$ .

An edge joining a vertex to itself is called a **loop**. Structures that permit both parallel edges and loops (including parallel loops) are sometimes called **pseudographs**. For emphasis then, every two vertices of a graph are joined by at most one edge and loops are not permitted. In a multigraph, every two vertices are permitted to be joined by more than one edge but this is not required. Also, no multigraph contains a loop. In a pseudograph, every two vertices are permitted to be joined by more than one edge and loops are permitted. However, parallel edges and loops are not required in pseudographs. There are authors who refer to multigraphs or pseudographs as graphs and those who refer to what we call graphs as **simple graphs**. Consequently, when reading any material written on graph theory, it is essential that there is a clear understanding of how the term *graph* is being used. According to the terminology introduced here then, every multigraph is a pseudograph and every graph is both a multigraph and a pseudograph.

In Figure 1.29,  $H_1$  and  $H_4$  are multigraphs while  $H_2$  and  $H_3$  are pseudographs. Of course,  $H_1$  and  $H_4$  are also pseudographs while  $H_4$  is the only graph

in Figure 1.29. For a vertex  $v$  in a multigraph  $G$ , the **degree**  $\deg v$  of  $v$  in  $G$  is the number of edges of  $G$  incident with  $v$ . In a pseudograph, each loop at a vertex contributes 2 to its degree. For the pseudograph  $H_3$  of Figure 1.29,  $\deg u = 5$  and  $\deg v = 2$ .




Figure 1.29: Multigraphs and pseudographs

The degree sequence of the multigraph  $G_1$  in Figure 1.30 is 5, 4, 3 and that of the multigraph  $G_2$  is 4, 3, 2, 1. That is,  $G_1$  and  $G_2$  are irregular multigraphs.




Figure 1.30: Two irregular multigraphs

The multigraphs  $G_1$  and  $G_2$  in Figure 1.30 illustrate the following result.

**Theorem 1.16** *For every connected graph  $G$  of order at least 3, there exists an irregular multigraph whose underlying graph is  $G$ .*

**Proof.** Let  $E(G) = \{e_1, e_2, \dots, e_m\}$ , where  $e_i = u_i v_i$  for  $i = 1, 2, \dots, m$ . Replacing  $e_i$  by  $2^{i-1}$  parallel edges joining  $u_i$  and  $v_i$  produces a multigraph  $H$ . Since every two vertices of  $G$  have distinct sets of edges incident with them and every positive integer has a unique base 2 representation, their degrees in  $H$  are distinct and so  $H$  is irregular. ■

When describing walks in multigraphs or in pseudographs, it is often necessary to list edges in the sequence as well as vertices in order to specify the edges being used in the walk. For example,

$$W = (u, e_1, u, e_3, v, e_6, w, e_6, v, e_7, w)$$

is a  $u - w$  walk in the pseudograph  $G$  of Figure 1.31.




Figure 1.31: Walks in a pseudograph



## Exercises for Chapter 1

### Section 1.1. Graphs

1. An electronics company keeps on hand wire segments of a fixed length and of different colors for various purposes. Each wire is either colored blue (b), green (g), purple (p), red (r), silver (s), white (w) or yellow (y). The company has many wire segments of each color. All of the wire segments have been randomly stored in a large barrel. Eight handfuls of wires are removed from the barrel and each collection of wires is placed in a box. The boxes are denoted by  $B_i$  ( $1 \leq i \leq 8$ ). The colors of the wire segments in each box are:

$$\begin{array}{llll} B_1 = \{b, r\} & B_2 = \{p, r, s, w\} & B_3 = \{p, w, y\} & B_4 = \{g, r, y\} \\ B_5 = \{g\} & B_6 = \{b, g, y\} & B_7 = \{g, p, s, w, y\} & B_8 = \{s, w, y\}. \end{array}$$

We are interested in those pairs of boxes containing at least one wire segment of the same color. Model this situation by a graph.

2. A graph  $G = (V, E)$  of order 8 has the power set of the set  $S = \{1, 2, 3\}$  as its vertex set, that is,  $V$  is the set of all subsets of  $S$ . Two vertices  $A$  and  $B$  of  $V$  are adjacent if  $A \cap B = \emptyset$ . Draw the graph  $G$ , determine the degree of each vertex of  $G$  and determine the size of  $G$ .

### Section 1.2. The Degree of a Vertex

3. A graph  $G$  of order 26 and size 58 has 5 vertices of degree 4, 6 vertices of degree 5 and 7 vertices of degree 6. The remaining vertices of  $G$  all have the same degree. What is this degree?
4. A graph  $G$  has order  $n = 3k + 3$  for some positive integer  $k$ . Every vertex of  $G$  has degree  $k + 1$ ,  $k + 2$  or  $k + 3$ . Prove that  $G$  has at least  $k + 3$  vertices of degree  $k + 1$  or at least  $k + 1$  vertices of degree  $k + 2$  or at least  $k + 2$  vertices of degree  $k + 3$ .
5. The degree of every vertex of a graph  $G$  is one of three consecutive integers. For each degree  $x$ , the graph  $G$  contains exactly  $x$  vertices of degree  $x$ . Prove that for every graph  $G$  with this property, two-thirds of the vertices of  $G$  have odd degree.
6. Show for every positive integer  $k$  that there exists a graph  $G$  of order  $2k$  containing two vertices of degree  $i$  for each  $i = 1, 2, \dots, k$ .

### Section 1.3. Isomorphic Graphs

7. Consider the pairs  $G_1, G_2$  and  $H_1, H_2$  of graphs in Figure 1.32.

- (a) Determine whether  $G_1 \cong G_2$ .  
 (b) Determine whether  $H_1 \cong H_2$ .




Figure 1.32: The graphs  $G_1, G_2, H_1, H_2$  in Exercise 7

8. (a) Determine all non-isomorphic graphs of order 5.  
 (b) Determine the minimum size of a graph  $G$  of order 5 such that every graph of order 5 and size 5 is isomorphic to some subgraph of  $G$ .
9. (a) Let  $G$  and  $H$  be two isomorphic graphs where one or more vertices of  $G$  (and of  $H$ ) have degree  $r$ . Let  $S$  be the set of vertices of degree  $r$  in  $G$  and  $T$  be the set of vertices of degree  $r$  in  $H$ . Prove that  $G[S] \cong H[T]$ .  
 (b) Use the result in (a) to show that the graphs  $G$  and  $H$  in Figure 1.33 are not isomorphic.




Figure 1.33: The graphs  $G$  and  $H$  in Exercise 9

10. (a) Give an example of three graphs of size 3, no two of which are isomorphic, such that in each graph, every two edge-induced subgraphs of the same size are isomorphic.  
 (b) Give an example of two graphs  $H$  and  $G$  of the same order and two spanning subgraphs  $F_1$  and  $F_2$  of  $G$  such that  $F_i \cong H$  for  $i = 1, 2$  and  $G - E(F_1) \not\cong G - E(F_2)$ .

### Section 1.4. Regular Graphs

11. Show that if  $G$  is a nonregular graph of order  $n$  and size  $rn/2$  for some integer  $r$  with  $1 \leq r \leq n-2$ , then  $\Delta(G) - \delta(G) \geq 2$ .
12. For each integer  $k \geq 2$ , give an example of  $k$  non-isomorphic regular graphs, all of the same order and same size.
13. Give an example of a nonregular graph  $G$  containing an edge  $e$  and a vertex  $u$  such that  $G - e$  and  $G - u$  are both regular.
14. (a) Give an example of two non-isomorphic regular graphs  $G_1$  and  $G_2$  of the same order and same size such that (1) for every two vertices  $v_1 \in V(G_1)$  and  $v_2 \in V(G_2)$ ,  $G_1 - v_1 \not\cong G_2 - v_2$  and (2) there exist 2-element subsets  $S_1 \subseteq V(G_1)$  and  $S_2 \subseteq V(G_2)$  such that  $G_1 - S_1 \cong G_2 - S_2$ .  
 (b) Give an example of two non-isomorphic regular graphs  $H_1$  and  $H_2$  of the same order and same size such that (1) for every 2-element subsets  $S_1 \subseteq V(H_1)$  and  $S_2 \subseteq V(H_2)$ ,  $H_1 - S_1 \not\cong H_2 - S_2$  and there exist 3-element subsets  $S'_1 \subseteq V(H_1)$  and  $S'_2 \subseteq V(H_2)$  such that  $H_1 - S'_1 \cong H_2 - S'_2$ .
15. Prove for every graph  $G$  and every integer  $r \geq \Delta(G)$  that there exists an  $r$ -regular graph containing  $G$  as an induced subgraph.
16. Let  $G$  be a graph of order  $n$  all of whose vertices have degree  $r$ , where  $r$  is a positive integer, except for exactly one vertex of each of the degrees  $r-1, r-2, \dots, r-j$ , where  $1 < j < r$ . Show, in fact, that there exists an  $r$ -regular graph of order  $2n$  containing  $G$  as an induced subgraph.
17. Let  $S = \{1, 2, 3, 4, 5\}$ . The vertex set of a graph  $G$  is the set of 2-element subsets of  $S$ . Two vertices of  $G$  are adjacent if the vertices are disjoint. What familiar graph is  $G$ ?
18. For positive integers  $k$  and  $n$  with  $n > 2k$ , the graph  $G_{n,k}$  is that graph whose vertices are the  $k$ -element subsets of an  $n$ -element set  $S = \{1, 2, \dots, n\}$  and where two vertices ( $k$ -element subsets)  $A$  and  $B$  are adjacent if  $A$  and  $B$  are disjoint. The graph  $G_{n,k}$  is called the **Kneser graph**.  
 (a) Determine the graphs  $G_{6,1}$  and  $G_{5,2}$ .  
 (b) Show that  $G_{n,k}$  is an  $r$ -regular graph for some integer  $r$ .

**Section 1.5 Bipartite Graphs**

19. A bipartite graph  $G$  of order  $n$  has partite sets  $U$  and  $W$  where  $|U| = 10$ . Every vertex of  $U$  has degree 6. In  $W$ , there are four vertices of degree 2 and three vertices of degree 4. All other vertices of  $G$  have degree 8. What is  $n$ ?
20. Show for each integer  $n \geq 2$  that there is exactly one bipartite graph of order  $n$  having size  $\lfloor n^2/4 \rfloor$ .
21. Prove for a 3-partite graph of order  $n = 3k$  and size  $m$  that  $m \leq 3k^2$ .
22. Let  $G$  be a nonempty graph with the property that whenever  $uv \notin E(G)$  and  $vw \notin E(G)$ , then  $uw \notin E(G)$ . Prove that  $G$  is a complete multipartite graph.

**Section 1.6. Operation on Graphs**

23. Determine all bipartite graphs  $G$  such that  $\overline{G}$  is bipartite.
24. Let  $G$  be a graph of odd order  $n = 2k + 1 \geq 3$  for some positive integer  $k$ . Prove that if the vertices of  $G$  have exactly the same degrees as the vertices of  $\overline{G}$ , then  $G$  has an odd number of vertices of degree  $k$ .
25. (a) Show that there are exactly two 4-regular graphs  $G$  of order 7.  
(b) How many 6-regular graphs of order 9 are there?
26. Prove that there is no regular self-complementary graph of even order.
27. We have seen that  $C_5$  is a self-complementary graph. Therefore, there is a regular self-complementary graph of order 5. Show that there is a regular self-complementary graph of order  $5^n$  for every positive integer  $n$ .
28. Let  $G_1$  and  $G_2$  be self-complementary graphs, where  $G_2$  has even order  $n$ . Let  $G$  be the graph obtained from  $G_1$  and  $G_2$  by joining each vertex of  $G_2$  whose degree is less than  $n/2$  to every vertex of  $G_1$ . Show that  $G$  is self-complementary.
29. Prove that there exists a self-complementary graph of order  $n$  for every positive integer  $n$  with  $n \equiv 0 \pmod{4}$  or  $n \equiv 1 \pmod{4}$ .
30. (a) Give an example of a graph  $G$  of order 6 and size 7 such that  $G$  is isomorphic to a subgraph  $H$  of  $\overline{G}$ .  
(b) Give an example of a graph  $G$  of order 7 and size 10 such that  $G$  is isomorphic to a subgraph  $H$  of  $\overline{G}$ .
31. Prove for every integer  $n \geq 3$  that there exists a graph  $G$  of order  $n$  and size  $\lfloor \binom{n}{2}/2 \rfloor$  that is isomorphic to a graph  $H \subseteq \overline{G}$ .

32. Let  $P$  be the Petersen graph. Show that  $\overline{P}$  contains a subgraph  $H$  such that  $H \cong P$ .
33. For  $i = 1, 2$ , let  $u_i$  be a vertex in a graph  $G_i$  of order  $n_i$  and size  $m_i$ .
- Determine the degree of  $u_1$  in  $G_1 + G_2$ .
  - Determine the degree of  $u_1$  in  $G_1 \vee G_2$ .
  - Determine the degree of  $(u_1, u_2)$  in  $G_1 \square G_2$ .
34. Determine the order and size of each of the graphs  $P_3 \vee 2P_3$ ,  $P_3 \square 2P_3$  and  $Q_1 + Q_2 + Q_3$ .

### Section 1.7. Degree Sequences

35. Find a sequence of 2-switches that transforms the graph  $G$  of Figure 1.34 into the graph  $H$ .




Figure 1.34: The graphs  $G$  and  $H$  in Exercise 35

36. For two pairs  $G_1, H_1$  and  $G_2, H_2$  of graphs shown in Figure 1.35, determine the minimum number of 2-switches required to transform
- $G_1$  into  $H_1$  and
  - $G_2$  into  $H_2$ .




Figure 1.35: The graphs in Exercise 36

37. Let  $s : 2, 2, 2, 2, 2, 2, 2, 2$  and let  $\mathcal{G}_s$  be the set of all graphs with degree sequence  $s$ . Let  $G$  be a graph with  $V(G) = \mathcal{G}_s$  where two vertices  $F$  and  $H$  in  $G$  are adjacent if  $F$  can be transformed into  $H$  by a single 2-switch. To which familiar graph is  $G$  isomorphic?

38. Give an example of a graphical sequence  $s$  (where  $\mathcal{G}_s$  is the set of all graphs with degree sequence  $s$ ) such that (1) the graph  $G$  has  $V(G) = \mathcal{G}_s$ , (2) two vertices  $F$  and  $H$  of  $G$  are adjacent if  $F$  can be transformed into  $H$  by a single 2-switch and (3)  $G$  contains a triangle.
39. Let  $s : d_1, d_2, \dots, d_n$  be a graphical sequence with  $\Delta = d_1 \geq d_2 \geq \dots \geq d_n$ . Show, for each integer  $k$  with  $1 \leq k \leq n$ , that there exists a graph  $G$  with  $V(G) = \{v_1, v_2, \dots, v_n\}$  where  $\deg v_i = d_i$  for  $1 \leq i \leq n$  having the property that  $v_k$  is adjacent to either (1) the vertices of  $\{v_1, v_2, \dots, v_{d_k}\}$  if  $k > d_k$  or (2) the vertices of  $\{v_1, v_2, \dots, v_{d_k+1}\} - \{v_k\}$  if  $1 \leq k \leq d_k$ .
40. Let  $G$  and  $H$  be two graphs that are neither empty nor complete. The graph  $H$  is said to be obtained from  $G$  by an **edge rotation** if  $G$  contains three vertices  $u, v$ , and  $w$  where  $uv \in E(G)$  and  $uw \notin E(G)$  and  $H \cong G - uv + uw$ .
- Show that the graph  $G_2$  of Figure 1.36 is obtained from  $G_1$  by an edge rotation.
  - Show that  $G_3$  of Figure 1.36 cannot be obtained from  $G_1$  by an edge rotation.
  - Show that for every two nonempty, noncomplete graphs  $G$  and  $H$  of the same order and same size, there exists a sequence  $G = G_0, G_1, \dots, G_k = H$  of graphs such that  $G_{i+1}$  is obtained from  $G_i$  by an edge rotation for  $i = 0, 1, \dots, k - 1$ .




Figure 1.36: The graphs in Exercise 40

41. Determine whether the following sequences are graphical. If so, construct a graph with the appropriate degree sequence.
- 4, 4, 3, 2, 1
  - 3, 3, 2, 2, 2, 2, 1, 1
  - 7, 7, 6, 5, 4, 4, 3, 2
  - 7, 6, 6, 5, 4, 3, 2, 1
  - 7, 4, 3, 3, 2, 2, 2, 1, 1, 1.
42. Prove that a sequence  $d_1, d_2, \dots, d_n$  is graphical if and only if  $n - d_1 - 1, n - d_2 - 1, \dots, n - d_n - 1$  is graphical.

43. Prove that for every integer  $x$  with  $0 \leq x \leq 5$ , the sequence  $x, 1, 2, 3, 5, 5$  is not graphical.
44. For which integers  $x$  ( $0 \leq x \leq 7$ ), if any, is the sequence  $7, 6, 5, 4, 3, 2, 1, x$  graphical?
45. Use Theorem 1.13 to determine whether the sequence  $s : 6, 6, 5, 4, 3, 2, 2$  is graphical.
46. We have seen that there is only one graphical sequence  $d_1, d_2, d_3, d_4, d_5$  with  $4 = d_1 \geq d_2 \geq d_3 \geq d_4 \geq d_5 = 1$  such that at least one term is 3 and at least one term is 2. How many graphical sequences  $d_1, d_2, d_3, d_4, d_5$  are there with  $4 = d_1 \geq d_2 \geq d_3 \geq d_4 \geq d_5 = 2$  such that at least one term is 3?
47. Show that for every finite set  $S$  of positive integers, there exists a positive integer  $k$  such that the sequence obtained by listing each element of  $S$  a total of  $k$  times is graphical. Find the minimum such  $k$  for  $S = \{2, 6, 7\}$ .
48. According to Theorem 1.15, for each integer  $n \geq 2$ , there exist exactly two distinct graphical sequences of length  $n$  having exactly two equal terms. What terms are equal for these two sequences?
49. Two finite sequences  $s_1$  and  $s_2$  of nonnegative integers are called **bigraphical** if there exists a bipartite graph  $G$  with partite sets  $V_1$  and  $V_2$  such that  $s_i$  lists the degrees of the vertices of  $G$  in  $V_i$  for  $i = 1, 2$ . Prove that the sequences  $s_1 : a_1, a_2, \dots, a_r$  and  $s_2 : b_1, b_2, \dots, b_t$  of nonnegative integers with  $r \geq 2$ ,  $a_1 \geq a_2 \geq \dots \geq a_r$ ,  $b_1 \geq b_2 \geq \dots \geq b_t$ ,  $0 < a_1 \leq t$  and  $0 < b_1 \leq r$  are bigraphical if and only if the sequences  $s'_1 : a_2, a_3, \dots, a_r$  and  $s'_2 : b_1 - 1, b_2 - 1, \dots, b_{a_1} - 1, b_{a_1+1}, \dots, b_t$  are bigraphical.
50. The graphs  $G$  and  $H$  of order 10 have vertex sets  $V(G) = \{u_1, u_2, \dots, u_{10}\}$  and  $V(H) = \{v_1, v_2, \dots, v_{10}\}$  and edge sets  $E(G) = \{u_i u_j : i + j \geq 11\}$  and  $E(H) = \{v_i v_j : i + j \geq 12\}$ . How are  $G$  and  $H$  related?
51. (a) Let  $n$  be a given positive integer and let  $r$  and  $s$  be nonnegative integers such that  $r + s = n$  and  $s$  is even. Give an example of a graph containing  $r$  even vertices and  $s$  odd vertices.  
 (b) Determine the minimum size of a graph  $G$  containing  $r$  even vertices and  $s$  odd vertices and satisfying the properties in (a).  
 (c) Determine the maximum size of a graph  $G$  containing  $r$  even vertices and  $s$  odd vertices and satisfying the properties in (a).
52. (a) Let  $G$  be a graph of order  $n \geq 4$ . Prove that if  $\deg v \geq \frac{2n+1}{3}$  for every vertex  $v$  of  $G$ , then every edge of  $G$  belongs to a complete subgraph of order 4.

- (b) Show that the result in (a) is best possible in general by showing that  $\frac{2n+1}{3}$  cannot be replaced by  $\frac{2n}{3}$ .

### Section 1.8. Multigraphs

53. We saw that the irregular multigraph  $G_2$  in Figure 1.30 has degree sequence 4, 3, 2, 1. Give an example of an irregular multigraph (if such a multigraph exists) having degree sequence
- (a) 5, 4, 3, 2, 1
  - (b) 6, 5, 4, 3, 2, 1
  - (c) 7, 6, 5, 4, 3, 2, 1.
54. Prove for every connected graph  $G$  of order  $n = 3$  or  $n = 4$  and size  $m$  that it is possible to label the edges of  $G$  by  $e_1, e_2, \dots, e_m$  and replace  $e_i$  by  $i$  parallel edges for each  $i$  ( $1 \leq i \leq m$ ) such that the degrees of the vertices of the resulting multigraph  $H$  are distinct.
55. Determine which of the following sequences are the degree sequences of a multigraph.
- (a)  $s_1 : 3, 2, 1$       (b)  $s_2 : 5, 2, 1$
  - (c)  $s_3 : 6, 4, 2$       (d)  $s_4 : 3, 2, 2$
  - (e)  $s_5 : 4, 4, 2, 2$     (f)  $s_6 : 5, 3, 2, 1$
  - (g)  $s_7 : 4, 4, 4, 4$     (h)  $s_8 : 7, 5, 3, 1$ .
56. Prove that a sequence  $s : d_1, d_2, \dots, d_n$  ( $n \geq 1$ ) of nonnegative integers with  $d_1 \geq d_2 \geq \dots \geq d_n$  is the degree sequence of a multigraph if and only if  $\sum_{i=1}^n d_i$  is even and  $d_1 \leq \frac{1}{2} \sum_{i=1}^n d_i$ .
57. Let  $G$  be a connected graph of order  $n$  where the vertices of  $G$  are labeled as  $v_1, v_2, \dots, v_n$  in some way. A multigraph  $H$  of size  $m$  with  $V(H) = V(G)$  is obtained by replacing each edge  $v_i v_j$  of  $G$  by  $\min\{i, j\}$  parallel edges.
- (a) Find  $m$  if  $G = K_5$ .
  - (b) Find sharp upper and lower bounds for  $m$  if  $G = C_5$ .
  - (c) Find the minimum value of  $m$  if  $G$  is bipartite.





# Bibliography

- [1] B. M. Ábrego, S. Fernández-Merchant and G. Salazar, The rectilinear crossing number of  $K_n$ : Closing in (or are we?) *Thirty essays on geometric graph theory* 5-18, Springer, New York, 2013. [286]
- [2] Y. Alavi, A. J. Boals, G. Chartrand, P. Erdős and O. R. Oellermann, The ascending subgraph decomposition problem. *Congr. Numer.* **58** (1987) 7-14. [345]
- [3] N. Alon, Choice numbers of graphs; a probabilistic approach, *Comb. Prop. Comput.* **1** (1992) 107-114. [409]
- [4] B. Alspach, Research problems, Problem 3. *Discrete Math.* **36** (1981) 333. [349]
- [5] B. Alspach, The wonderful Walecki construction. *Bull. Inst. Combin. Appl.* **52** (2008) 7- 20. [341]
- [6] B. Alspach and H. Gavlas, Cycle decompositions of  $K_n$  and  $K_n - I$ . *J. Combin. Theory Ser. B.* **81** (2001) 77-99. [348, 350]
- [7] V. I. Arnautov, Estimation of the exterior stability number of a graph by means of the minimal degree of the vertices. (Russian) *Prikl. Mat. i Programirovanie Vyp.* **11** (1974) 3-8. [325, 550]
- [8] I. Anderson, Perfect matchings of a graph. *J. Combin. Theory Ser. B* **10** (1971) 183-186. [311]
- [9] K. Appel and W. Haken, Every planar map is four colorable. *Bull. Amer. Math. Soc.* **82** (1976) 711-712. [427]
- [10] D. Archdeacon and P. Huneke, A Kuratowski theorem for nonorientable surfaces. *J. Combin. Theory Ser. B* **46** (1989) 173-231. [300]
- [11] A. T. Balaban, Trivalent graphs of girth nine and eleven, and relationships among cages. *Rev. Roumaine Math. Pures Appl.* **18** (1973) 1033-1043. [516]

- [12] E. Bannai and T. Ito, On finite Moore graphs. *J. Fac. Sci. Univ. Tokyo, Sect. IA* **20** (1973) 191-208. [518]
- [13] J. Battle, F. Harary and V. Kodama, Every planar graph with nine points has a nonplanar complement. *Bull. Amer. Math. Soc.* **68** (1962) 569-571. [293]
- [14] D. Bauer, H. J. Broersma and H. J. Veldman, Not every 2-tough graph is Hamiltonian. *Discrete Appl. Math.* **99** (2000) 317-321. [138]
- [15] M. Behzad, *Graphs and Their Chromatic Numbers*. Ph.D. Thesis, Michigan State University (1965). [496]
- [16] M. Behzad, G. Chartrand and J. K. Cooper, Jr., The colour numbers of complete graphs. *J. London Math. Soc.* **42** (1967) 226-228. [497]
- [17] L. W. Beineke, Characterizations of derived graphs. *J. Combin. Theory* **9** (1970) 129-135. [149]
- [18] L. W. Beineke and F. Harary, The genus of the  $n$ -cube. *Canad. J. Math.* **17** (1965) 494-496. [295]
- [19] L. W. Beineke and R. D. Ringeisen, On the crossing numbers of products of cycles and graphs of order four. *J. Graph Theory* **4** (1980) 145-155. [282, 283]
- [20] C. T. Benson, Minimal regular graphs of girths eight and twelve. *Canad. J. Math.* **18** (1966) 1091-1094. [516]
- [21] C. Berge, Two theorems in graph theory. *Proc. Nat. Acad. Sci. U.S.A.* **43** (1957) 842-844. [307]
- [22] C. Berge, *Théorie des Graphes et Ses Applications*. Dunod, Paris (1958). [99, 323]
- [23] C. Berge, Sur le couplage maximum d'un graphe. *C. R. Acad. Sci. Paris* **247** (1958) 258-259. [317]
- [24] C. Berge, Some classes of perfect graphs. *Six Papers on Graph Theory*. Indian Statistical Institute, Calcutta (1963) 1-21. [396]
- [25] C. Berge, *Graphs and Hypergraphs*. North-Holland, Amsterdam-London (1973). [20, 326]
- [26] H. Bielak and M. M. Syslo, Peripheral vertices in graphs. *Studia Sci. Math. Hungar.* **18** (1983) 269-275. [49]
- [27] G. D. Birkhoff, A determinant formula for the number of ways of coloring a map. *Ann. of Math.* **14** (1912) 42-46. [438]

- [28] D. Blanuša, Problem cetiriju boja. *Glasnik Mat. Fiz. Astr.* **1** (1946) 31-42. [474]
- [29] A. Blass and F. Harary, Properties of almost all graphs and complexes. *J. Graph Theory.* **3** (1979) 225-240. [556, 558]
- [30] J. Blažek and M. Koman, A minimal problem concerning complete plane graphs. *Theory of Graphs and its Applications*. Publ. House. Czechoslovak Acad. Sci. Prague (1964) 113-117. [277]
- [31] B. Bollobás and E. J. Cockayne, The irredundance number and maximum degree of a graph. *Discrete Math.* **49** (1984) 197-199. [325]
- [32] B. Bollobás and A. J. Harris, List-colourings of graphs. *Graphs Combin.* **1** (1985) 115-127. [492]
- [33] J. A. Bondy, Pancyclic graphs. I. *J. Combin. Theory Ser. B* **11** (1971) 80-84. [145]
- [34] J. A. Bondy and V. Chvátal, A method in graph theory. *Discrete Math.* **15** (1976) 111-136. [130, 131]
- [35] O. Borůvka, O jistém Problému minimálním. *Práce Mor. Přírodověd. Spol. v Brně (Acta Societ. Scient. Natur. Moraviae)* **3** (1926) 37-58. [81]
- [36] R. C. Brigham and R. D. Dutton, A compilation of relations between graph invariants. *Networks* **15** (1985) 73-107. [381]
- [37] A. Brodsky, S. Durocher and E. Gethner, Toward the rectilinear crossing number of  $K_n$ : new drawings, upper bounds, and asymptotics. *Discrete Math.* **262** (2003) 59-77. [286]
- [38] R. L. Brooks, On coloring the nodes of a network. *Proc. Cambridge Philos. Soc.* **37** (1941) 194-197. [378]
- [39] D. Bryant, D. Horsley and W. Pettersson, Cycle decompositions V: Complete graphs into cycles of arbitrary lengths. *Proc. London Math. Soc.* To appear. [349, 350, 491]
- [40] F. Buckley, Z. Miller and P. J. Slater, On graphs containing a given graph as center. *J. Graph Theory* **5** (1981) 427-434. [48]
- [41] P. Camion, Chemins et circuits hamiltoniens des graphes complets. *C. R. Acad. Sci. Paris* **249** (1959) 2151-2152. [181]
- [42] P. A. Catlin, Hajós' graph-coloring conjecture: variations and counterexamples. *J. Combin. Theory Ser. B* **26** (1979) 268-274. [435]
- [43] A. Cayley, On the theory of analytical forms called trees. *Philos. Mag.* **13** (1857) 19-30. [64]

- [44] A. Cayley, A theorem on trees. *Quart. J. Math.* **23** (1889) 376-378. [73]
- [45] G. Chartrand, F. Fujie and P. Zhang On an extension of an observation of Hamilton. *J. Combin. Math. Combin. Comput.* To appear. [142]
- [46] G. Chartrand and F. Harary, Graphs with prescribed connectivities, *Theory of Graphs*. Academic Press, New York (1968) 61-63. [96]
- [47] G. Chartrand, A. M. Hobbs, H. A. Jung, S. F. Kapoor and C. St. J. A. Nash-Williams, The square of a block is Hamiltonian connected. *J. Combin. Theory Ser. B* **16** (1974) 290-292. [147]
- [48] G. Chartrand, H. Jordon and P. Zhang, A cycle decomposition conjecture for Eulerian graphs. *Australas. J. Combin.* **58** (2014) 48-59. [351]
- [49] G. Chartrand and J. Mitchem, Graphical theorems of the Nordhaus-Gaddum class. *Recent Trends in Graph Theory*. Springer, Berlin (1971) 55-61. [380]
- [50] G. Chartrand, A. D. Polimeni and M. J. Stewart, The existence of 1-factors in line graphs, squares, and total graphs. *Nedrl. Akad. Wetensch. Proc. Ser. A* 76 *Indag. Math.* **35** (1973) 228-232. [120]
- [51] G. Chartrand and C. E. Wall, On the Hamiltonian index of a graph. *Studia Sci. Math. Hungar.* **8** (1973) 43-48. [151]
- [52] A. G. Chetwynd and A. J. W. Hilton, Star multigraphs with three vertices of maximum degree. *Math. Proc. Cambridge Philos. Soc.* **100** (1986) 303-317. [339, 466]
- [53] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem. *Ann. of Math.* **164** (2006) 51-229. [405]
- [54] V. Chvátal, On Hamilton's ideals. *J. Combin. Theory Ser. B* **12** (1972) 163-168. [131]
- [55] V. Chvátal, Tough graphs and Hamiltonian circuits. *Discrete Math.* **5** (1973) 215-228. [136, 138]
- [56] V. Chvátal, A combinatorial theorem in plane geometry. *J. Combin. Theory Ser. B* **18** (1975) 39-41. [285]
- [57] V. Chvátal, Tree-complete graph Ramsey numbers. *J. Graph Theory* **1** (1977) 93. [534]
- [58] V. Chvátal and P. Erdős, A note on hamiltonian circuits. *Discrete Math.* **2** (1972) 111-113. [133]
- [59] A. Clebsch, Über die Flächen vierter Ordnung, welche eine Doppelcurve zweiten Grades besitzen. *J. für Math.* **69** (1868) 142-184. [532]

- [60] E. J. Cockayne and S. T. Hedetniemi, Towards a theory of domination in graphs. *Networks* **7** (1977) 247-261. [323]
- [61] S. Cook, The complexity of theorem proving procedures. *Proceedings of the Third Annual ACM Symposium on Theory of Computing* (1971) 151-158. [140]
- [62] R. Courant and H. Robbins, *What is Mathematics?* Oxford University Press, London (1941). [446]
- [63] B. Csaba, D. Kühn, A. Lo, D. Osthus and A. Treglown, Proof of the 1-factorization and Hamiltonian Decomposition Conjectures. *Memoirs of the AMS*. To appear. [339, 343]
- [64] R. M. Damerell, On Moore graphs. *Proc. Cambridge Philos. Soc.* **74** (1973) 227-236. [518]
- [65] C. F. de Jaenisch, *Applications de l'Analyse Mathématique au Jeu des Echecs*. Petrograd (1862). [322]
- [66] B. Descartes, A three colour problem. *Eureka* **9** (1947) 21. [394]
- [67] B. Descartes, Network-colouring. *Math. Gazette* **32** (1948) 67-69. [474]
- [68] B. Descartes, The expanding unicurse. *Proof Techniques in Graph Theory* (F. Harary, ed.). Academic Press, New York (1969) 25. [199]
- [69] E. A. Dinic, An algorithm for the solution of the problem of maximal flow in a network with power estimation. *Soviet Math. Dokl.* **11** (1970) 1277-1280. [201]
- [70] G. A. Dirac, Some theorems on abstract graphs. *Proc. London Math. Soc.* **2** (1952) 69-81. [129]
- [71] G. A. Dirac, A property of 4-chromatic graphs and some remarks on critical graphs. *J. London Math. Soc.* **27** (1952) 85-92. [434]
- [72] G. A. Dirac, In abstrakten Graphen vorhandene vollständige 4-Graphen und ihre Unterteilungen. *Math. Nachr.* **22** (1960) 61-85. [107]
- [73] G. A. Dirac, On rigid circuit graphs. *Abh. Math. Sem. Univ. Hamburg* **25** (1961) 71-76. [400]
- [74] J. Edmonds and R. A. Karp, Theoretical improvements in algorithmic efficiency for network flow problems. *Journal of ACM.* **19** (1972) 248-264. [201, 202]
- [75] E. Egerváry, On combinatorial properties of matrices (Hungarian). *Mat. Lapok* **38** (1931) 16-28. [320]

- [76] R. B. Eggleton and R. K. Guy, The crossing number of the  $n$ -cube. *Notices Amer. Math. Soc.* **17** (1970) 757. [281]
- [77] P. Elias, A. Feinstein and C. E. Shannon A note on the maximum flow through a network. *IRE Trans. on Inform. Theory* **IT 2** (1956) 117-119. [199]
- [78] P. Erdős, Some remarks on the theory of graphs. *Bull. Amer. Math. Soc.* **53** (1947) 292-294. [530]
- [79] P. Erdős, Graph theory and probability II. *Canad. J. Math* **13** (1961) 346-352. [396, 551]
- [80] P. Erdős, Extremal problems in graph theory. *A Seminar on Graph Theory*. Holt, Rinehart and Winston, New York (1967) 54-59. [505]
- [81] P. Erdős and T. Gallai, Graphs with prescribed degrees of vertices (Hungarian). *Mat. Lapok* **11** (1960) 264-274. [23]
- [82] P. Erdős, A. L. Rubin and H. Taylor, Choosability in graphs. *Congr. Numer.* **126** (1980) 125-157. [406, 428]
- [83] P. Erdős and H. Sachs, Reguläre Graphen gegebener Taillenweite mit minimaler Knotenzahl. (German) *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* **12** (1963) 251-257. [511]
- [84] P. Erdős and G. Szekeres, A combinatorial problem in geometry. *Compositio Math.* **2** (1935) 463-470. [526]
- [85] A. Errera, Une demonstration du theoreme de Petersen. *Mathesis* **36** (1922) 56-61. [315]
- [86] L. Euler, Solutio problematis ad geometriam situs pertinentis. *Comment. Academiae Sci. I. Petropolitanae* **8** (1736) 128-140. [6, 115, 118]
- [87] L. Euler, Elementa doctrinae solidorum. *Novi Comm. Acad. Sci. Imp. Petropol.* **4** (1752-1753) 109-140. [240]
- [88] L. Euler, Demonstratio nonnullarum insignium proprietatum quibus solida hedris planis inclusa sunt praedita. *Novi Comm. Acad. Sci. Imp. Petropol.* **4** (1752-1753) 140-160. [240]
- [89] I. Fáry, On straight line representation of planar graphs. *Acta Univ. Szeged Sect. Sci. Math.* **11** (1948) 229-233. [283]
- [90] H. J. Finck, On the chromatic numbers of a graph and its complement. *Theory of Graphs*. Academic Press, New York (1968) 99-113. [381]
- [91] S. Fisk, A short proof of Chvátal's watchman theorem. *J. Combin. Theory Ser. B* **24** (1978) 374. [285]

- [92] H. Fleischner, The square of every two-connected graph is Hamiltonian. *J. Combin. Theory Ser. B* **16** (1974) 29-34. [147]
- [93] L. R. Ford, Jr. and D. R. Fulkerson, Maximal flow through a network. *Canad. J. Math.* **8** (1956) 399-404. [198, 199]
- [94] L. R. Ford, Jr. and D. R. Fulkerson, A simple algorithm for finding maximal network flows and an application to the Hitchcock problem. *Canad. J. Math.* **9** (1957) 210-218. [200]
- [95] P. Franklin, The four color problem. *Amer. J. Math.* **44** (1922) 225-236. [252]
- [96] O. Frink and P. A. Smith, Irreducible non-planar graphs. *Bull. Amer. Math. Soc.* **36** (1930) 214. [254]
- [97] R. Frucht, Herstellung von Graphen mit vorgegebener abstrakter Gruppe. *Compositio Math.* **6** (1938) 239-250. [227]
- [98] T. Gallai, Maximum-minimum Sätze über Graphen. *Acta Math. Acad. Sci. Hungar.* **9** (1958) 395-434. [396]
- [99] T. Gallai, Über extreme Punkt- und Kantenmengen. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* **2** (1959) 133-138. [319]
- [100] T. Gallai, On directed paths and circuits. *Theory of Graphs: Proceedings of the Colloquium held at Tihany, Hungary, 1969* (P. Erdős and G. Katona, eds). Academic Press, New York (1969) 115-118. [383]
- [101] F. Galvin, The list chromatic index of a bipartite multigraph. *J. Combin. Theory Ser. B* **63** (1995) 153-158. [493]
- [102] D. P. Geller, Problem 5713. *Amer. Math. Monthly* **77** (1970) 85. [379]
- [103] A. Georgakopoulos, A short proof of Fleischner's theorem. *Discrete Math.* **309** (2009) 6632-6634. [147]
- [104] A. Ghouila-Houri, Une condition suffisante d'existence d'un circuit Hamiltonien. *C. R. Acad. Sci. Paris* **251** (1960) 495-497. [169]
- [105] E. N. Gilbert, Random graphs. *Ann. Math. Stat.* **30** (1959) 1141-1144. [556]
- [106] S. W. Golomb, How to number a graph. *Graph Theory and Computing*. Academic Press, New York (1972) 23-37. [351, 356]
- [107] R. L. Graham and P. Hell, On the history of the minimum spanning tree problem. *Annals of the History of Computing* **7** (1985) 43-57. [81]
- [108] R. E. Greenwood and A. M. Gleason, Combinatorial relations and chromatic graphs. *Canad. J. Math.* **7** (1955) 1-7. [532]



- [109] E. J. Grinberg, Plane homogeneous graphs of degree three without hamiltonian circuits. *Latvian Math. Yearbook* **4** (1968) 51-58. [265]
- [110] R. P. Gupta, The chromatic index and the degree of a graph. *Notices Amer. Math. Soc.* **13** (1966) 719. [455, 459]
- [111] R. K. Guy, A combinatorial problem. *Bull. Malayan Math. Soc.* **7** (1960) 68-72. [277]
- [112] R. K. Guy, Crossing numbers of graphs. *Graph Theory and Applications*. Springer-Verlag, New York (1972) 111-124. [276, 278, 286]
- [113] H. Hadwiger, Über eine Klassifikation der Streckenkomplexe. *Vierteljahrsschr. Naturforsch. Ges. Zürich* **88** (1943) 133-142. [437]
- [114] R. Haggkvist and A. G. Chetwynd, Some upper bounds on the total and list chromatic numbers of multigraphs. *J. Graph Theory* **16** (1992) 503-516. [492]
- [115] A. Hajnal and J. Surányi, Über die Auflösung von Graphen in vollständige Teilgraphen. *Ann. Univ. Sci. Budapest Eötvös. Sect. Math.* **1** (1958) 113-121. [400]
- [116] G. Hajós, Über eine Konstruktion nicht  $n$ -färbbarer Graphen. *Wiss. Z. Martin-Luther Univ. Halle-Wittenberg. Math.-Nat. Reihe.* **10** (1961) 116-117. [435]
- [117] S. L. Hakimi, On the realizability of a set of integers as degrees of the vertices of a graph. *SIAM J. Appl. Math.* **10** (1962) 496-506. [21]
- [118] P. Hall, On representation of subsets. *J. London Math. Soc.* **10** (1935) 26-30. [308]
- [119] F. Harary, The maximum connectivity of a graph. *Proc. Nat. Acad. Sci. U.S.A.* **48** (1962) 1142-1146. [100]
- [120] F. Harary, P. C. Kainen and A. J. Schwenk, Toroidal graphs with arbitrarily high crossing numbers. *Nanta Math.* **6** (1973) 58-67. [281]
- [121] F. Harary and L. Moser, The theory of round robin tournaments. *Amer. Math. Monthly* **73** (1966) 231-246. [178, 183]
- [122] F. Harary and C. St. J. A. Nash-Williams, On eulerian and hamiltonian graphs and line graphs. *Canad. Math. Bull.* **8** (1965) 701-709. [149]
- [123] F. Harary and R. Z. Norman, The dissimilarity characteristic of Husimi trees. *Ann. of Math.* **58** (1953) 134-141. [61]
- [124] V. Havel, A remark on the existence of finite graphs (Czech.) *Časopis Pěst. Mat.* **80** (1955) 477-480. [21]

- [125] P. J. Heawood, Map colour theorems. *Quart. J. Math.* **24** (1890) 332-338.  
[422, 426, 444, 445]
- [126] P. J. Heawood, On the four-colour map theorem. *Quart. J. Pure Appl. Math.* **29** (1898) 270-285. [427]
- [127] L. Heffter, Über das Problem der Nachbargebiete. *Math. Ann.* **38** (1891) 477-508. [446]
- [128] C. Hierholzer, Über die Möglichkeit, einen Linienzug ohne Wiederholung und ohne Unterbrechnung zu umfahren. *Math. Ann.* **6** (1873) 30-32. [120]
- [129] A. J. Hoffman and R. R. Singleton, On Moore graphs with diameter 2 and 3. *IBM J. Res. Develop.* **4** (1960) 497-504. [516, 518]
- [130] I. Holyer, The NP-completeness of edge colouring. *SIAM J. Comput.* **4** (1981) 718-720. [460]
- [131] J. Hopcroft and R. E. Tarjan, Efficient planarity testing. *J. Assoc. Comput. Mach.* **21** (1974) 549-568. [257]
- [132] R. Isaacs, Infinite families of nontrivial trivalent graphs which are not Tait colorable. *Amer. Math. Monthly* **82** (1975) 221-239. [474]
- [133] H. Izbicki, Reguläre Graphen beliebigen Grades mit vorgegebenen Eigenschaften. *Monatsh. Math.* **64** (1960) 15-21. [228]
- [134] B. Jackson, Hamilton cycles in regular 2-connected graphs. *J. Combin. Theory Ser. B* **29** (1980) 27-46. [132]
- [135] F. Jaeger, A survey of the cycle double cover conjecture. *Cycles in Graphs*. North-Holland, Amsterdam (1985) 1-12. [491]
- [136] V. Jarník, O jistém problému min imálním. (Czech) *Acta Societ. Scient. Natur. Moraviae* **6** (1930) 57-63. [83]
- [137] T. Jensen and G. F. Royle, Small graphs with chromatic number 5: a computer search. *J. Graph Theory* **19** (1995) 107-116. [395]
- [138] J. J. Karaganis, On the cube of a graph. *Canad. Math. Bull.* **11** (1968) 295-296. [146]
- [139] R. M. Karp, Reducibility among combinatorial problems, in *Complexity Computer Computations* (R. E. Miller and J. W. Thatcher, eds.). Plenum, New York (1972) 85-103. [140]
- [140] F. Kárteszi, Piani finiti ciclici come risoluzioni di un certo problema di minimo. *Boll. Un. Mat. Ital.* **15** (1960) 522-528. [510]
- [141] J. B. Kelly and L. M. Kelly, Paths and circuits in critical graphs. *Amer. J. Math.* **76** (1954) 786-792. [394]

- [142] G. Kirchhoff, Über die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Verteilung galvanischer Ströme geführt wird. *Ann. Phys. Chem.* **72** (1847) 497-508. [64, 77]
- [143] T. P. Kirkman, On a problem in combinatorics. *Cambridge and Dublin Math. J.* **2** (1847) 191-204. [347]
- [144] D. J. Kleitman, The crossing number of  $K_{5,n}$ . *J. Combin. Theory* **9** (1970) 315-323. [279]
- [145] W. L. Kocay, On Stockmeyer's nonreconstructible tournaments. *J. Graph Theory* **9** (1985) 473-476. [230]
- [146] D. König, Über Graphen und ihre Anwendung auf Determinantentheorie und Mengenlehre. *Math. Ann.* **77** (1916) 453-465. [336, 461]
- [147] D. König, Graphen und Matrizen. *Math. Lapok.* **38** (1931) 116-119. [416]
- [148] D. König, *Theorie der endlichen und unendlichen Graphen*. Akademische Verlagsgesellschaft, Leipzig (1936). [103, 226]
- [149] A. D. Korshunov, A solution of a problem of P. Erdős and A. Rényi about Hamiltonian cycles in non-oriented graphs, Metody Diskr. Anal. Teoriy, Upr. Syst., Sb. Trudov Novosibirsk. **31** (1977) 17-56 (in Russian). [560]
- [150] J. B. Kruskal, On the shortest spanning tree of a graph and the traveling salesmen problem. *Proc. Amer. Math. Soc.* **7** (1956) 48-50. [82]
- [151] K. Kuratowski, Sur le problème des courbes gauches en topologie. *Fund. Math.* **15** (1930) 271-283. [254]
- [152] H. G. Landau, On dominance relations and structure of animal societies. III. The condition for a score structure. *Bull. Math. Biophys.* **15** (1953) 143-148. [176, 179]
- [153] A.-M. Legendre, *Éléments de Géométrie*. F. Didot, Paris (1794). [240]
- [154] L. Lesniak, Results on the edge-connectivity of graphs. *Discrete Math.* **8** (1974) 351-354. [102]
- [155] L. Lesniak, Eccentric sequences in graphs. *Period. Math. Hungar.* **6** (1975) 287-293. [75]
- [156] S. A. J. Lhuillier, Mémoire sur la polyédrométrie, contenant une démonstration directe du théorème d'Euler sur les polyèdres, et un examen de diverses exceptions auxquelles ce théorème est assujéti. *Ann. Math. Pures et Appliquées* **3** (1812-1813) 169-189. [290]
- [157] D. R. Lick, Characterizations of  $n$ -connected and  $n$ -line-connected graphs. *J. Combin. Theory Ser. B* **14** (1973) 122-124. [107]

- [158] L. Lovász, On chromatic number of finite set-systems. *Acta. Math. Acad. Sci. Hungar.* **79** (1967) 59-67. [396, 405, 551]
- [159] L. Lovász, A characterization of perfect graphs. *J. Combin. Theory Ser. B* **13** (1972) 95-98. [402]
- [160] L. Lovász, Normal hypergraphs and the perfect graph conjecture. *Discrete Math.* **2** (1972) 253-267. [402]
- [161] W. Mantel, Problem 28. *Wiskundige Opgaven* **10** (1907) 60-61. [15, 503]
- [162] B. Manvel, On Reconstruction of Graphs. Doctoral thesis, University of Michigan (1970). [232]
- [163] M. M. Matthews and D. P. Sumner, Hamiltonian results in  $K_{1,3}$ -free graphs. *J. Graph Theory* **8** (1984) 139-146. [137, 157]
- [164] D. W. Matula, The cohesive strength of graphs. *The Many Facets of Graph Theory*. Springer, Berlin (1969) 215-221. [373]
- [165] W. F. McGee, A minimal cubic graph of girth seven. *Canad. Math. Bull.* **3** (1960) 149-152. [514]
- [166] B. D. McKay, Computer reconstruction of small graphs. *J. Graph Theory* **1** (1977) 281-283. [229]
- [167] B. D. McKay, W. Myrvold and J. Nadon, Fast backtracking principles applied to find new cages. *Proceedings of the Ninth Annual ACM-SIAM Symposium on Discrete Algorithms*. ACM, New York (1998) 188-191. [516]
- [168] T. A. McKee, Recharacterizing Eulerian: intimations of new duality. *Discrete Math.* **51** (1984) 237-242. [122]
- [169] K. Menger, Zur allgemeinen Kurventheorie. *Fund. Math.* **10** (1927) 95-115. [102]
- [170] M. Meyniel, Une condition suffisante d'existence d'un circuit Hamiltonien dans un graphe orienté. *J. Combin. Theory Ser. B* **14** (1973) 137-147. [168]
- [171] M. Mirzakhani, A small non-4-choosable planar graph. *Bull. Inst. Combin. Appl.* **17** (1996) 15-18. [430]
- [172] M. Molloy and B. Reed, A bound on the total chromatic number. *Combinatorica* **18** (1998) 241-280. [498]
- [173] J. W. Moon, On subtournaments of a tournament. *Canad. Math. Bull.* **9** (1966) 297-301. [183]
- [174] J. Mycielski, Sur le coloriage des graphes. *Colloq. Math.* **3** (1955) 161-162. [394]

- [175] C. St. J. A. Nash-Williams, Edge-disjoint Hamiltonian circuits in graphs with the vertices of large valency, in *Studies in Pure Mathematics*. Academic Press (1971) 157-183. [343]
- [176] C. St. J. A. Nash-Williams, Hamiltonian arcs and circuits, in *Recent Trends in Graph Theory*. Springer Verlag (1971) 197-210. [343]
- [177] A. Nijenhuis, Note on the unique determination of graphs by proper subgraphs. *Notices Amer. Math. Soc.* **24** (1977) A-290. [229]
- [178] E. A. Nordhaus and J. W. Gaddum, On complementary graphs. *Amer. Math. Monthly* **63** (1956) 175-177. [380]
- [179] O. Ore, Note on Hamilton circuits. *Amer. Math. Monthly* **67** (1960) 55. [129]
- [180] O. Ore, *Theory of Graphs*. Amer. Math. Soc. Colloq. Pub., Providence, RI (1962). [323, 324]
- [181] O. Ore, Hamilton connected graphs. *J. Math. Pures Appl.* **42** (1963) 21-27. [140]
- [182] S. Pan and R. B. Richter, The crossing number of  $K_{11}$  is 100. *J. Graph Theory* **56** (2007) 128-134. [278]
- [183] C. Payan, Sur le nombre d'absorption d'un graphe simple. (French) Colloque sur la Théorie des Graphes (Paris, 1974). Cahiers Centre Etudes Recherche Opér. **17** (1975) 307-317. [325, 550]
- [184] C. Payan and N. H. Xuong, Domination-balanced graphs. *J. Graph Theory* **6** (1982) 23-32. [325]
- [185] E. Pegg, Review of *The Colossal Book of Mathematics* by Martin Gardner. *Notices of the AMS.* **49** (2002) 1085-1086. [475]
- [186] J. Petersen, Die Theorie der regulären Graphen. *Acta Math.* **15** (1891) 193-220. [13, 313, 339]
- [187] J. Petersen, Sur le théorème de Tait. *L'Intermédiaire des Mathématiciens* **5** (1898) 225-227. [314, 336]
- [188] J. Plesník, Critical graphs of given diameter. *Acta Fac. Rerum Natur. Univ. Comenian. Math.* **30** (1975) 71-93. [101]
- [189] L. Pósa, Hamiltonian circuits in random graphs. *Discrete Math.* **14** (1976) 359-364. [560]
- [190] R. C. Prim, Shortest connection networks and some generalizations. *Bell Syst. Tech. J.* **36** (1957) 1389-1401. [83]

- [191] H. Prüfer, Neuer Beweis eines Satzes über Permutationen. *Arch. Math. Phys.* **27** (1918) 142-144. [69]
- [192] R. Rado, A theorem on independence relations. *Quart. J. Math. Oxford Ser.* **13** (1942) 83-89. [321]
- [193] F. P. Ramsey, On a problem of formal logic. *Proc. London Math. Soc.* **30** (1930) 264-286. [523]
- [194] F. P. Ramsey, *The Foundations of Mathematics, and Other Logical Essays*. Harcourt, Brace and Company, New York (1931). [523]
- [195] D. K. Ray-Chaudhuri and R. M. Wilson, Solution of Kirkman's school-girl problem. *Combinatorics (Proc. Sympos. Pure Math., Vol. XIX, Univ. California, Los Angeles, Calif., 1968)*, Amer. Math. Soc., Providence, RI. (1971) 187-203. [348]
- [196] R. C. Read, An introduction to chromatic polynomials. *J. Combin. Theory* **4** (1968) 52-71. [438, 440]
- [197] L. Rédei, Ein kombinatorischer Satz. *Acta Litt. Szeged* **7** (1934) 39-43. [180, 181]
- [198] B. Reed,  $\omega$ ,  $\Delta$ , and  $\chi$ . *J. Graph Theory* **27** (1998) 177-212. [374, 381]
- [199] S. Říha, A new proof of the theorem by Fleischner. *J. Combin. Theory Ser. B* **52** (1991) 117-123. [147]
- [200] R. D. Ringeisen and L. W. Beineke, The crossing number of  $C_3 \times C_n$ . *J. Combin. Theory Ser. B* **24** (1978) 134-136. [281]
- [201] G. Ringel, Über drei kombinatorische Probleme am  $n$ -dimensionalen Würfel und Würfelgitter. *Abh. Math. Sem. Univ. Hamburg* **20** (1955) 10-19. [295]
- [202] G. Ringel, Problem 25. *Theory of Graphs and its Applications*. Nakl. ČSAV, Prague (1964) 162. [353]
- [203] G. Ringel, Das Geschlecht des vollständigen paaren Graphen. *Abh. Math. Sem. Univ. Hamburg* **28** (1965) 139-150. [294]
- [204] G. Ringel and J. W. T. Youngs, Solution of the Heawood map-coloring problem. *Proc. Nat. Acad. Sci. USA* **60** (1968) 438-445. [294, 447]
- [205] H. E. Robbins, A theorem on graphs, with an application to a problem in traffic control. *Amer. Math. Monthly* **46** (1939) 281-283. [166]
- [206] N. Robertson and P. D. Seymour, Graph minors. XX. Wagner's conjecture. *J. Combin. Theory Ser. B* **92** (2004) 325-357. [300]

- [207] N. Robertson, P. Seymour and R. Thomas, Hadwiger's conjecture for  $K_6$ -free graphs. *Combinatorica* **13** (1993) 279-361. [437]
- [208] N. Robertson, P. D. Seymour and R. Thomas, Tutte's edge-colouring conjecture. *J. Combin. Theory Ser. B* **70** (1997) 166-183. [489]
- [209] N. Robertson, P. D. Seymour and R. Thomas, Excluded minors in cubic graphs. arXiv:1403.2118v1. [489]
- [210] N. Robertson, P. D. Seymour and R. Thomas, Cyclically 5-connected cubic graphs. arXiv:1503.02298. [489]
- [211] A. Rosa, On certain valuations of the vertices of a graph. *Theory of Graphs*. Gordon and Breach, New York (1967) 349-355. [351, 352, 354, 355]
- [212] B. Roy, Nombre chromatique et plus longs chemins d'un graph. *Rev AFIRO* **1** (1967) 127-132. [383]
- [213] M. Šajna, Cycle Decompositions III: Complete graphs and fixed length cycles. *J. Combin. Des.* **10** (2002) 27-78. [348, 350]
- [214] D. P. Sanders, P. D. Seymour and R. Thomas, Edge 3-coloring cubic doublecross graphs. Preprint. [489]
- [215] D. P. Sanders and R. Thomas, Edge 3-coloring cubic apex graphs. Preprint. [489]
- [216] D. P. Sanders and Y. Zhao, Planar graphs of maximum degree seven are class I. *J. Combin. Theory Ser. B* **83** (2001) 201-212. [473]
- [217] S. Schuster, Interpolation theorem for the number of end-vertices of spanning trees. *J. Graph Theory* **7** (1983) 203-208. [75]
- [218] M. Sekanina, On an ordering of the set of vertices of a connected graph. *Spisy Přírod Fak. Univ. Brno.* (1960) 137-141. [146]
- [219] P. D. Seymour, Sums of circuits. *Graph Theory and Related Topics*. Academic Press, New York (1979) 341-355. [491]
- [220] P. D. Seymour, Nowhere-zero 6-flows. *J. Combin. Theory Ser. B* **30** (1981) 130-135. [489]
- [221] C. E. Shannon, A theorem on coloring the lines of a network. *J. Math. Physics* **28** (1949) 148-151. [459]
- [222] T. Slivnik, Short proof of Galvin's theorem on the list-chromatic index of a bipartite multigraph. *Combin. Probab. Comput.* **5** (1996) 91-94. [493]
- [223] J. Spencer, Ramsey's theorem - a new lower bound. *J. Combin. Theory* **18 A** (1975) 108-115. [547]

- [224] J. Spencer, Ten Lectures on the Probabilistic Method, 2ed. CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM 1994. [547, 560]
- [225] S. K. Stein, Convex maps. *Proc. Amer. Math. Soc.* **2** (1951) 464-466. [283]
- [226] B. M. Stewart, On a theorem of Nordhaus and Gaddum. *J. Combin. Theory* **6** (1969) 217-218. [381]
- [227] P. K. Stockmeyer, The falsity of the reconstruction conjecture for tournaments. *J. Graph Theory* **1** (1977) 19-25. [230]
- [228] P. K. Stockmeyer, Erratum to: "The falsity of the reconstruction conjecture for tournaments." *J. Graph Theory* **62** (2009) 199-200. [230]
- [229] G. Szekeres, Polyhedral decompositions of cubic graphs. *Bull. Austral. Math. Soc.* **8** (1973) 367-387. [474, 491]
- [230] G. Szekeres and H. S. Wilf, An inequality for the chromatic number of a graph. *J. Combin. Theory* **4** (1968) 1-3. [377]
- [231] T. Szele, Kombinatorische Untersuchungen über gerichtete vollständige Graphen. *Mat. Fiz. Lapok* **50** (1943) 223-256. [181, 549]
- [232] P. G. Tait, Remarks on the colouring of maps. *Proc. Royal Soc. Edinburgh* **10** (1880) 729. [468]
- [233] C. Thomassen, Landau's characterization of tournament score sequences, in *The Theory and Applications of Graphs*, Wiley, New York (1981) 589-591. [176]
- [234] C. Thomassen, Every planar graph is 5-choosable. *J. Combin. Theory Ser. B* **62** (1994) 180-181. [428]
- [235] C. Thomassen, Some remarks on Hajós' conjecture. *J. Combin. Theory Ser. B* **93** (2005) 95-105. [436]
- [236] S. Toida, Properties of an Euler graph. *J. Franklin Inst.* **295** (1973) 343-345. [122]
- [237] P. Turán, Eine Extremalaufgabe aus der Graphentheorie. *Mat. Fiz. Lapok* **48** (1941) 436-452. [504]
- [238] P. Turán, A note of welcome. *J. Graph Theory* **1** (1977) 7-9. [278]
- [239] W. T. Tutte, On hamiltonian circuits. *J. London Math. Soc.* **21** (1946) 98-101. [266]
- [240] W. T. Tutte, A family of cubical graphs. *Proc. Cambridge Philos. Soc.* **43** (1947) 459-474. [510, 514]



- [241] W. T. Tutte, On the imbedding of linear graphs in surfaces. *Proc. London Math. Soc.* **51** (1949) 474-483. [489]
- [242] W. T. Tutte, A short proof of the factor theorem for finite graphs. *Canad. J. Math.* **6** (1954) 347-352. [311]
- [243] W. T. Tutte, A theorem of planar graphs. *Trans. Amer. Math. Soc.* **82** (1956) 99-116. [266]
- [244] O. Veblen, An application of modular equations in analysis situs. *Ann. of Math.* **14** (1912) 86-94. [121]
- [245] L. M. Vitaver, Determination of minimal coloring of vertices of a graph by means of Boolean powers of the incidence matrix. (Russian) *Dokl. Akad. Nauk. SSSR* **147** (1962) 758-759. [383]
- [246] V. G. Vizing, On an estimate of the chromatic class of a  $p$ -graph. (Russian) *Diskret. Analiz.* **3** (1964) 25-30. [455, 459, 496]
- [247] V. G. Vizing, Critical graphs with given chromatic class. *Metody Diskret. Analiz.* **5** (1965) 9-17. [473]
- [248] V. G. Vizing, Coloring the vertices of a graph in prescribed colors (Russian) *Diskret. Analiz* **29** (1976) 3-10. [406, 428]
- [249] M. Voigt, List colourings of planar graphs. *Discrete Math.* **120** (1993) 215-219. [428]
- [250] K. Wagner, Bemerkungen zum Vierfarbenproblem. *Jber. Deutsch. Math-Verein.* **46** (1936) 26-32. [283]
- [251] K. Wagner, Über eine Eigenschaft der ebene Komplexe. *Math. Ann.* **114** (1937) 570-590. [260, 437]
- [252] D. J. A. Welsh and M. B. Powell, An upper bound for the chromatic number of a graph and its application to timetabling problems. *Computer J.* **10** (1967) 85-86. [377]
- [253] P. Wernicke, Über den kartographischen Vierfarbensatz. *Math. Ann.* **58** (1904) 413-426. [250]
- [254] A. T. White, *Graphs of Groups on Surfaces, Interactions and Models*. North-Holland, Amsterdam (2001). [280]
- [255] H. Whitney, Congruent graphs and the connectivity of graphs. *Amer. J. Math.* **54** (1932) 150-168. [98, 106, 149]
- [256] H. Whitney, The coloring of graphs. *Ann. of Math.* **33** (1932) 688-718. [438, 441]

- [257] J. E. Williamson, Panconnected graphs. II. *Period. Math. Hungar.* **8** (1977) 105-116. [144]
- [258] R. M. Wilson, Decompositions of complete graphs into subgraphs isomorphic to a given graph. *Congr. Numer.* **15** (1976) 647-659. [345]
- [259] D. R. Woodall, Sufficient conditions for circuits in graphs. *Proc. London Math. Soc.* **24** (1972) 739-755. [169]
- [260] D. R. Woodall, Cyclic-order graphs and Zarankiewicz's crossing-number conjecture. *J. Graph Theory* **17** (1993) 657-671. [279]
- [261] J. W. T. Youngs, Minimal imbeddings and the genus of a graph. *J. Math. Mech.* **12** (1963) 303-315. [292]
- [262] K. Zarankiewicz, On a problem of P. Turán concerning graphs. *Fund. Math.* **41** (1954) 137-145. [279]
- [263] A. A. Zykov, On some properties of linear complexes (Russian). *Mat. Sbornik N. S.* **24** (1949) 163-188. [394, 440]

## Supplemental References

- N. Alon and J. H. Spencer, *The Probabilistic Method, Third Edition*. Wiley, New York (2008).
- A. Benjamin, G. Chartrand and P. Zhang, *The Fascinating World of Graph Theory*. Princeton University Press, Princeton, NJ (2015).
- N. L. Biggs, E. K. Lloyd and R. J. Wilson, *Graph Theory, 1736–1936*. Oxford University Press, London (1976).
- J. A. Bondy and U. S. R. Murty, *Graph Theory*. Springer, New York (2008).
- G. Chartrand and P. Zhang, *Chromatic Graph Theory*. Chapman & Hall/CRC Press, Boca Raton, FL (2009).
- R. Diestel, *Graph Theory, Third Edition*. Springer-Verlag, Berlin (2005).
- J. Gross, J. Yellen and P. Zhang (editors), *Handbook of Graph Theory, Second Edition*. CRC Press, Boca Raton, FL (2014).
- D. B. West, *Introduction to Graph Theory, Second Edition*. Prentice-Hall, Upper Saddle River, NJ (2001).
- R. Wilson, *Four Colors Suffice: How the Map Problem Was Solved*. Princeton University Press, Princeton, NJ (2002).