

Quantum Monte Carlo calculations of energy gaps from first-principles

Ryan J. Hunt

Department of Physics,
Lancaster University


Lancaster
University



What's the problem?

We'd like to be able to predictively model the (opto)electronic behaviour of materials. Because this could be *useful*.

- ▶ Specifically, $\Delta_{\text{Ex.}}$, Δ_{QP} , and E_{B}^{X} in semiconductors.¹
- ▶ Don't define “material heaven”, but are a start.
- ▶ (The blue LED is blue for a reason.)



¹FYI work discussed here is in: **Hunt et al.**, Phys. Rev. B **98(7)** (2018).

The **quasiparticle gap**, Δ_{QP} , is defined as the difference between the CBM and the VBM:

$$\begin{aligned}\Delta_{\text{QP}}(\mathbf{k}_f, \mathbf{k}_t) &= \mathcal{E}_{\text{CBM}}(\mathbf{k}_t) - \mathcal{E}_{\text{VBM}}(\mathbf{k}_f) \\ &= [E_{N+1}(\mathbf{k}_t) - E_N(\mathbf{k}_t)] - [E_N(\mathbf{k}_f) - E_{N-1}(\mathbf{k}_f)] \\ &= E_{N+1}(\mathbf{k}_t) + E_{N-1}(\mathbf{k}_f) - E_N(\mathbf{k}_t) - E_N(\mathbf{k}_f),\end{aligned}\tag{1}$$

The **excitonic gap**, $\Delta_{\text{Ex.}}$, is defined as the energy difference between an excited N-electron state and the ground N-electron state:

$$\Delta_{\text{Ex.}}(\mathbf{k}_f, \mathbf{k}_t) = E_N^+(\mathbf{k}_f, \mathbf{k}_t) - E_N,\tag{2}$$

Their difference is the **exciton binding**.²

² The interaction energy of a quasielectron at \mathbf{k}_t and a quasihole at \mathbf{k}_f .

The Theory of Everything

$$i\hbar \frac{\partial \Psi}{\partial t} = \mathcal{H} \Psi$$

$$\mathcal{H} = - \sum_j^N \frac{\hbar^2}{2m_j} \nabla_j^2 - \sum_\alpha^M \frac{\hbar^2}{2m_\alpha} \nabla_\alpha^2 - \sum_j^N \sum_\alpha^M \frac{Z_\alpha e^2}{|r_j - R_\alpha|} \\ + \sum_{j < k}^N \frac{e^2}{|r_j - r_k|} + \sum_{\alpha < \beta}^M \frac{Z_\alpha Z_\beta e^2}{|R_\alpha - R_\beta|}$$

- | | | | |
|----------|-----------|--------------|------------------|
| * Air | * Steel | * Paper | * Vitamins |
| * Water | * Plastic | * Dynamite | * Ham Sandwiches |
| * Fire | * Glass | * Antifreeze | * Ebola Virus |
| * Rocks | * Wood | * Glue | * Economists |
| * Cement | * Asphalt | * Dyes | * ... |

Figure 1: Introductory slide from Laughlin's Nobel lecture.

What else?

- ▶ Density functional theory (or HF | hybrids)
 - ▶ Take differences in Kohn-Sham (Hartree-Fock) SP eigenvalues.
- ▶ Many-body perturbation theory (GW | GW -BSE | MPn)
 - ▶ QP energies from QP equation (feat. self-energy, $\Sigma(\mathbf{k}, \omega)$).
- ▶ Quantum chemistry (post HF | CC | CI | FCI)
 - ▶ Most similar to present: direct calculation of total energies.


Either too crude, too scattered, or too expensive.

QMC methods:

- ✓ Are highly accurate, and systematically improvable.
- ✓ Are non-perturbative, and treat correlation effects exactly.
- ✓ Have $\mathcal{O}(N_e^3)$ cost, not much worse in “abnormal” cases.

Proof? Lots available, see reviews,³ or below.⁴

Figure 2: The basis of much modern (computational) electronic structure theory.



³ W. M. C. Foulkes et al., Rev. Mod. Phys. 73 (2001), R. J. Needs et al., J. Phys. Condens. Matter 22 (2009).

⁴ D. M. Ceperley and B. J. Alder, Phys. Rev. Lett. 45 (1980).

Variational Monte Carlo

- ▶ Endow a *trial* wavefunction with variational freedom:

$$\Psi(\mathbf{R}) = \underbrace{\exp \left[\mathcal{J}_{\{\alpha\}}(\mathbf{R}) \right]}_{\text{Our additions}} \times \underbrace{\mathcal{D}(\mathbf{R})}_{\text{DFT, HF, ...}}, \quad (3)$$

and pick $\{\alpha\}$.

- ▶ MC integration used, for example, to evaluate

$$\langle \Psi | \hat{\mathcal{H}} | \Psi \rangle = \int d\mathbf{R} |\Psi(\mathbf{R})|^2 \left[\frac{\hat{\mathcal{H}}\Psi(\mathbf{R})}{\Psi(\mathbf{R})} \right] \approx \sum_i \frac{\mathcal{H}(\mathbf{R}_i)\Psi(\mathbf{R}_i)}{\Psi(\mathbf{R}_i)}, \quad (4)$$

($\{\mathbf{R}_i\}$ distributed as $|\Psi(\mathbf{R})|^2$).

⁵ W. M. C. Foulkes et al., Rev. Mod. Phys. 73 (2001).

Diffusion Monte Carlo

- DMC is a stochastic projector-based method for solving

$$\hat{\mathcal{H}} \Psi(\mathbf{R}, \tau) = (E_T - \partial_\tau) \Psi(\mathbf{R}, \tau), \quad (5)$$

or, if you like

$$\Psi(\mathbf{R}, \tau + \Delta\tau) = \int G(\mathbf{R} \leftarrow \mathbf{R}', \Delta\tau) \Psi(\mathbf{R}', \tau) d\mathbf{R}'. \quad (6)$$

- **Separable** ($\partial_\tau \hat{\mathcal{H}} = 0$)⁶

$$\Psi(0) = \sum_n c_n \Phi_n \implies \Psi(\tau) = \sum_n c_n \Phi_n \exp[-(\mathcal{E}_n - E_T)\tau] \quad (7)$$

⁶ $\{\Phi_i\} \rightarrow$ complete basis of eigenstates of the interacting problem.

Effectively we take:

$$\lim_{\tau \rightarrow \infty} \Psi(\tau) \sim \Phi_0, \quad (8)$$

by having the DMC Green's function take configurations $\mathbf{R}' \rightarrow \mathbf{R}$, with caveats:

- ▶ **Time steps**: know $G(\mathbf{R} \leftarrow \mathbf{R}', \Delta\tau)$ in limit of small $\Delta\tau$.
- ▶ **Population control**: number of walkers in DMC fluctuates. Control mechanism introduces a bias.
- ▶ **Finite-size (FS) effects**: extrapolation to TD limit a necessity.
- ▶ **Fixed-node approximation**: (non-local) antisymmetry enforced by (local) boundary condition ($\Psi = 0$ surface is fixed).

Gaps: expect some of these to matter less!

Briefly:


- ▶ QP gap $\rightarrow E_{N,N\pm 1}$ (VP on each *ground* state).
- ▶ Ex. gap \rightarrow *may* have VP on E^+ . *May* only have at VMC level. FN-DMC VP obtained in special circumstances.⁷
- ▶ (FN constraint means effective VP)

⁷ W. M. C. Foulkes et al., Phys. Rev. B **60** (1999).

We've studied Si, α -SiO₂, and cubic BN in the current work. Previous QMC studies had claimed success in evaluation of “QMC band structures”,⁸ minus discussions of:

- ▶ Finite-size errors.
- ▶ Fixed-nodal errors.
- ▶ Δ_{QP} vs. $\Delta_{\text{Ex.}}$.

Will concentrate on Si here, exploring the above.



⁸ P. R. C. Kent et al., Phys. Rev. B 57 (1998), A. J. Williamson et al., Phys. Rev. B 57 (1998).

- Able only to simulate a finite *chunk* of material (supercell), under PBCs. Excitations “1/N” effects. Need statistical accuracy + careful FS treatment.




Figure 3: Uncorrected SJ-DMC gaps of Si. FS effect characteristic and quantifiable, largely from image-interactions.

- Then why do Δ_{QP} & Δ_{Ex} behave same?

- ▶ Probe with Backflow transformation:

$$\mathbf{r}_i \rightarrow \mathbf{x}_i = \mathbf{r}_i + \boldsymbol{\xi}_i(\mathbf{R}) \quad (9)$$

which can change nodal surface.⁹

- ▶ Tested $\Delta_{\text{QP/Ex}}(\Gamma_v \rightarrow \Gamma_c)$ and $\Delta_{\text{QP}}(\Gamma_v \xrightarrow{\sim} \text{CBM})$, in $2 \times 2 \times 2$ supercell.
- ▶ We find that backflow leads to a reduction in gaps, of at least 0.2 eV, but upto 0.3–0.4 eV when one re-optimises $\boldsymbol{\xi}_i$.¹⁰

⁹ P. Lopez Rios et al., Phys. Rev. E 74 (2006).

¹⁰ C.f. controllable uncertainty: $\mathcal{O}(0.1 \text{ eV})$ for each of pseudopotentials, statistics, NLO FS effects (?).

A direct gap 2D semiconductor, with large exciton binding energy.¹¹





Figure 4: PL measurements of Phosphorene n -layers.

- ▶ Do **not** expect $\Delta_{QP} \sim \Delta_{Ex}$.
- ▶ FS effects in $\Delta_{QP/Ex}$ much more important.

¹¹ J. Yang et al., Light Sci. Appl. 4 (2015).



- ▶ We want to model a *free* excitonic complex
- ▶ Perform supercell calculation (SC characteristic size L), subject to periodic BCs
- ▶ Hence incur an unphysical image-interaction

- ▶ Need to remove E_{int} . **How does it scale?**




Figure 5: Unscreened (left) and screened (right) field lines from point charges at $\rho = z = 0$.

- With 2D screening (Keldysh interaction), charge-quadrupole interaction¹² leads to expected scaling which is $\mathcal{O}(L^{-2})$.

¹² Note no dipole in inversion symmetric system!

Δ_{QP}


- ▶ Similar image effects, easier to manage.
- ▶ Subtract single-particle v_{M} ($\mathcal{O}(L^{-1})$).
- ▶ From regularized lattice sum over screened interaction (\sim Ewald sum).

$$\sum_{\mathbf{R}} W(\mathbf{r} - \mathbf{R}) \rightarrow \overbrace{\sum_{\mathbf{R}} V(\mathbf{r} - \mathbf{R})}^{\text{Ewald}} + \underbrace{\sum_{\mathbf{R}} \delta V(\mathbf{r} - \mathbf{R})}_{\text{safe}}. \quad (10)$$

- ▶ “*Safety*”: $\lim_{r \rightarrow \infty} \delta V(r) = 0$.

- ▶ FSE appear to scale as argued.
- ▶ Big gaps ($\epsilon!$),¹³ but good agreement w/ Gaufres *et al.*¹⁴
- ▶ Phonon renormalisation ~ 0.17 eV @ 300K.¹⁵

Figure 6: QMC energy gaps in phosphorene vs. system size.



¹³ Also, this is not due to FN error! Backflow has $\mathcal{O}(0.05$ eV) effect here.

¹⁴ This result is unpublished, so far, but was presented at GW 2018 by A. Loiseau. $\Delta_{EX} = 1.95$ eV.

¹⁵ Via Tomeu Monserrat, also as yet unpublished.

Another approach is to consider passivated (finite) clusters. Here FS effect is kinetic in origin (confinement).¹⁶

- ▶ FS converge faster (QP gap $\mathcal{O}(L^{-2})$ by default), **but...**
- ▶ State under study may not be relevant¹⁷...




Figure 7: Band charge densities in $C_{29}H_{36}$.

¹⁶ T. Frank et al., arXiv:1805.10823 (2018).

¹⁷ N. D. Drummond et al., Phys. Rev. Lett. **95** (2005).

Frank *et al.* have also studied phosphorene. We're dissatisfied with their approach. Why?

- ▶ Used cluster calculations to argue scaling in bulk calculations.
- ▶ Calculated the *wrong gap*:

Figure 8: Excerpt from preprint.

by QMC methods within the error bars²⁸. The gap Δ_f was extracted as the singlet-singlet vertical excitation energy. Here $\Delta_f \approx E_v^{ss} = E_1^s - E_0^s$, with E_0 and E_1 being, respectively, the ground- and the first excited-states obtained by fixed-node QMC³⁸ not allowing any

- ▶ Our excitonic gap (2.2(2) eV) agrees with their “quasiparticle” gap (2.4 eV) ☺.

18

¹⁸ Guessed wrong scaling exponent (1/N) for QP gap, but this isn't a QP gap! Just so happen to have calculated and taken TD limit for an excitonic gap. Assuming they've done the calculations correctly, a good test of our result!

- ▶ QMC methods offer a direct, real-space approach to the many-body problem.
- ▶ They allow for accurate determination of energy gaps from first-principles in one, two and three-dimensional systems.
- ▶ They can be systematically extended, and treat various important pieces of physics **exactly**.

Lancaster:

- ▶ Neil Drummond
- ▶ Marcin Szyniszewski

Japan (JAIST):

- ▶ Ryo Maezono
- ▶ Genki Prayogo

Thanks for your attention!