

Modelling charge complexes in 2D semiconductors, and their heterostructures

CDT Summer Conference 2018

Ryan J. Hunt¹, Neil D. Drummond¹
and Vladimir I. Fal'ko²



¹Department of Physics,
Lancaster University

²National Graphene Institute,
University of Manchester



2nd July, 2018

“Modelling”

For me, this means using *quantum Monte Carlo* (QMC) methods. QMC relies on random sampling techniques, which can be used (with CASINO¹) to solve the many-body Schrödinger equation:

$$\left[-\frac{1}{2} \nabla_{\mathbf{R}}^2 + V(\mathbf{R}) \right] \Psi(\mathbf{R}) = E \Psi(\mathbf{R}), \quad \mathbf{R} = \{\mathbf{r}_1, \dots, \mathbf{r}_N\} \quad (1)$$

Main point: This is a **hard problem**. N might be 6,² but it might be 60, or even 600³...

- ▶ QMC deals with it, without *excessively muddying* $V(\mathbf{R})$.

¹ R. J. Needs et al., J. Phys. Condens. Matter **22** (2009).


² \sim charge complex in the effective mass approximation

³ \sim charge complex in “heroic” QMC

Quintons in monolayer TMDs⁶

A.k.a. “charged biexciton”, but that’s no fun.

Figure 1: A *quinton* (XX^- , pictured) is a *hypothetical* bound state of five particles in a semiconductor.



We predicted that quintons would be stable in monolayers of TMDCs. Recent experiments corroborate this claim.^{4,5}

Disclaimer: Our model neglects valley effects, these would act to alter interactions at short-range.

⁴ K. Hao et al., Nat. Commun. **8** (2017).

⁵ 1806.03775, 1805.04950, 1802.10247

⁶ E. Mostaani et al., Phys. Rev. B **96** (2017).

Interlayer complexes in heterobilayers of TMDs⁷




Figure 2: (one) Diagram (of four) representing D^0 recombination with h^+ .


⁷ M. Danovich et al., Phys. Rev. B **97** (2018).

“Heroic” electronic structure calculations⁸

Q Can we access the optoelectronic properties of a material by knowing only the atomic numbers of its constituents?

A Yeah, but beware:

- ▶ It's not cheap (CPU time).
 - ▶ Technical difficulties: finite size effects, variational collapse.
 - ▶ Sometimes, the electronic problem is not the whole story...
-
- ▶ Paper is mostly technical, we do consider excitons in phosphorene.




⁸ R. J. Hunt et al., arXiv:1806.04750 (submitted to PRB) (2018).

The Future

My next 2D materials project will concern charge complexes at finite density.

- ▶ Superfluidity of excitons in vdW crystals? (needs thought)
- ▶ The **Mahan exciton** (ME) in 2D “metals”?⁹ (thinking done)



Coulomb 2DEG: $?_1 \approx 2a_0^*$, $?_2 \approx 4a_0^*$

⁹ 2D semiconductors, at high density.

The Mahan Exciton

“Excitons in Metals”

Formation coincides with onset of absorption in metals,¹⁰ in a feature known as the “Fermi Edge Singularity” (FES).¹¹

Anderson and **Mahan** studied this, FES boils down to spectral function:

$$A(\omega) = A^{(0)}(\omega) \underbrace{\Theta(\omega - \omega_T)}_{\text{Orth. Catastrophe}} \overbrace{\left(\frac{\xi_0}{\omega - \omega_T} \right)^\alpha}^{\text{e-h int.}}. \quad (2)$$

(In practice, you’d expect to see an increasingly asymmetric peak appear in PL measurements at lower and lower temperatures.)

¹⁰ Most clearly at low temperature...

¹¹ G. D. Mahan, (Plenum, New York, 2000), G. D. Mahan, Phys. Rev. Lett. **18** (1967).

ME II

- ▶ ME a distinctly many-body effect – “exciton” is a **misnomer**.
- ▶ To properly describe ME in doped 2D SCs, would need to account for all electrons, ions... Such “heroic”-scale realistic computations are hard.¹²
- ▶ Can we do *something*...? **I think so**.
- ▶ ME has been studied in (Coulomb) 2DEG,¹³ but key ingredient in (*bona fide*) 2D is screening. Options:
 1. “Heroic” calculations → screening **emergent** from $V(\mathbf{R})$.
 2. Devise a clever model → screening **modelled**. (Other clever models exist for the crossover...¹⁴)

¹² At least for mere mortals with finite computer power.

¹³ G. G. Spink et al., Phys. Rev. B **94** (2016).

¹⁴ D. K. Efimkin and A. H. MacDonald, Phys. Rev. B **95** (2017), D. K. Efimkin and A. H. MacDonald, Phys. Rev. B **97** (2018).

Thank you all for listening, and thanks to my collaborators:

Manchester (NGI):

- ▶ M. Danovich
- ▶ D. A. Riuz-Tijerina
- ▶ V. I. Fal'ko

Cambridge (CGC):

- ▶ E. Mostaani







Japan (JAIST):

- ▶ R. Maezono
- ▶ G. Prayogo

Lancaster:

- ▶ D. M. Thomas
- ▶ M. Szyniszewski
- ▶ N. D. Drummond
- ▶ O. Witham (Frankfurt)

References I

-  R. J. Needs, M. D. Towler, N. D. Drummond, and P. López Ríos, “Continuum variational and diffusion quantum monte carlo calculations”, J. Phys. Condens. Matter **22**, 023201 (2009).
-  K. Hao et al., “Neutral and charged inter-valley biexcitons in monolayer mose₂”, Nat. Commun. **8**, 15552 (2017).
-  E. Mostaani, M. Szyniszewski, C. H. Price, R. Maezono, M. Danovich, R. J. Hunt, N. D. Drummond, and V. I. Fal’ko, “Diffusion quantum monte carlo study of excitonic complexes in two-dimensional transition-metal dichalcogenides”, Phys. Rev. B **96**, 075431 (2017).
-  M. Danovich, D. A. Ruiz-Tijerina, R. J. Hunt, M. Szyniszewski, N. D. Drummond, and V. I. Fal’ko, “Localized interlayer complexes in heterobilayer transition metal dichalcogenides”, Phys. Rev. B **97**, 195452 (2018).
-  R. J. Hunt, M. Szyniszewski, G. I. Prayogo, R. Maezono, and N. D. Drummond, “Quantum Monte Carlo calculations of energy gaps from first principles”, arXiv:1806.04750 (submitted to PRB) (2018).
-  G. D. Mahan, Many Particle Physics, Third Edition, (Plenum, New York, 2000).

References II



G. D. Mahan, “Excitons in metals”, Phys. Rev. Lett. **18**, 448–450 (1967).



G. G. Spink, P. López Ríos, N. D. Drummond, and R. J. Needs, “Trion formation in a two-dimensional hole-doped electron gas”, Phys. Rev. B **94**, 041410 (2016).



D. K. Efimkin and A. H. MacDonald, “Many-body theory of trion absorption features in two-dimensional semiconductors”, Phys. Rev. B **95**, 035417 (2017).



D. K. Efimkin and A. H. MacDonald, “Exciton-polarons in doped semiconductors in a strong magnetic field”, Phys. Rev. B **97**, 235432 (2018).