

Recent findings on genetics of systemic autoimmune diseases

Angélica Delgado-Vega¹, Elena Sánchez², Sara Löfgren¹,
 Casimiro Castillejo-López¹ and Marta E Alarcón-Riquelme^{1,2,3}

Association studies of over 1 million SNPs capturing most of the human genome common variation became possible thanks to the information provided by the HapMap International project and the development of high-throughput genotyping technologies at accessible prices. Genome-wide scans analyzing thousands of individuals have now identified most if not all of the major genes involved in susceptibility for several systemic autoimmune diseases. In particular, results for rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), and systemic sclerosis (SSc) are reviewed here. While most genes are shared between diseases, few seem to be unique reflecting that we still are long before knowing all genes, their interactions with other genes and the environment and their impact on biological functions.

Addresses

¹ Department of Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Dag Hammarskjölds väg 20, 751 85 Uppsala, Sweden

² Arthritis and Immunology Program, Oklahoma Medical Research Foundation, 825 NE 13th St., Oklahoma City 73104, OK, United States

³ Center for Genomics and Oncological Research Pfizer, University of Granada, Junta de Andalucía, Avenida de la Ilustración 114, Granada 18007, Spain

Corresponding author: Alarcón-Riquelme, Marta E (alarconm@omrf.org)

Current Opinion in Immunology 2010, 22:698–705

This review comes from a themed issue on
 Autoimmunity
 Edited by Kazuhiko Yamamoto and Mark Shlomchik

Available online 20th October 2010

0952-7915/\$ – see front matter

© 2010 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.coi.2010.09.002

Introduction

Previous studies on the genetics of complex diseases used small sample sizes leading to inconclusive results, with the exception of the strong genetic association between genes within the major histocompatibility complex (MHC) and the various autoimmune diseases. While GWAS have provided some surprises, they have also confirmed a few of the old results, but not all. The most important genes identified previously, *IRF5* and *STAT4* for SLE and *PTPN22* and *STAT4* for RA stand [1]. In this review we focus only on those genes identified in the past two to three years. As GWAS with extensive coverage have been used, except for a few examples (*PDCD1* is a relevant one, as SNPs within this gene are lacking in the

GWAS arrays), most of the genome has been screened with thousands of samples providing an overall picture of genetic susceptibility. We present only those genes that have been clearly replicated and whose role in genetic susceptibility is beyond doubt or the result of powerful GWAS with replication using large sample sizes.

Genetics of SLE

SLE is considered as a prototypic autoimmune disease, characterized by production of antinuclear autoantibodies, immune-complex deposition, and subsequent multiple organ damage.

The understanding of the genetic basis of SLE has expanded enormously over the past couple of years, driven principally by technological advances and the assessment of six genome-wide association studies (GWAS) in the past two years, in Caucasians and Asians [2,3,4,5,6,7].

The great majority of the identified genes are involved in innate and adaptive immune responses. The convincingly associated genes, as summarized in Table 1, are mainly implicated in immune-complex (IC) processing, T/B cell signaling and/or and Toll-like receptor (TLR), and type I interferon signaling. The strongest association in the GWAS era is undoubtedly the MHC region, but because it consists of strongly correlated polymorphisms in a ~6 Mb region involving more than 400 genes it remains a challenge. Recent observations by Fernando *et al.* have shown at least two independent genetic effects within the MHC region in SLE: one signal is provided by the *HLA-DRB1*0301* allele, and the second within the class III region specifically detecting the 6th intron of the *SKIVL2* gene [8]. A more recent study genotyping over 1400 variants within the MHC region and over 10,000 individuals with various autoimmune diseases including SLE and RA showed the top disease-specific signal for SLE to be SNP [8,9] located between *TNXB* and *CREBL1* and the *HLA DRB1*0301* [9].

Impaired IC clearance and deposition is an important pathological aspect in SLE. Susceptibility genes with important roles in IC processing known from previous studies are the FcGR family of genes and more recently *ITGAM* [10] coding for the surface antigen CD11b (or CR3). Signal transduction in immune cells, particularly T and B cells, is another pathway that has revealed to contain multiple lupus susceptibility genes, modulating T cell signaling such as *TNFSF4* (OX40L) [11,12] and

Table 1

Susceptibility Genes for SLE

Chromosome	Gene	SNPs	Population	References
6p21	<i>HLA region</i>	DRB1*0301 and several other alleles	European, several Asian, African-American, mixed European-Amerindian	[8,9]
7q32	<i>IRF5</i>	5bp promoter indel, rs2004640, rs2070197, rs10954213	European, several Asian, mixed European-Amerindian, African-American	[3*,4**,5**,7,20**]
2q32	<i>STAT4</i>	rs7574865, rs3821236, rs7601754	European, mixed European-Amerindian, several Asian, African-American	[3*,4**,5**,7,20**]
6q23	<i>TNFAIP3</i>	rs5029939, rs2230926	European, Asian, African-American	[2,3*,4**,5**,7,17*,20**]
16p11	<i>ITGAM</i>	rs9888739, rs1143679, rs4548893	European, mixed European-Amerindian, Asian, African-American	[3*,4**,5**,7,10**,20**]
4q24	<i>BANK1</i>	rs10516487, rs17266594, rs3733197	European, European-Amerindian, Asian	[3*,6**,7,14]
1p13	<i>PTPN22</i>	rs2476601	European	[4**]
8p23	<i>BLK</i>	rs13277113, rs2736340	European, several Asian	[3*,4**,5**,7,15,20**]
2q37	<i>PDCD1 (CD279)</i>	PD1.3A	European, European-Amerindian, Chinese	[49]
1q25	<i>TNFSF4</i>	Risk haplotype; rs3850641	European, Asian	[3*,7,11,12,14,20**]
18q22.3	<i>CD226</i>	rs763361, rs727088	European, European-Amerindian	[13*]
1q21-23	<i>FCGR2A</i>	ARG131HIS	European, European-Amerindian, African American	[4**,5**,20**]
19p13.2	<i>TYK2</i>	rs280519	European	[20**]
3p21.3	<i>TREX1</i>	rs72556554, R114H and other 11 nonsynonymous substitutions	European	[16]
Xq28	<i>MECP2-IRAK1</i>	rs2269368, rs17435	European, Chinese, Korean, European-Amerindian (Mexican)	[18,19,20**]
3p14.3	<i>PXK</i>	rs6445975, rs2176082	European	[4**,20**]
2q24	<i>IFIH1</i>	rs1990760	European	[20**]
11p15.5	<i>KIAA1542 (PHRF1)</i>	rs4963128	European	[4**]
8p23.1	<i>XKR6</i>	rs6985109	European	[4**]
6q21	<i>ATG5-PRMD1</i>	rs6568431, rs2245214	European, Chinese	[4**,20**]
22q11.2	<i>UBE2L3</i>	rs5754217	European, Chinese	[2*,20**]
5q33.3	<i>PTTG1</i>	rs2431099	European	[2*,20**]
6p21	<i>UHRF1BP1</i>	rs11755393	European	[20**]
5q32	<i>TNIP1</i>	rs7708392	European, Chinese, Thai	[3*,20**]
7p15.2	<i>JAZF1</i>	rs849142	European	[20**]
7p21.3	<i>ICA1</i>	rs10156091	European	[4**,20**]
1q24	<i>IL10</i>	rs3024505	European	[20**]
1q25.3	<i>NMNAT2</i>	rs2022013	European, Chinese	[3*,4**]
11q23.3	<i>ETS1</i>	rs6590330	Chinese, Thai	[3*,7]
10q11.23	<i>WDFY4</i>	rs877819	Chinese, Thai	[3*,7]
7p12.2	<i>IKZF1</i>	rs4917014	Chinese	[3*]
12q24.32	<i>SLC15A4</i>	rs10847697, rs1385374	Chinese	[3*]
2p22.3	<i>RASGRP3</i>	rs13385731	Chinese	[3*]

CD226 on NKT cells [13*]. More recently *BANK1* and *BLK*, thought to be involved in B cell activation and tolerance [5**,6**], respectively have been now clearly established [3*,14,15].

One pathway that has been biologically and genetically strongly related to SLE pathogenesis is the type I interferon (IFN) pathway. Several genes for factors upstream and downstream of IFN production, such as *IRF5*, *STAT4*, and more recently *TNFAIP3*, *TYK2*, and *TREX1*, have been associated to susceptibility to SLE [2*,16,17*]. *IRAK1* has been an interesting candidate but it is closely linked to *MECP2*, a gene that can regulate expression of *IRAK1* also associated. It has proven as yet impossible to

discern the genetic effects between these X chromosome genes [18*,19*,20**]. *TREX1* is mainly represented by rare but penetrant and mutations leading to high levels of type I interferon found in few patients with lupus, suggesting a potentially important role of rare variants that have remained undetected with the use of common variation mapping.

Other recently identified loci, such as *PXK*, *XKR6*, and *KIAA1542* close to *IRF7* [4**], with no known function or correlation to SLE pathology, have the potential to lead to the discovery of novel pathways involved in SLE. It is unclear if the genetic association of *KIAA1542* indeed represents an association with *IRF7*.

Table 2

Susceptibility Genes for RA

Chromosome	Gene	SNPs	Population	Reference
6p21.32	<i>HLA-DRB1</i>	rs615672, rs660895, rs64576200, rs6910071, rs13192471	European, Japanese	[21–23,24**,25*]
1p13.2	<i>PTPN22</i>	rs6679677, rs2476601	European	[21–23,24**,30*]
2q32.3	<i>STAT4</i>	rs7574865	European, Japanese	[24**,25*]
9q34	<i>TRAF1-C5</i>	rs3761847, rs881375	European	[22,24**,30*]
6q23.3	<i>TNFAIP3, OLIG3</i>	rs10499194, rs6920220	European, Japanese	[23,24**,25*,29*]
6p21.32	<i>HLA-DQA1, HLA-DQA2</i>	rs6457617	European	[31]
18q23	<i>SALL3</i>	rs2002842	European	[31]
20q13.12	<i>CD40</i>	rs4810485	European	[23,24**]
9p13.3	<i>CCL21</i>	rs2812378	European	[23]
12q13.3	<i>KIF5A,PIP4K2C</i>	rs1678542	European	[23]
1p36.32	<i>TNFRSF14</i>	rs3890745	European	[23,24**]
10p15.1	<i>PRKCQ</i>	rs4750316	European	[23,24**]
7q21.2	<i>CDK6</i>	rs42041	European	[23]
2p16.1	<i>REL</i>	rs13017599, rs13031237	European	[24**,30*]
2q33.2	<i>CTLA4</i>	rs231735, rs3087243	European	[24**,30*]
8p23.1	<i>BLK</i>	rs2736340	European	[30*]
2q11.2	<i>AFF3</i>	rs11676922,rs10865035	European	[24**]
5q11.2	<i>ANKRD55,IL6ST</i>	rs6859219	European	[24**]
14q24.3	<i>BATF</i>	rs7155603	European	[24**]
5q21.1	<i>C5orf30</i>	rs26232	European	[24**]
9p13.3	<i>CCL21</i>	rs951005	European	[24**]
6q27	<i>CCR6</i>	rs3093023, rs3093024	Japanese	[24**,25*]
1q24.2	<i>CD247</i>	rs840016	European	[24**]
17q12	<i>IKZF3</i>	rs2872507	European	[24**]
4q27	<i>IL2,IL21</i>	rs13119723	European	[24**]
10p15.1	<i>IL2RA</i>	rs706778	European	[24**]
7q32.1	<i>IRF5</i>	rs10488631	European	[24**]
15q23	<i>KIF3</i>	rs17374222	European	[24**]
1p34.3	<i>POU3F1</i>	rs12131057	European	[24**]
3p14.3	<i>PXK</i>	rs13315591	European	[24**]
4p15.2	<i>RBPJ</i>	rs874040	European	[24**]
12q24.12	<i>SH2B3</i>	rs3184504	European	[24**]
2p14	<i>SPRED2</i>	rs934734	European	[24**]
21q22.3	<i>UBASH3A</i>	rs11203203	European	[24**]

Finally, a very large replication study identified and replicated several genes among which are *JAZF1*, *TNIP1*, *PRDM1* (or *BLIMP1*), *UHRF1BP1*, *PTTG1* [2*], *UBE2L3* [2*], *IL10*, *IL21*, and the *IL21R* [20**]. Importantly, several other genes were also confirmed in this study such as *ATG5*, *ICA1*, and *NMNAT2* found in a previous GWAS from the SLEGEN consortium [4**]. *ATG5* is an important component of the autophagy pathway. *JAZF1* and *UHRF1BP1* are transcription factors, while *TNIP1* interacts with *TNFAIP3*, its function in regulating *TNFAIP3* is not known. *TNFAIP3* (or A20) regulates inflammation by turning off NFκB through polyubiquitination and degradation.

Studies in Asian populations have identified new susceptibility genes for lupus and replication of the hitherto identified genes in Europeans has revealed the presence of some genes but not others [3*,7,14]. Two genes were clearly identified, *ETS1* and *WDFY4* [3*,7]. *ETS1* is involved in the development of TH17 cells while *WDFY4* codes for a protein of unknown function. *STAT4*, *IRF5*, *BANK1*, *BLK*, *TNFAIP3*, and *TNFSF4* have been

confirmed in Asians [3*,7]. Studies on African Americans and European-Amerindian admixed populations are ongoing.

Genetics of RA

Multiple GWAS have corroborated the MHC genes as major genetic contributors to the risk of developing RA [21–23,24**,25*]. Within the MHC, the strongest contribution to risk is given by the *HLA-DRB1* gene, which codes for the third hypervariable region of the HLA-DR molecule β chain. Since the original report of Peter Stastny, diverse *HLA-DRB1* alleles have been associated in European, Asian, African, and European-Amerindian populations [26]. All the associated alleles were unified by Peter Gregersen under the hypothesis of the shared epitope (SE). SE alleles are associated with anti-citrulline antibody production, a major biomarker for RA and determine severity. A study analyzing over 1400 SNPs within the HLA regions from the IMAGEN Consortium found the peak association for RA between the gene *BTNL2* and *HLA-DRA* (SNP rs2395175) and the allele for the *DQA1* gene *DQA1*0301* [9]. Dense typing of the

MHC have revealed several *DRB1*-independent associations, including a signal at *MICA*, one in the border between class I and class III region and some in the class I region [27].

Although any other association outside the MHC is rather modest in RA, and the SE accounts for 18–37% of the genetic heritability [28], fine mapping of candidate non-MHC linkage regions successfully identified important susceptibility genes such as *PTPN22* (1p13), *PADI4* (1p36), and *STAT4* (2q32). To date, seven GWAS conducted in collections of thousands of patients with RA and healthy controls reliably detect several new susceptibility genes [21–23,25*,29*,30*,31] (Table 2). The loci supported by the best evidence are the *TRAF1-C5* [22,24**,30*] and the 6q23 region [23,24**,25,29*], both with strongest effect in anti-CCP+ patients. GWAS studies have also confirmed the association of *PTPN22* and *STAT4*, as well as previously reported candidate genes identified by studies that did not have enough power such as *CTLA4* and *CD40*. The maximum power has been achieved by a recent meta-analysis of GWAS conducted in a total of 12,307 patients and 28,975 controls of European ancestry [24**]. This allowed the identification and confirmation of *IL6ST*, *SPRED2*, *RBPJ*, *CCR6* [25*], *IRF5*, and *PXK* [24**]. While *SPRED2* has been found to be a negative regulator of the Ras-ERK cascade, *IL6ST* and *CCR6* are inflammation regulators and *RBPJ* is a transcription factor important in dendritic cell function.

It is important to note that major differences across racial groups have been noticed; for example, *PTPN22* and *CTLA4* were associated in Europeans whereas *PADI4* and *SLC22A4* are confirmed only in Asian population groups. By contrast, the *STAT4* association is valid for European, Asian, and European-Amerindian but not African Americans [32]. All these findings add evidence of the complexity and the heterogeneity of the genetic basis of

RA and justify the study of diverse populations. One GWAS has been performed in Japanese [25*] but none in admixed populations of European-Amerindian or African American. The observed heterogeneity highlights the importance of conducting well-powered GWAS in non-European populations in order to dissect all the genetic contribution to the disease.

Genetics of systemic sclerosis

Systemic sclerosis (SSc) or scleroderma is an autoimmune disease characterized by an extensive fibrotic process that affects multiple organs and tissues. Until now genetic studies have not been particularly successful in the identification of risk factors for SSc. The controversial results found for the majority of genes, such as, *PTPN22*, *CTGF*, or *TGF- β* , suggest that those studies were often limited by small sample size and clinical heterogeneity. Some were finally replicated when large sample sizes were used (Table 3). One of the first discoveries was the contribution, as in other autoimmune diseases, of HLA class II genes, which seem to be predominantly associated with the presence of specific autoantibodies rather than with SSc itself. These associations have been confirmed in a recent GWAS in SSc in the Korean population [33]. The first GWAS in a sample of European ancestry including 2296 SSc patients and 5171 controls has firmly established the role of the *HLA* genes in SSc [34**]. *STAT4* and *IRF5* that had previously been identified as risk factors for SSc through candidate gene studies [35,36], were also identified in the GWAS of European patients with SSc [34**]. The consistent association of these genes with SSc susceptibility provides compelling evidence that variation in genes with key functions in the innate immune system are involved in the pathogenesis of the disease.

The identification of the association of connective tissue growth factor (*CTGF*) gene with risk to SSc provides one of the most conflicting results to date. The potential

Table 3

Susceptibility Genes for SSc

Chromosome	Gene	SNPs	Population	Reference
6p21.32	<i>HLA-DPB1,DPB2</i>	rs3128930, rs7764491, rs7763822, rs3128965, rs3117230, rs7763822, rs7764491, rs3117230,rs 3128965	Korean, European	[33,34]
2q32.3	<i>HLA-DQB1</i> <i>STAT4</i>	rs6457617 rs7574865, rs11889341, rs8179673, rs10181656, rs3821236	European European, Japanese	[34] [34,44]
7q32	<i>TNPO3-IRF5</i>	rs2004640, rs2280714, rs10954213, rs10488631, rs12537284, rs4728142	European, Japanese	[34–36]
4q24	<i>BANK1</i>	rs10516487, rs17266594	European	[41]
8p23.1	<i>C8orf13-BLK</i>	rs2736340, rs13277113	European	[42,43]
17q21.32	<i>TBX21</i>	rs11650354, rs17699436	European	[44]
1q25.1	<i>TNFSF4</i>	rs1234314, rs2205960, rs844644, rs844648	European	[47]
10q24	<i>FAS</i>	rs1800682 (G-670A)	European	[45,46]
6q23.1	<i>CTGF</i>	rs6918698 (G-945C), rs9399005	European, Japanese	[37–40]
1q22–23	<i>CD247</i>	rs2056626	European	[34]

functional -945 G allele of *CTGF* gene was primarily associated with susceptibility to SSc in Europeans and Japanese [37]. However, three additional studies in Europeans failed to replicate the association [38–40]. A recent study suggests that variant rs9399005 in the 3'UTR region of the *CTGF* gene is associated with both subtypes of SSc [39]. The lack of replication of the *CTGF* polymorphisms in three large cohorts of SSc suggests that *CTGF* may not be a strong genetic determinant for SSc susceptibility.

Genes involved in B cell receptor signaling contribute to SSc susceptibility. Association with diffuse SSc (dcSSc) has been identified with *BANK1* [41]. The *C8orf13-BLK* region has been associated in European and Japanese patients with SSc [42,43]. These findings suggest an important role of B cells in the pathogenesis of SSc.

A large multicenter study in Europeans found two polymorphisms (rs11650354 and rs17699436) in the *TBX21* gene associated with SSc [44]. In addition, they also showed a gene–gene interaction between the *TBX21* and *STAT4*.


Some other genes that appear to play a role in susceptibility to SSc are *FAS*, *TNFSF4*, and *CD247*. The promoter rs1800682 polymorphism in the *FAS* gene is a confirmed susceptibility variant for SSc in different populations

[45,46]. In addition, the SLE susceptibility gene *TNFSF4* [11,12,14] was associated with SSc in a large case–control study [47]. The *CD247* gene, which encodes the T cell receptor zeta (CD3 ζ) subunit, was a new susceptibility gene for SSc in a GWAS and this association has been confirmed in an independent cohort [34^{••}].

Conclusions

The genetics results deriving from GWAS are just the beginning of a new era of research, but new insight on disease genetics has been acquired. First, the risk alleles identified for these diseases explain only between 5 and 15% of the whole genetic contribution to disease and with odds ratios ranging from 1.01 to 2.4 at the most. Clearly, gene–gene interactions, gene–environment interactions, and other genomic structural variation such as copy number variation, the role of rare variants, and epigenetics need to be addressed. In this regard, the recent identification of rare, but highly penetrant mutations in the sialic acid acetyltransferase gene (*SIAE*) involved in SLE and RA susceptibility and with an important functional impact in the gene [48^{••}] suggests that rare variants may have an important role not yet fully comprehended. Second, most autoimmune diseases, in particular systemic autoimmune diseases, share several susceptibility genes. The differences seem to reside on the contribution of each gene in each disease. HLA alleles have been known to be differ-

Figure 1



Unique and shared genes between SLE, RA and SSc.

ent from disease to disease, but the risk alleles for non-MHC genes appear, until now to be the same. However while *IRF5* and *STAT4* are prominent genes in SLE, *PTPN22* and *TNFAIP3* are major genes in RA and *CD247* in SSc. Very few genes are unique for each disease (Figure 1) and whether some genes may predominate in individuals with certain clinical manifestations is still a difficult nut to crack. Phenotyping of samples studied in genetics lack the detail required to define the correlation between clinical manifestations and genes. One possible exception is *ITGAM* in SLE, a unique gene for lupus importantly associated with kidney disease. Third, studies in different populations are important. Differences and similarities will not only lead to a comprehensive picture of genetic susceptibility, but may also pave the way to the very needed studies on gene–environmental interactions in autoimmune disease, a theme about which we know practically nothing.

Genetics studies are a starting point to cell biology and immunology studies aimed at understanding disease pathogenesis and the influence of susceptibility genes on cell function. Indeed, we have challenging and exciting times ahead of us.

Acknowledgements

This work has been supported by the NIH grants AI083194, CA141700, and AR058621, the Instituto de Salud Carlos III (FIS), and the Consejería de Salud de Andalucía to MEAR. Funding from the Swedish Research Council, the Swedish Association Against Rheumatism, the King Gustaf Vth-80th Jubilee Fund, and the Royal Swedish Academy of Sciences to MEAR for work supported in Sweden are greatly and sincerely acknowledged.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Wong M, Tsao BP: **Current topics in human SLE genetics.** *Springer Semin Immunopathol* 2006, **28**:97-107.
2. Graham RR, Cotsapas C, Davies L, Hackett R, Lessard CJ, Leon JM, Burt NP, Guiducci C, Parkin M, Gates C et al.: **Genetic variants near *TNFAIP3* on 6q23 are associated with systemic lupus erythematosus.** *Nat Genet* 2008, **40**:1059-1061.
3. Han JW, Zheng HF, Cui Y, Sun LD, Ye DQ, Hu Z, Xu JH, Cai ZM, Huang W, Zhao GP et al.: **Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus.** *Nat Genet* 2009, **41**:1234-1237.
4. Harley JB, Alarcon-Riquelme ME, Criswell LA, Jacob CO, Kimberly RP, Moser KL, Tsao BP, Vyse TJ, Langefeld CD, Nath SK et al.: **Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in *ITGAM*, *PXK*, *KIAA1542* and other loci.** *Nat Genet* 2008, **40**:204-210.
- This study is one of the two first GWAS for SLE. Primarily *ITGAM* was identified as a major gene for lupus.
5. Hom G, Graham RR, Modrek B, Taylor KE, Ortmann W, Garnier S, Lee AT, Chung SA, Ferreira RC, Pant PV et al.: **Association of systemic lupus erythematosus with *C8orf13*-BLK and *ITGAM*-*ITGAX*.** *N Engl J Med* 2008, **358**:900-909.
- The other first GWAS in SLE. The risk polymorphisms close to the promoter of BLK and 3' of *C8orf13* (now *FAM146A*) correlate with expression levels of both genes.
6. Kozyrev SV, Abelson AK, Wojcik J, Zaghlool A, Linga Reddy MV, Sanchez E, Gunnarsson I, Svenungsson E, Sturfelt G, Jonsen A et al.: **Functional variants in the B-cell gene *BANK1* are associated with systemic lupus erythematosus.** *Nat Genet* 2008, **40**:211-216.
- This study identified *BANK1* as a gene for lupus and also defined the functional variants involved in the risk for lupus. Two variants were identified, one leading to differences in isoform expression and a coding variant R61H substitution located in the second exon. The short isoform lacks the second exon and its expression correlated with protection.
7. Yang W, Shen N, Ye DQ, Liu Q, Zhang Y, Qian XX, Hirankarn N, Ying D, Pan HF, Mok CC et al.: **Genome-wide association study in Asian populations identifies variants in *ETS1* and *WDFY4* associated with systemic lupus erythematosus.** *PLoS Genet* 2010, **6**:e1000841.
8. Fernando MM, Stevens CR, Sabeti PC, Walsh EC, McWhinnie AJ, Shah A, Green T, Rioux JD, Vyse TJ: **Identification of two independent risk factors for lupus within the MHC in United Kingdom families.** *PLoS Genet* 2007, **3**:e192.
9. Rioux JD, Goyette P, Vyse TJ, Hammarstrom L, Fernando MM, Green T, De Jager PL, Foisy S, Wang J, de Bakker PI et al.: **Mapping of multiple susceptibility variants within the MHC region for 7 immune-mediated diseases.** *Proc Natl Acad Sci USA* 2009, **106**:18680-18685.
10. Nath SK, Han S, Kim-Howard X, Kelly JA, Viswanathan P, Gilkeson GS, Chen W, Zhu C, McEver RP, Kimberly RP et al.: **A nonsynonymous functional variant in integrin- α (M) (encoded by *ITGAM*) is associated with systemic lupus erythematosus.** *Nat Genet* 2008, **40**:152-154.
- This study identified a potential functional variant for *ITGAM*. In particular a coding R-H substitution is defined as the risk variants for SLE. The substitution is not conserved however and does not reside in a binding region for *ITGAM* (also known as *CD11b* or *CR3*).
11. Delgado-Vega AM, Abelson AK, Sanchez E, Witte T, D'Alfonso S, Galeazzi M, Jimenez-Alonso J, Pons-Estel BA, Martin J, Alarcon-Riquelme ME: **Replication of the *TNFSF4* (*OX40L*) promoter region association with systemic lupus erythematosus.** *Genes Immun* 2009, **10**:248-253.
12. Cunningham-Graham DS, Graham RR, Manku H, Wong AK, Whittaker JC, Gaffney PM, Moser KL, Rioux JD, Altschuler D, Behrens TW et al.: **Polymorphism at the TNF superfamily gene *TNFSF4* confers susceptibility to systemic lupus erythematosus.** *Nat Genet* 2008, **40**:83-89.
13. Lofgren SE, Delgado-Vega AM, Gallant CJ, Sanchez E, Frostegard J, Truedsson L, Garrido ED, Sabio JM, Gonzalez-Escribano MF, Pons-Estel BA et al.: **A 3'-untranslated region variant is associated with impaired expression of *CD226* in T and natural killer T cells and is associated with susceptibility to systemic lupus Erythematosus.** *Arthritis Rheum* 2010, **62**:3404-3414.
14. Chang YK, Yang W, Zhao M, Mok CC, Chan TM, Wong RW, Lee KW, Mok MY, Wong SN, Ng IO et al.: **Association of *BANK1* and *TNFSF4* with systemic lupus erythematosus in Hong Kong Chinese.** *Genes Immun* 2009, **10**:414-420.
15. Ito I, Kawasaki A, Ito S, Hayashi T, Goto D, Matsumoto I, Tsutsumi A, Hom G, Graham RR, Takasaki Y et al.: **Replication of the association between the *C8orf13*-BLK region and systemic lupus erythematosus in a Japanese population.** *Arthritis Rheum* 2009, **60**:553-558.
16. Lee-Kirsch MA, Gong M, Chowdhury D, Senenko L, Engel K, Lee YA, de Silva U, Bailey SL, Witte T, Vyse TJ et al.: **Mutations in the gene encoding the 3'-5' DNA exonuclease *TREX1* are associated with systemic lupus erythematosus.** *Nat Genet* 2007, **39**:1065-1067.
17. Musone SL, Taylor KE, Lu TT, Nititham J, Ferreira RC, Ortmann W, Shifrin N, Petri MA, Kamboh MI, Manzi S et al.: **Multiple polymorphisms in the *TNFAIP3* region are independently associated with systemic lupus erythematosus.** *Nat Genet* 2008, **40**:1062-1064.

18. Sawalha AH, Webb R, Han S, Kelly JA, Kaufman KM, Kimberly RP, Alarcon-Riquelme ME, James JA, Vyse TJ, Gilkeson GS *et al.*: **Common variants within MECP2 confer risk of systemic lupus erythematosus.** *PLoS ONE* 2008, **3**:e1727.
 19. Jacob CO, Zhu J, Armstrong DL, Yan M, Han J, Zhou XJ, Thomas JA, Reiff A, Myones BL, Ojwang JO *et al.*: **Identification of IRAK1 as a risk gene with critical role in the pathogenesis of systemic lupus erythematosus.** *Proc Natl Acad Sci USA* 2009, **106**:6256-6261.
 20. Gateva V, Sandling JK, Hom G, Taylor KE, Chung SA, Sun X, Ortmann W, Kosoy R, Ferreira RC, Nordmark G *et al.*: **A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus.** *Nat Genet* 2009, **41**:1228-1233.
- A second GWAS and replication study identifying novel lupus susceptibility genes in Europeans with a large replication cohort. This study shows how large case control sets lead to identification of new susceptibility genes.
21. WTCCC: **Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls.** *Nature* 2007, **447**:661-678.
 22. Plenge RM, Seielstad M, Padyukov L, Lee AT, Remmers EF, Ding B, Liew A, Khalili H, Chandrasekaran A, Davies LR *et al.*: **TRAF1-C5 as a risk locus for rheumatoid arthritis—a genome-wide study.** *N Engl J Med* 2007, **357**:1199-1209.
 23. Raychaudhuri S, Remmers EF, Lee AT, Hackett R, Guiducci C, Burtt NP, Gianniny L, Korman BD, Padyukov L, Kurreeman FA *et al.*: **Common variants at CD40 and other loci confer risk of rheumatoid arthritis.** *Nat Genet* 2008, **40**:1216-1223.
 24. Stahl EA, Raychaudhuri S, Remmers EF, Xie G, Eyre S, Thomson BP, Li Y, Kurreeman FA, Zernakova A, Hinks A *et al.*: **Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci.** *Nat Genet* 2010, **42**:508-514.
- Study showing how the inclusion of large population sets leads to the identification of a large number of new disease loci.
25. Kochi Y, Okada Y, Suzuki A, Ikari K, Terao C, Takahashi A, Yamazaki K, Hosono N, Myouzen K, Tsunoda T *et al.*: **A regulatory variant in CCR6 is associated with rheumatoid arthritis susceptibility.** *Nat Genet* 2010, **42**:515-519.
 26. Kochi Y, Suzuki A, Yamada R, Yamamoto K: **Genetics of rheumatoid arthritis: underlying evidence of ethnic differences.** *J Autoimmun* 2009, **32**:158-162.
 27. Lee HS, Lee AT, Criswell LA, Seldin MF, Amos CI, Carulli JP, Navarrete C, Remmers EF, Kastner DL, Plenge RM *et al.*: **Several regions in the major histocompatibility complex confer risk for anti-CCP-antibody positive rheumatoid arthritis, independent of the DRB1 locus.** *Mol Med* 2008, **14**:293-300.
 28. van der Woude D, Houwing-Duistermaat JJ, Toes RE, Huizinga TW, Thomson W, Worthington J, van der Helm-van Mil AH, de Vries RR: **Quantitative heritability of anti-citrullinated protein antibody-positive and anti-citrullinated protein antibody-negative rheumatoid arthritis.** *Arthritis Rheum* 2009, **60**:916-923.
 29. Plenge RM, Cotsapas C, Davies L, Price AL, de Bakker PI, Maller J, Pe'er I, Burtt NP, Blumenstiel B, DeFelice M *et al.*: **Two independent alleles at 6q23 associated with risk of rheumatoid arthritis.** *Nat Genet* 2007, **39**:1477-1482.
 30. Gregersen PK, Amos CI, Lee AT, Lu Y, Remmers EF, Kastner DL, Seldin MF, Criswell LA, Plenge RM, Holers VM *et al.*: **REL, encoding a member of the NF-kappaB family of transcription factors, is a newly defined risk locus for rheumatoid arthritis.** *Nat Genet* 2009, **41**:820-823.
 31. Julia A, Ballina J, Canete JD, Balsa A, Tornero-Molina J, Naranjo A, Alperi-Lopez M, Erra A, Pascual-Salcedo D, Barcelo P *et al.*: **Genome-wide association study of rheumatoid arthritis in the Spanish population: KLF12 as a risk locus for rheumatoid arthritis susceptibility.** *Arthritis Rheum* 2008, **58**:2275-2286.
 32. Kelley JM, Hughes LB, Malik A, Danila MI, Edberg Y, Alarcon GS, Conn DL, Jonas BL, Callahan LF, Smith EA *et al.*: **Genetic variants of STAT4 associated with rheumatoid arthritis in persons of Asian and European ancestry do not replicate in African Americans.** *Ann Rheum Dis* 2010, **69**:625-626.
 33. Zhou X, Lee JE, Arnett FC, Xiong M, Park MY, Yoo YK, Shin ES, Reveille JD, Mayes MD, Kim JH *et al.*: **HLA-DPB1 and DPB2 are genetic loci for systemic sclerosis: a genome-wide association study in Koreans with replication in North Americans.** *Arthritis Rheum* 2009, **60**:3807-3814.
 34. Radstake TR, Gorlova O, Rueda B, Martin JE, Alizadeh BZ, Palomino-Morales R, Coenen MJ, Vonk MC, Voskuyl AE, Schuerwegh AJ *et al.*: **Genome-wide association study of systemic sclerosis identifies CD247 as a new susceptibility locus.** *Nat Genet* 2010, **42**:426-429.
- The first GWAS for SSc. Two large replication sets were used. Only one gene, CD247 (CD3z) was identified.
35. Dieude P, Guedj M, Wipff J, Avouac J, Fajardy I, Diot E, Granel B, Sibilia J, Cabane J, Mouthon L *et al.*: **Association between the IRF5 rs2004640 functional polymorphism and systemic sclerosis.** *Arthritis Rheum* 2009, **60**:225-233.
 36. Ito I, Kawaguchi Y, Kawasaki A, Hasegawa M, Ohashi J, Hikami K, Kawamoto M, Fujimoto M, Takehara K, Sato S *et al.*: **Association of a functional polymorphism in the IRF5 region with systemic sclerosis in a Japanese population.** *Arthritis Rheum* 2009, **60**:1845-1850.
 37. Kawaguchi Y, Ota Y, Kawamoto M, Ito I, Tsuchiya N, Sugiyama T, Katsumata Y, Soejima M, Sato S, Hasegawa M *et al.*: **Association study of a polymorphism of the CTGF gene and susceptibility to systemic sclerosis in the Japanese population.** *Ann Rheum Dis* 2009, **68**:1921-1924.
 38. Rueda B, Simeon C, Hesselstrand R, Herrick A, Worthington J, Ortego-Centeno N, Riemekasten G, Fonollosa V, Vonk MC, van den Hoogen FH *et al.*: **A large multicentre analysis of CTGF -945 promoter polymorphism does not confirm association with systemic sclerosis susceptibility or phenotype.** *Ann Rheum Dis* 2009, **68**:1618-1620.
 39. Granel B, Argiro L, Hachulla E, Fajardy I, Weiller PJ, Durand JM, Frances Y, Dombey AM, Marquet S, Lesavre N *et al.*: **Association between a CTGF gene polymorphism and systemic sclerosis in a French population.** *J Rheumatol* 2010, **37**:351-358.
 40. Gourh P, Mayes MD, Arnett FC: **CTGF polymorphism associated with systemic sclerosis.** *N Engl J Med* 2008, **358**:308-309 author reply 309.
 41. Rueda B, Gourh P, Broen J, Agarwal SK, Simeon C, Ortego-Centeno N, Vonk MC, Coenen M, Riemekasten G, Hunzelmann N *et al.*: **BANK1 functional variants are associated with susceptibility to diffuse systemic sclerosis in Caucasians.** *Ann Rheum Dis* 2010, **69**:700-705.
 42. Ito I, Kawaguchi Y, Kawasaki A, Hasegawa M, Ohashi J, Kawamoto M, Fujimoto M, Takehara K, Sato S, Hara M *et al.*: **Association of the FAM167A-BLK region with systemic sclerosis.** *Arthritis Rheum* 2010, **62**:890-895.
 43. Gourh P, Agarwal SK, Martin E, Divecha D, Rueda B, Bunting H, Assassi S, Paz G, Shete S, McNearney T *et al.*: **Association of the C8orf13-BLK region with systemic sclerosis in North-American and European populations.** *J Autoimmun* 2010, **34**:155-162.
 44. Gourh P, Agarwal SK, Divecha D, Assassi S, Paz G, Arora-Singh RK, Reveille JD, Shete S, Mayes MD, Arnett FC *et al.*: **Polymorphisms in TBX21 and STAT4 increase the risk of systemic sclerosis: evidence of possible gene-gene interaction and alterations in Th1/Th2 cytokines.** *Arthritis Rheum* 2009, **60**:3794-3806.
 45. Liakouli V, Manetti M, Pacini A, Tolusso B, Fatini C, Toscano A, Cipriani P, Guiducci S, Bazzichi L, Codullo V *et al.*: **The -670G>A polymorphism in the FAS gene promoter region influences the susceptibility to systemic sclerosis.** *Ann Rheum Dis* 2009, **68**:584-590.
 46. Broen J, Gourh P, Rueda B, Coenen M, Mayes M, Martin J, Arnett FC, Radstake TR: **The FAS -670A>G polymorphism influences susceptibility to systemic sclerosis phenotypes.** *Arthritis Rheum* 2009, **60**:3815-3820.
 47. Gourh P, Arnett FC, Tan FK, Assassi S, Divecha D, Paz G, McNearney T, Draeger H, Reveille JD, Mayes MD *et al.*: **Association of TNFSF4 (OX40L) polymorphisms with**

susceptibility to systemic sclerosis. *Ann Rheum Dis* 2010, **69**:550-555.

48. Surolia I, Pirnie SP, Chellappa V, Taylor KN, Cariappa A, Moya J, Liu H, Bell DW, Driscoll DR, Diederichs S *et al.*: **Functionally defective germline variants of sialic acid acetyltransferase in autoimmunity.** *Nature* 2010, **466**:243-247.

The discovery of the gene SIAE, an enzyme that negatively controls B cell signaling, as related to autoimmunity. Eight homozygous mutations are found in patients with rheumatoid arthritis and lupus.

49. Lee YH, Woo JH, Choi SJ, Ji JD, Song GG: **Association of programmed cell death 1 polymorphisms and systemic lupus erythematosus: a meta-analysis.** *Lupus* 2009, **18**:9-15.