

Distinct roles of IL-22 in human psoriasis and inflammatory bowel disease

Wenjun Ouyang

Department of Immunology, Genentech, Inc., 1 DNA Way, M/S 34, South San Francisco, CA 94080, USA

ARTICLE INFO

Article history:

Available online 23 November 2010

Keywords:

IL-22

IBD

Psoriasis

Th17

ABSTRACT

IL-22, an IL-10 family cytokine, is produced by different leukocyte subsets, including T cells, NK cells and lymphoid tissue inducer (LTi) cells. IL-22 mediates the crosstalk between leukocytes and tissue epithelia because its receptor is preferentially expressed on various tissue epithelial cells. IL-22 is essential for host defense against infections of extracellular pathogens, such as bacteria and yeasts, by eliciting various innate defensive mechanisms from tissue epithelial cells and promoting wound-healing responses. In autoimmune diseases, however, diverse tissue microenvironments and underlying pathogenic mechanisms may result in opposing contributions of IL-22 in disease progression. For example, in psoriasis, IL-22 can synergize with other proinflammatory cytokines to induce many of the pathogenic phenotypes from keratinocytes and exacerbate disease progression. In contrast, IL-22 plays a beneficial role in IBD by enhancing barrier integrity and epithelial innate immunity of intestinal tract.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

1.1. IL-22 and IL-22 receptor

IL-22 was originally cloned by Renauld and colleagues from murine lymphoma cell line treated with IL-9 [1]. Because it shares sequence homology with IL-10, they named this cytokine as IL-10 related T cell-derived inducible factor (IL-TIF) [1], which was subsequently renamed as IL-22 [2]. Human IL-22 cDNA encodes a protein with 179 amino acids and share 23% sequence homology with human IL-10 gene [2]. Interestingly, structure analyses indicate that IL-22 does not form domain swapped dimer as that of IL-10 [3]. But its structure displays the characteristic α -helical fold consisting of six helices labeled A to F and connecting loops, which is shared by all Class II cytokines [4,5]. Despite the structural differences between IL-22 and IL-10, these two cytokines share a common receptor subunit, IL-10R2 chain [2]. In addition, IL-22 and IL-10 require their own unique receptors, IL-22R1 and IL10R1, respectively, together with IL-10R2, to deliver the downstream signaling. IL-22R1, like IL-10R1 and IL-10R2, belongs to Class II cytokine receptor family [6,7].

Upon binding to the heterodimeric receptor complex, IL-22R1 and IL-10R2, IL-22 activates the Jak1 and Tyk2 [8], which further phosphorylate downstream Stat molecules, Stat1, Stat3, and Stat5 [2,8]. Similar to other IL-10 family cytokines, IL-22 predominately utilizes Stat3 to mediate its functions. Interestingly, there is a unique domain located at the c-terminal of IL-22R1 intracellular tail, which can recruit Stat3, through the interaction of the coiled-

coil domain of Stat3, to the receptor independent of 8 tyrosine residues in the receptor [9]. The biological importance of this novel mechanism of Stat3 recruitment is still unclear. In addition to IL-22R1, there is also a decoy receptor, referred as IL-22BP or IL-22R2, which is structurally homologous to the extracellular domain of IL-22R1 but without the transmembrane and intracellular domains. IL-22BP binds IL-22 with high affinity and can block the binding of IL-22 to IL-22R1 [10–12].

1.2. Cellular sources and targets of IL-22

The early work by Renauld and colleagues has suggested that IL-22 is a cytokine produced by T cells [1]. They showed that IL-22 is detected from IL-9 treated T helper cell line, TS2, and from concanavalin A (ConA) stimulated mouse splenocytes, presumably mainly consisting of activated T cells. These results further confirmed by Wolk and colleagues by demonstrating that activated mouse T cells, especially Th1 cells, preferentially express IL-22 [13]. In 2005, a novel T helper subset, Th17, was identified [14,15]. These cells preferentially express IL-17 (or IL-17A), a signature cytokine distinguishing these cells from other T helper subsets, including Th1, Th2 and Treg cells. Th17 cells play essential roles in host defense against infections by various pathogens, especially extracellular bacteria, and participate in the pathogenesis of many autoimmune diseases [16]. It was soon discovered that IL-22 is another cytokine preferentially expressed by Th17 cells [17–19]. The level of IL-22 produced by Th17 cells is much higher than that produced by Th1 cells or undifferentiated Th0 cells. But the regulation of IL-22 and IL-17 produced in T cells is not parallel. IL-6 and TGF- β are both required for IL-17 induction from naïve T cells, whereas IL-6 alone is sufficient to promote the expression of IL-22

E-mail address: ouyang@gene.com.

[18,20,21]. In fact, TGF- β inhibits IL-22 production in a dose dependent manner [18]. In addition to CD4 T cells, CD8 T cells, $\gamma\delta$ T cells, and NKT cells also express high level of IL-22 upon activation, especially in conjunction with IL-23 treatment [18,22]. NK cells treated with IL-2 and IL-12 also express IL-22 [13]. Recently, lymphoid tissue induce (LTI) cells and developmental related NK like cells (NK22), which express NK marker NKp46, were demonstrated to be the major innate sources of IL-22, especially in the intestine [23–27]. Finally, subsets of myeloid cells express IL-23R and response to IL-23 treatment to produce lower level of IL-22 [18,28–30].

The IL-22 expressing human leukocytes are similar to those in mouse. Human T cells activated by anti-CD3 or ConA produce IL-22 [2]. Human Th17 cells generated in vitro or purified ex vivo from blood, based on lineage marker CCR6 and CCR4, preferentially express IL-22 [31,32]. CD161 is another surface marker that defines IL-17 and IL-22 expressing cells in human PBMC [33]. In addition to Th17 cells, CD161+ human CD8+ cells can also produce IL-22 as well as IL-17 [34]. Recently, a unique IL-22 producing CD4+ T helper subset, which express neither IL-17 nor IFN- γ , has been identified in human blood [35–37]. These cells express CCR10 and preferentially home to skin. Similar to what has been reported in the mouse system, these IL-22 producing human T cells can be generated from naïve CD4+ T cells in the presence of IL-6, but not TGF- β [18,35]. Interestingly, human Langerhans cells are able to differentiate T cells into the IL-22 only producing T helper phenotype in vitro [38]. Finally, innate human immune cell types such as NK cells and LTI cells are also produce IL-22 [39,40]. Human immature NK cells defined as CD161+CD117+CD34–CD94– express aryl hydrocarbon receptor (AHR) and IL-22 [41]. The murine equivalent NKp46+ NK like cells are developmentally linked to lymphoid tissue inducer cells [23,42,43]. These cells produce high level of IL-22 and may be the main sources of IL-22 in mucosal immunity.

Contrary to its production from leukocytes, IL-22 targets mainly tissue epithelial cells but not immune cells [44]. While IL-10R2 is ubiquitously expressed, IL-22R1 is restrictedly detected on cells with epithelial origins. Upon binding to its receptors on epithelial cells, IL-22 promotes cell proliferation and differentiation, and induces genes involved in host defense and wound-healing responses [16,45]. These cellular functions of IL-22 endow its essential role in epithelial defense mechanisms, especially against extracellular bacteria. Indeed, studies in preclinical models have demonstrated indispensable functions of IL-22 in mucosal immune responses to Gram-bacteria such as *Klebsiella pneumonia* and *Citrobacter rodentium* [28,46]. In addition, IL-22 is involved in the development of many human autoimmune diseases [16]. IL-22 expression is upregulated in autoimmune diseases, such as rheumatoid arthritis (RA), inflammatory bowel disease (IBD), and psoriasis. However, in autoimmune diseases, IL-22 can exert both protective and pathogenic functions. This review discusses the potential mechanisms underlying the distinct functions of IL-22 in psoriasis and IBD.

2. Pathogenic functions of IL-22 in psoriasis

Psoriasis is a common chronic inflammatory skin disease [47,48]. Psoriasis vulgaris is the predominant form of psoriasis, featured as erythematous and raised plaque scattered with silvery scales. Histologically, psoriatic skin is manifested as thickening of epidermis (acanthosis) with elongation of rete ridges. Keratinocytes in epidermis are hyperproliferative and associated with pre-maturation and incomplete cornification in the stratum corneum (parakeratosis). In addition, there is vascular elongation and dilation as well as significant leukocytes infiltration in both the dermis and epidermis of the skin lesions. Although, historically psoriasis has been primarily considered as keratinocytes disorder,

evidence from clinic, human genetics, and preclinical models has strongly supported a central role of the immune system in the pathogenesis of psoriasis [48,49]. It has been proposed that various cytokines from infiltrating leukocytes orchestrate the development of psoriasis.

What molecular links mediate the crosstalks between the infiltration of leukocytes and the abnormalities of keratinocytes in psoriatic skin is one of the important questions underlying the pathogenesis of psoriasis. Progress in the past several years has identified IL-22 as one of the key mediators that regulates many of the important pathogenic features of epidermis in psoriatic skin [50]. As discussed in detail later, IL-22 is augmented in psoriatic skin and targets epidermal keratinocytes that express IL-22 receptor complex. The expression of IL-22 from leukocytes in psoriatic skin is directly regulated by IL-23, whose indispensable role in psoriasis has been proven by clinic data with therapeutics blocking IL-23 pathway and by genome-wide genetic association data [51,52]. Finally, IL-22 induces hyperplasia, abnormal differentiation, and the expression of many psoriatic marker genes such as psoriasin (S100A7) from keratinocytes [53,54].

Elevated expression of IL-22 mRNA has been found in lesional skin [44]. Consistently, there is also increased IL-22 protein detected in the serum samples from psoriasis patients [55]. Recently, studies have indicated that T cells might be the major sources of IL-22 in psoriasis [56,57]. Leukocytes infiltrating the psoriatic skin consist of T cells, dendritic cells (DCs), and macrophages in the dermis as well as neutrophils and some T cells, mainly CD8+ T cells, in the epidermis [47,48]. Although myeloid cells and neutrophils play essential roles in the pathophysiology in psoriasis, several lines of evidence support T cells as central players during the progression of the diseases [49]. First, the only consistently identified genetic locus, PSORS1, is the MHC locus residing on chromosome 6p21, although the potential antigens in psoriasis triggering T cells activation in skin lesions are still unclear [47]. Activated CD4+ and CD8+ T cells have been identified in psoriatic plaques and blood of psoriatic patients [58]. Second, in models using xenograft transplantation where uninjured skin from psoriatic patients are grafted to severe combined immunodeficient (SCID) mice, injection of T cells from the donors into these mice is required for the development of psoriatic lesions. More specially, CD4+ T cell, but not CD8+ T cells are necessary and sufficient to induce the psoriatic phenotypes of the skin grafts. Finally, therapeutics that specifically target T cells, such as cyclosporine, tacrolimus, or antibodies targeting CD3 or CD4, resulted in clinical improvement of psoriasis, at least in some patients [59].

It has been proposed that psoriasis is a 'type 1' disease since both CD4+ and CD8+ T cells in psoriatic skin demonstrated IFN- γ expressing Th1 and Tc1 phenotypes, respectively [58]. The discovery of the IL-17 producing Th17 subset prompted people to investigate the association of Th17 cells with psoriasis [60]. Indeed, Th17 cells and downstream cytokine, IL-17, are also elevated in the lesional psoriatic skin [57,61]. As we discussed above, both Th1 and Th17 cells are potential sources of IL-22. CD8 cells from inflamed psoriatic skin also produce IL-22, in addition to other Th1/Th17 cytokines such as IFN- γ , TNF- α , IL-17, and IL-21 [62]. Recently, IL-22 but not IL-17 producing T helper cells have also been identified in psoriatic skin samples. These IL-22 producing cells express skin homing chemokines receptor CCR10 [35,36]. Finally, it is currently unclear whether innate immune cells such as NK cells contribute to IL-22 expression in psoriatic skin, although these cells are known sources of IL-22.

Keratinocytes are the major downstream targets of IL-22 in the skin [53,54,63]. Many of the downstream phenotypes induced by IL-22 from keratinocytes are also observed in epidermal




Fig. 1. Pathogenic role of IL-22 in psoriasis. In psoriasis, IL-23 promotes the production of IL-22 from various T cell subsets, including Th1, Th17, Th22, as well as CD8 T cells. IL-22 induces many histological features of psoriasis in the epidermis, such as acanthosis, hypogranulosis, and parakeratosis. In addition, IL-22 also stimulates the expression of antimicrobial peptides, chemokines, and proteins involved in wound healing.

keratinocytes from lesional psoriatic skin. For example, in cultured reconstituted human epidermis (RHE), IL-22 induces prolonged Stat3 activation, which is readily detectable in involved psoriatic epidermis [53,64]. In vitro, IL-22 treatment results in pronounced proliferation and altered differentiation process of the cultured epidermal keratinocytes, resembling acanthosis of psoriatic skin [53,54]. Normal epidermis contains four major keratinocyte layers, basal layer, spinous layer, granular layer, and cornified layer (Fig. 1) [47]. In cultured epidermis, IL-22 enhances the proliferative capacity of epidermal keratinocytes, leading to increase overall thickness of the RHE [53,54]. Similar to what observed in psoriatic epidermis, there is an increase of total spinous layers when RHE is treated with IL-22. Interestingly, IL-22 treatment also resulted in the loss of granular layer (hypogranulosis), which is characterized with cytoplasmic keratohyalin granules, in the RHE. Hypogranulosis is a key histological feature of psoriatic skin. Finally, IL-22 endows the development of parakeratosis, another histological feature of psoriatic skin, in the cornified layer in culture [65].

In addition to these cellular changes, IL-22 induces the expression of many genes that involved in innate host defense, inflammatory response, wound healing, re-epithelialization and differentiation from RHE [53,54,63]. For examples, IL-22 regulates the expression of many antimicrobial peptides such as S100 family proteins (e.g., S100A7, S100A8) and β -defensin family proteins. In addition, it induces chemokines including IL-8, CXCL1, CXCL7 and CCL20 from keratinocytes. IL-22 also promotes the expression of autocrine growth factors, such as IL-20 and HB-EGF, from these cells [53,66]. Finally, many genes, including kallikrein subgroup of serine proteases and serpin family of protease inhibitors, induced by IL-22, may participate in wound healing and tissue repairing.

In the pathogenesis of psoriasis, many other cytokines are also upregulated [48,49]. Many downstream effects and genes induced by IL-22 from keratinocytes can be regulated by other cytokines [53]. It is relatively easy to differentiate the functions of IL-22 from those factors that regulate only inflammation or epithelial homeostasis. For instance, IL-1 β and TNF- α can induce the expression of many of the same antimicrobial peptides and

chemokines. But these two cytokines do not promote keratinocyte hyperplasia and differentiation. On the other hand, EGF and TGF- α are strong stimuli for the proliferation of keratinocytes, whereas both of them do not induce hypogranulosis, persistent Stat3 activation, and the expression of many antimicrobial peptides, such as S100A7 [53]. In fact, the overall gene signatures regulated by EGF and TGF- α are negatively correlated with genes altered in psoriatic skin, while genes regulated by IL-22 are strongly correlative [53].

Nevertheless, the overall functions of IL-22 on keratinocytes are shared by other Stat3-activating cytokines, including proteins in its own family, IL-19, IL-20 and IL-24, as well as Oncostatin M (OSM) and IL-6 [53,67]. All these cytokines are upregulated in psoriatic skin. One feature makes IL-22 unique among these cytokines is its tight association with IL-23 pathway. IL-23 is a heterodimeric cytokine composed of its unique p19 subunit and the p40 subunit, which is shared with IL-12 [68]. IL-23 signals through IL-23R chain and IL-12R β 1 chain that are expressed primarily on various leukocyte subsets. IL-23 is upregulated in psoriatic skin and plays an indispensable pathogenic role in the development of psoriasis [69,70]. Genome-wide association studies have linked both *IL23R* and *IL12B* (encoded p40) genes to psoriasis [51]. In preclinical mouse models, overexpression of IL-23 or injection of IL-23 into the skin leads to the development of inflammatory skin phenotypes bearing some features of human psoriatic skin [18,71,72]. Most importantly, anti-p40 antibodies, which neutralizes both IL-12 and IL-23, in clinic have generated great efficacy in the treatment of psoriasis [52]. We have shown that IL-23 treatment in vitro can induce IL-22 production from many different leukocytes [18]. Furthermore, in multiple in vivo models, the induction of IL-22 is compromised in mice deficient of IL-23, suggesting that IL-22 is one of the direct downstream effector cytokines of IL-23 [28,46,73]. In preclinical psoriatic models such as the CD45Rb^{hi} T cell transfer model of psoriasis, blocking IL-22 or IL-23 results in similar efficacy in ameliorating the skin symptoms [74]. IL-23 triggered skin inflammation and keratinocytes hyperplasia is reduced in IL-22 deficient mice or in WT mice treated with anti-IL-22 antibody, again supporting IL-22 as a direct downstream mediator of IL-23 pathway




Fig. 2. Protective function of IL-22 in IBD. In the intestine, both innate and adaptive immune cells are capable in making IL-22 in responding to the stimulation of IL-23. NK like cells and LTi cells are the major innate sources of IL-22. IL-22 induces the expression of various antimicrobial peptides from intestinal epithelial cells. IL-22 also helps the regeneration of goblet cells and the production of mucin proteins. In addition, IL-22 promotes the wound healing responses and help to restore the integrity of the epithelial layer.

in psoriasis [18]. Taken together, IL-22 mediates the crosstalk between leukocytes and keratinocytes in psoriasis.

In spite of its functional importance, IL-22 is probably not sufficient in triggering psoriasis by itself. IL-22 primarily targets epithelial cells and is not able to profoundly amplify and sustain the inflammatory responses in disease [44]. The first evidence linking IL-20 subfamily cytokines including IL-22 to psoriasis came from the phenotypes observed in IL-20 transgenic mice [75]. These mice die soon after birth with skin phenotypes similar to psoriatic skin. The infiltration of leukocytes, especially T cells, however, is not prominently observed in the affected skin. Similar phenotypes have also been observed in IL-22 transgenic mice, which suggest IL-22 alone is not sufficient in initiation and amplification of inflammation [76]. In fact, in many other tissues, the role of IL-22 in tissue protection and in innate host defense helps to dampen the inflammatory responses [77–79]. By comparing the functions of IL-22 with those of IL-17, another tissue targeting cytokine, it is clear that IL-22 induces less proinflammatory responses from keratinocytes [65]. For example, the expression of chemokines such as CXCL1 and CXCL3 induced by IL-17 is much higher than that by IL-22 from keratinocytes. In addition, IL-17 but not IL-22 induces high level of IL-6 and G-CSF from primary fibroblasts. Therefore, in order to sustain the inflammatory cascades during psoriasis, IL-22 needs to synergize with other cytokines. It has been reported that IL-22 synergizes with IL-17 to enhance the induction of antimicrobial proteins from keratinocytes [17]. Furthermore, IL-22, IL-17, OSM, IL-1 α and TNF- α can synergistically induce some skin features recapitulating psoriasis closely both in vitro and in vivo [80].

Taken together, current data support that IL-22, although is not sole actor, plays an essential role in the pathogenesis of psoriasis. IL-22 synergizes with other leukocyte cytokines to form a cytokine

network that orchestrates the progression of many different pathogenic features of psoriasis.

3. Protective role of IL-22 in IBD

IBD is chronic, relapsing inflammatory diseases of gastrointestinal tract. Ulcerative colitis (UC) and Crohn's disease (CD) are two idiopathic forms of IBD, which have distinct clinical and histological features [81,82]. Inflammation in CD is usually transmural and discontinuous and involves the ileum and colon, while in UC inflammation is typically superficial and only affects colon [82]. Although the exact causes of IBD are still unclear, both environmental factors and genetic predisposition are involved in the onset and progression of the disease. Genetic evidence and data from preclinical models suggest that the inflammatory responses in IBD may result from inappropriate activation of the immune system by the normal luminal flora. Recent genome-wide association studies in IBD have identified the major genetic associations, which can be divided into genes that contribute to the epithelial integrity, innate immune response and adaptive immune response. UC and CD share some common genetic association alleles such as *IL23R* and *IL12B*, as well as alleles that are unique for each disease [83–86].

Detailed functional analysis of the SNPs associated *IL23R* and data from preclinical studies support a pathogenic role of IL-23 pathway in the pathogenesis of IBD [87]. First, T cells from people bearing the protective SNP (R381Q) in *IL23R* are less responsive to the stimulation by IL-23 in vitro, resulting in decreased Stat3 activation and IL-22 production, than those from control group (Pidashveva S. Manuscript submitted). Second, elevated IL-23 expression has been detected in CD and UC patients with activated

disease [88,89]. Finally, blocking IL-23 pathway lead to reduced disease severity in several preclinical IBD models [90–93]. Taken together, one would expect that blocking IL-23 might provide promising therapeutic benefits for the treatment of IBD in human as well. However, the initial data from clinic trials with anti-p40 antibody (ustekinumab) for the treatment of CD indicate only mild efficacy, less impressive than that in psoriasis [94]. Although further clinic trials are required to explore the potential therapeutic value of anti-p40 in the treatment of IBD, it is important to understand the downstream mechanisms, which may result in different clinic outcomes in psoriasis and IBD, of IL-23 pathway.

As we discussed above, IL-22 is a direct downstream effector cytokine of IL-23. Unlike its pathogenic functions in psoriasis, IL-22, supported by accumulating data, may exert a protective role in IBD. In the gastrointestinal tract, IL-23 is indispensable for IL-22 production, as in IL-23 deficient mice the induction of IL-22 triggered by various infections, for example, *C. rodentium* or *Toxoplasma gondii*, are abolished [73,95]. Although IL-22 and its receptor have not been linked with IBD genetically, expression of IL-22 is augmented in patients with IBD [96,97]. Increased IL-22, but not IL-17 and IFN- γ , has been reported in serum samples from CD patients [98]. The IL-22 levels also correlate with disease severity [99]. Interestingly, there is a correlation of IL-22 expression with IL-23R minor alleles. The proactive allele of IL-23R, which leads to hypo-responsiveness of T cells to IL-23 as discussed above, results in lower IL-22 expression in patients [99].

Despite the fact that IL-22 is correlated with disease activities in IBD patients, data from several preclinical models suggest that IL-22 exerts protective functions (Fig. 2). IL-22 deficient mice or WT mice treated with IL-22 neutralizing antibody lead to more severe weight loss, delayed recovery, and augmented epithelial damage and inflammation in the colon during dextran sodium sulfate (DSS) induced colitis [29,79,100,101]. In addition, T cells from IL-22 $^{-/-}$ mice cause a more severe disease in the T cell transfer model of IBD [101]. On the contrary, overexpression of IL-22 in TCR α deficient mice reduces colonic thickness and disease score [79]. Finally, IL-22 plays an essential role in infection induced colitis as well. *C. rodentium* is bacterium that specifically attacks murine colon epithelial cells and causes severe colitis within two weeks following infection. But the infection and colitis can be self-resolved in WT mice upon the development of the host adaptive immune responses, especially the generation of anti- *C. rodentium* antibodies. However, in IL-22 deficient mice, all mice develop exacerbated colitis during the infection and most of the mice succumb to the infection in the second week of infection [95]. These data suggest IL-22 endows essential defense mechanisms against invading pathogens in the intestine.

What are the mechanisms that cause the apparently different functions of IL-22 in psoriasis and IBD? The distinct pathogenic features and the different cellular sources of IL-22 in these diseases may provide possible explanation. Although the exact environmental triggers for psoriasis and IBD are still unclear, the presence of vast microbial microbes in the gut is essential for the development of IBD [81,82]. The inappropriate activation of mucosal immune system by the massive luminal microbial flora underlies the progression of IBD. On the other hand, while streptococcal infection has been linked with the onset of psoriasis, psoriatic skin is considered relatively sterile [102]. Therefore, the protective role of IL-22 in IBD is likely linked to its functions in enhancing epithelial innate defense mechanisms and barrier integrity, which sequester the direct interactions between the immune system and luminal commensal bacteria (Fig. 2). Similar protective functions of IL-22, nonetheless, are not obvious in the pathogenesis of psoriasis. In the intestinal tract, IL-22 induces the proliferation and reconstitution of mucosal epithelial cells through

the activation of Stat3 [29,79,96,100]. This increased healing response prevents further penetration of microbes into the epithelial layer. Second, IL-22 helps goblet cell to regenerate and produce mucus-associated proteins, which form the essential static external barrier separating intestinal flora with intestinal epithelial cells [79]. Finally, IL-22 stimulates the production and secretion of various antimicrobial peptides, including defensins, cathelicidins and C-type lectins, from intestinal epithelial cells and Paneth cells [28,29,79,100,101]. These antimicrobial peptides further sequester and directly kill the invading pathogens through various mechanisms [103].

Another distinction of IL-22 in psoriasis and IBD is its cellular sources. In human psoriasis, various T cell subsets are the major sources of IL-22, whereas in IBD innate immune cells as well as T cells in the intestine produce IL-22 upon activation [23,42,43]. T cells, upon activation concurrently produce other proinflammatory cytokines such as IL-17, IFN- γ and TNF- α , all of which can synergize with IL-22 in promoting the psoriatic features of keratinocytes and inflammatory cascades as discussed before. In the intestine, innate immune cells, such as LT α cells, NK cell subsets, and DCs may produce IL-22 in response to IL-23 stimulation alone [23,27,42,43,95]. It is unclear whether these cells can secrete other proinflammatory cytokines under the same condition. It is possible that IL-22 is the dominant cytokine from these cells under certain conditions in disease. As a result, IL-22 itself will not amplify the inflammatory responses in the absence of concurrent expression of other cytokines. Instead, its major function in gut is protective through promoting epithelial defense and healing process.

4. Conclusion


Clearly, we are still at the early stage of understanding the functional roles of IL-22 in various human diseases. Future studies will focus on dissecting specific biology of IL-22 in a given tissue environment and disease context, since its complicated functions so far forbid us to generalize its role. The protective functions of IL-22 in IBD also suggest IL-23 may rely on other factors mediating its pathogenic functions. Identifying these factors may offer better therapeutic opportunities in IBD, in which one only needs to block these factors downstream of IL-23 pathway, but leave the protective functions of IL-22 intact.

References

- [1] Dumoutier L, Louahed J, Renauld JC. Cloning and characterization of IL-10-related T cell-derived inducible factor (IL-TIF), a novel cytokine structurally related to IL-10 and inducible by IL-9. *J Immunol* 2000;164:1814–9.
- [2] Xie MH, Aggarwal S, Ho WH, Foster J, Zhang Z, Stinson J, et al. Interleukin (IL)-22, a novel human cytokine that signals through the interferon receptor-related proteins CRF2-4 and IL-22R. *J Biol Chem* 2000;275:31335–9.
- [3] Nagem RA, Colau D, Dumoutier L, Renauld JC, Ogata C, Polikarpov I. Crystal structure of recombinant human interleukin-22. *Structure* 2002;10:1051–62.
- [4] Pestka S, Krause CD, Sarkar D, Walter MR, Shi Y, Fisher PB. Interleukin-10 and related cytokines and receptors. *Annu Rev Immunol* 2004;22:929–79.
- [5] Krause CD, Mei E, Mirochnitchenko O, Lavnikova N, Xie J, Jia Y, et al. Interactions among the components of the interleukin-10 receptor complex. *Biochem Biophys Res Commun* 2006;340:377–85.
- [6] Jones BC, Logsdon NJ, Walter MR. Structure of IL-22 bound to its high-affinity IL-22R1 chain. *Structure* 2008.
- [7] Bleicher L, de Moura PR, Watanabe L, Colau D, Dumoutier L, Renauld JC, et al. Crystal structure of the IL-22/IL-22R1 complex and its implications for the IL-22 signaling mechanism. *FEBS Lett* 2008;582:2985–92.
- [8] Lejeune D, Dumoutier L, Constantinescu S, Kruijer W, Schuringa JJ, Renauld JC. Interleukin-22 (IL-22) activates the JAK/STAT, ERK, JNK, and p38 MAP kinase pathways in a rat hepatoma cell line. Pathways that are shared with and distinct from IL-10. *J Biol Chem* 2002;277:33676–82.
- [9] Bartlett NW, Dumoutier L, Renauld JC, Kotenko SV, McVey CE, Lee HJ, et al. A new member of the interleukin 10-related cytokine family encoded by a poxvirus. *J Gen Virol* 2004;85:1401–12.
- [10] Kotenko SV, Izotova LS, Mirochnitchenko OV, Esterova E, Dickensheets H, Donnelly RP, et al. Identification, cloning, and characterization of a novel soluble receptor that binds IL-22 and neutralizes its activity. *J Immunol* 2001;166:7096–103.

- [11] Dumoutier L, Lejeune D, Hor S, Fickscher H, Renauld JC. Cloning of a new type II cytokine receptor activating signal transducer and activator of transcription (STAT)1, STAT2 and STAT3. *Biochem J* 2003;370:391–6.
- [12] Xu W, Presnell SR, Parrish-Novak J, Kindsvogel W, Jaspers S, Chen Z, et al. A soluble class II cytokine receptor, IL-22RA2, is a naturally occurring IL-22 antagonist. *Proc Natl Acad Sci U S A* 2001;98:9511–6.
- [13] Wolk K, Kunz S, Asadullah K, Sabat R. Cutting edge: immune cells as sources and targets of the IL-10 family members? *J Immunol* 2002;168:5397–402.
- [14] Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al. Interleukin 17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol* 2005;6:1123–32.
- [15] Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang YH, et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nat Immunol* 2005;6:1133–41.
- [16] Ouyang W, Kolls JK, Zheng Y. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* 2008;28:454–67.
- [17] Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* 2006;203:2271–9.
- [18] Zheng Y, Danilenko D, Valdez P, Kasman I, Eastham-Anderson J, Wu J, et al. Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* 2007;445:648–51.
- [19] Chung Y, Yang X, Chang SH, Ma L, Tian Q, Dong C. Expression and regulation of IL-22 in the IL-17-producing CD4⁺ T lymphocytes. *Cell Res* 2006;16:902–7.
- [20] Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGFβ in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 2006;24:179–89.
- [21] Mangan PR, Harrington LE, O'Quinn DB, Helms WS, Bullard DC, Elson CO, et al. Transforming growth factor-β induces development of the T(H)17 lineage. *Nature* 2006;441:231–4.
- [22] Goto M, Murakawa M, Kadoshima-Yamaoka K, Tanaka Y, Nagahira K, Fukuda Y, et al. Murine NKT cells produce Th17 cytokine interleukin-22. *Cell Immunol* 2009;254:81–4.
- [23] Cella M, Fuchs A, Vermi W, Facchetti F, Otero K, Lennerz JK, et al. A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature* 2009;457:722–5.
- [24] Luci C, Reynders A, Ivanov II, Cognet C, Chiche L, Chasson L, et al. Influence of the transcription factor RORγ₁ on the development of Nkp46⁺ cell populations in gut and skin. *Nat Immunol* 2009;10:75–82.
- [25] Sanos SL, Bui VL, Mortha A, Oberle K, Heners C, Johnner C, et al. RORγ₁ and commensal microflora are required for the differentiation of mucosal interleukin 22-producing Nkp46⁺ cells. *Nat Immunol* 2009;10:83–91.
- [26] Satoh-Takayama N, Vosschenrich CAJ, Lesjean-Pottier S, Sawa S, Lochner M, Rattis F, et al. Microbial flora drives interleukin 22 production in intestinal Nkp46⁺ cells that provide innate mucosal immune defense. *Immunity* 2008;29:958–70.
- [27] Takatori H, Kanno Y, Watford WT, Tato CM, Weiss G, Ivanov II, et al. Lymphoid tissue inducer-like cells are an innate source of IL-17 and IL-22. *J Exp Med* 2009;206:35–41.
- [28] Zheng Y, Valdez P, Danilenko D, Hu Y, Sa S, Gong Q, et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat Med* 2008;14:282–9.
- [29] Pickert G, Neufert C, Leppkes M, Zheng Y, Wittkopf N, Warntjen M, et al. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J Exp Med* 2009;206:1465–72.
- [30] De Luca A, Zelante T, D'Angelo C, Zagarella S, Fallarino F, Spreca A, et al. IL-22 defines a novel immune pathway of antifungal resistance. *Mucosal Immunol* 2010;3:361–73.
- [31] Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia A, et al. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol* 2007;8:639–46.
- [32] Wilson NJ, Boniface K, Chan JR, McKenzie BS, Blumenschein WM, Mattson JD, et al. Development, cytokine profile and function of human interleukin 17-producing helper T cells. *Nat Immunol* 2007;8:950–7.
- [33] Kleinschek MA, Boniface K, Sadekova S, Grein J, Murphy EE, Turner SP, et al. Circulating and gut-resident human Th17 cells express CD161 and promote intestinal inflammation. *J Exp Med* 2009;206:525–34.
- [34] Billerbeck E, Kang YH, Walker L, Lockstone H, Grafmueller S, Fleming V, et al. Analysis of CD161 expression on human CD8⁺ T cells defines a distinct functional subset with tissue-homing properties. *Proc Natl Acad Sci U S A* 2010;107:3006–11.
- [35] Duhon T, Geiger R, Jarrossay D, Lanzavecchia A, Sallusto F. Production of interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T cells. *Nat Immunol* 2009;10:857–63.
- [36] Eyerich S, Eyerich K, Pennino D, Carbone T, Nasorri F, Pallotta S, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* 2009;119:3573–85.
- [37] Trifari S, Kaplan C, Tran E, Crellin N, Spits H. Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)-1 and T(H)-2 cells. *Nat Immunol* 2009.
- [38] Fujita H, Nogales KE, Kikuchi T, Gonzalez J, Carucci JA, Krueger JG. Human Langerhans cells induce distinct IL-22-producing CD4⁺ T cells lacking IL-17 production. *Proc Natl Acad Sci U S A* 2009;106:21795–800.
- [39] Vivier E, Spits H, Cupedo T. Interleukin-22-producing innate immune cells: new players in mucosal immunity and tissue repair? *Nat Rev Immunol* 2009;9:229–34.
- [40] Colonna M. Interleukin-22-producing natural killer cells and lymphoid tissue inducer-like cells in mucosal immunity. *Immunity* 2009;31:15–23.
- [41] Hughes T, Becknell B, McClory S, Briercheck E, Freud AG, Zhang X, et al. Stage 3 immature human natural killer cells found in secondary lymphoid tissue constitutively and selectively express the TH 17 cytokine interleukin-22. *Blood* 2009;113:4008–10.
- [42] Crellin NK, Trifari S, Kaplan CD, Cupedo T, Spits H. Human Nkp44 + IL-22+ cells and LT_i-like cells constitute a stable RORC⁺ lineage distinct from conventional natural killer cells. *J Exp Med* 2010;207:281–90.
- [43] Cupedo T, Crellin NK, Papazian N, Rombouts EJ, Weijer K, Grogan JL, et al. Human fetal lymphoid tissue-inducer cells are interleukin 17-producing precursors to RORC⁺ CD127⁺ natural killer-like cells. *Nat Immunol* 2009;10:66–74.
- [44] Wolk K, Kunz S, Witte E, Friedrich M, Asadullah K, Sabat R. IL-22 increases the innate immunity of tissues. *Immunity* 2004;21:241–54.
- [45] Ouyang W, Valdez P. IL-22 in mucosal immunity. *Mucosal Immunol* 2008;1:335–8.
- [46] Aujla SJ, Chan YR, Zheng M, Fei M, Askew DJ, Pociask DA, et al. IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nat Med* 2008;14:275–81.
- [47] Bowcock AM, Krueger JG. Getting under the skin: the immunogenetics of psoriasis. *Nat Rev Immunol* 2005;5:699–711.
- [48] Nestle FO, Kaplan DH, Barker J. N Engl J Med 2009;361:496–509.
- [49] Nickoloff BJ, Nestle FO. Recent insights into the immunopathogenesis of psoriasis provide new therapeutic opportunities. *J Clin Invest* 2004;113:1664–75.
- [50] Nickoloff BJ. Cracking the cytokine code in psoriasis. *Nat Med* 2007;13:242–4.
- [51] Cargill M, Schrodi SJ, Chang M, Garcia VE, Brandon R, Callis KP, et al. A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 2007;80:273–90.
- [52] Krueger GG, Langley RG, Leonardi C, Yeilding N, Guzzo C, Wang Y, et al. A human interleukin-12/23 monoclonal antibody for the treatment of psoriasis. *N Engl J Med* 2007;356:580–92.
- [53] Sa SM, Valdez PA, Wu J, Jung K, Zhong F, Hall L, et al. The effects of IL-20 subfamily cytokines on reconstituted human epidermis suggest potential roles in cutaneous innate defense and pathogenic adaptive immunity in psoriasis. *J Immunol* 2007;178:2229–40.
- [54] Boniface K, Bernard FX, Garcia M, Gurney AL, Lecron JC, Morel F. IL-22 inhibits epidermal differentiation and induces proinflammatory gene expression and migration of human keratinocytes. *J Immunol* 2005;174:3695–702.
- [55] Lo YH, Torii K, Saito C, Furuhashi T, Maeda A, Morita A. Serum IL-22 correlates with psoriatic severity and serum IL-6 correlates with susceptibility to phototherapy. *J Dermatol Sci* 2010;58:225–7.
- [56] Boniface K, Guignouard E, Pedretti N, Garcia M, Delwail A, Bernard FX, et al. A role for T cell-derived interleukin 22 in psoriatic skin inflammation. *Clin Exp Immunol* 2007;150:407–15.
- [57] Lowes MA, Kikuchi T, Fuentes-Duculan J, Cardinale I, Zaba LC, Haider AS, et al. Psoriasis vulgaris lesions contain discrete populations of Th1 and Th17 T cells. *J Invest Dermatol* 2008;128:1207–11.
- [58] Lew W, Bowcock AM, Krueger JG. Psoriasis vulgaris: cutaneous lymphoid tissue supports T-cell activation and “type 1” inflammatory gene expression. *Trends Immunol* 2004;25:295–305.
- [59] Lowes MA, Bowcock AM, Krueger JG. Pathogenesis and therapy of psoriasis. *Nature* 2007;445:866–73.
- [60] Kagami S, Rizzo HL, Lee JJ, Koguchi Y, Blauvelt A. Circulating Th17, Th22, and Th1 cells are increased in psoriasis. *J Invest Dermatol* 2010;130:1373–83.
- [61] Haider AS, Lowes MA, Suarez-Farinas M, Zaba LC, Cardinale I, Khatcherian A, et al. Identification of cellular pathways of “type 1”, Th17 T cells, and TNF- and inducible nitric oxide synthase-producing dendritic cells in autoimmune inflammation through pharmacogenomic study of cyclosporine A in psoriasis. *J Immunol* 2008;180:1913–20.
- [62] Ortega C, Fernandez AS, Carrillo JM, Romero P, Molina JJ, Moreno JC, et al. IL-17-producing CD8⁺ T lymphocytes from psoriasis skin plaques are cytotoxic effector cells that secrete Th17-related cytokines. *J Leukoc Biol* 2009;86:435–43.
- [63] Wolk K, Witte E, Wallace E, Docke WD, Kunz S, Asadullah K, et al. IL-22 regulates the expression of genes responsible for antimicrobial defense, cellular differentiation, and mobility in keratinocytes: a potential role in psoriasis. *Eur J Immunol* 2006;36:1309–23.
- [64] Sano S, Chan KS, Carbajal S, Clifford J, Peavey M, Kiguchi K, et al. Stat3 links activated keratinocytes and immunocytes required for development of psoriasis in a novel transgenic mouse model. *Nat Med* 2005;11:43–9.
- [65] Nogales KE, Zaba LC, Guttman-Yassky E, Fuentes-Duculan J, Suarez-Farinas M, Cardinale I, et al. Th17 cytokines interleukin (IL)-17 and IL-22 modulate distinct inflammatory and keratinocyte-response pathways. *Br J Dermatol* 2008;159:1092–102.
- [66] Wolk K, Witte E, Warszawska K, Schulze-Tanzil G, Witte K, Philipp S, et al. The Th17 cytokine IL-22 induces IL-20 production in keratinocytes: a novel immunological cascade with potential relevance in psoriasis. *Eur J Immunol* 2009;39:3570–81.
- [67] Boniface K, Diveu C, Morel F, Pedretti N, Froger J, Rapon E, et al. Oncostatin M secreted by skin infiltrating T lymphocytes is a potent keratinocyte activator involved in skin inflammation. *J Immunol* 2007;178:4615–22.

- [68] Oppmann B, Lesley R, Blom B, Timans JC, Xu Y, Hunte B, et al. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 2000;13:715–25.
- [69] Lee E, Trepicchio WL, Oestreicher JL, Pittman D, Wang F, Chamian F, et al. Increased expression of interleukin 23 p19 and p40 in lesional skin of patients with psoriasis vulgaris. *J Exp Med* 2004;199:125–30.
- [70] Di Cesare A, Di Meglio P, Nestle FO. The IL-23/Th17 axis in the immunopathogenesis of psoriasis. *J Invest Dermatol* 2009;129:1339–50.
- [71] Kopp T, Lenz P, Bello-Fernandez C, Kastelein RA, Kupper TS, Stingl G. IL-23 production by cosecretion of endogenous p19 and transgenic p40 in keratin 14/p40 transgenic mice: evidence for enhanced cutaneous immunity. *J Immunol* 2003;170:5438–44.
- [72] Chan JR, Blumenschein W, Murphy E, Diveu C, Wiekowski M, Abbondanzo S, et al. IL-23 stimulates epidermal hyperplasia via TNF and IL-20R2-dependent mechanisms with implications for psoriasis pathogenesis. *J Exp Med* 2006;203:2577–87.
- [73] Munoz M, Heimesaat MM, Danker K, Struck D, Lohmann U, Plickert R, et al. Interleukin (IL)-23 mediates *Toxoplasma gondii*-induced immunopathology in the gut via matrix metalloproteinase-2 and IL-22 but independent of IL-17. *J Exp Med* 2009;206:3047–59.
- [74] Ma HL, Liang S, Li J, Napierata L, Brown T, Benoit S, et al. IL-22 is required for Th17 cell-mediated pathology in a mouse model of psoriasis-like skin inflammation. *J Clin Invest* 2008;118:597–607.
- [75] Blumberg H, Conklin D, Xu WF, Grossmann A, Brender T, Carollo S, et al. Interleukin 20: discovery, receptor identification, and role in epidermal function. *Cell* 2001;104:9–19.
- [76] Wolk K, Haugen HS, Xu W, Witte E, Waggie K, Anderson M, et al. IL-22 and IL-20 are key mediators of the epidermal alterations in psoriasis while IL-17 and IFN- γ are not. *J Mol Med* 2009;87:523–36.
- [77] Zenewicz LA, Yancopoulos GD, Valenzuela DM, Murphy AJ, Karow M, Flavell RA. Interleukin-22 but not interleukin-17 provides protection to hepatocytes during acute liver inflammation. *Immunity* 2007;27:647–59.
- [78] Radaeva S, Sun R, Pan HN, Hong F, Gao B. Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine hepatitis: IL-22 is a survival factor for hepatocytes via STAT3 activation. *Hepatology* 2004;39:1332–42.
- [79] Sugimoto K, Ogawa A, Mizoguchi E, Shimomura Y, Andoh A, Bhan AK, et al. IL-22 ameliorates intestinal inflammation in a mouse model of ulcerative colitis. *J Clin Invest* 2008;118:534–44.
- [80] Guilloteau K, Paris I, Pedretti N, Boniface K, Juchaux F, Huguier V, et al. Skin inflammation induced by the synergistic action of IL-17A, IL-22, oncostatin M, IL-1 α , and TNF- α recapitulates some features of psoriasis. *J Immunol* 2010;184:5263–70.
- [81] Xavier RJ, Podolsky DK. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448:427–34.
- [82] Podolsky DK. Inflammatory bowel disease. *N Engl J Med* 2002;347:417–29.
- [83] Rioux JD, Xavier RJ, Taylor KD, Silverberg MS, Goyette P, Huett A, et al. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet* 2007;39:596–604.
- [84] Barrett JC, Hansoul S, Nicolae DL, Cho JH, Duerr RH, Rioux JD, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat Genet* 2008;40:955–62.
- [85] Barrett JC, Lee JC, Lees CW, Prescott NJ, Anderson CA, Phillips A, et al. Genome-wide association study of ulcerative colitis identifies three new susceptibility loci, including the HNF4A region. *Nat Genet* 2009;41:1330–4.
- [86] Cho JH. The genetics and immunopathogenesis of inflammatory bowel disease. *Nat Rev Immunol* 2008;8:458–66.
- [87] Duerr RH, Taylor KD, Brant SR, Rioux JD, Silverberg MS, Daly MJ, et al. A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. *Science* 2006;314:1461–3.
- [88] Fuss IJ, Becker C, Yang Z, Groden C, Hornung RL, Heller F, et al. Both IL-12p70 and IL-23 are synthesized during active Crohn's disease and are down-regulated by treatment with anti-IL-12 p40 monoclonal antibody. *Inflamm Bowel Dis* 2006;12:9–15.
- [89] Schmidt C, Giese T, Ludwig B, Mueller-Molaian I, Marth T, Zeuzem S, et al. Expression of interleukin-12-related cytokine transcripts in inflammatory bowel disease: elevated interleukin-23p19 and interleukin-27p28 in Crohn's disease but not in ulcerative colitis. *Inflamm Bowel Dis* 2005;11:16–23.
- [90] Kullberg MC, Jankovic D, Feng CG, Hue S, Gorelick PL, McKenzie BS, et al. IL-23 plays a key role in *Helicobacter hepaticus*-induced T cell-dependent colitis. *J Exp Med* 2006;203:2485–94.
- [91] Hue S, Ahern P, Buonocore S, Kullberg MC, Cua DJ, McKenzie BS, et al. Interleukin-23 drives innate and T cell-mediated intestinal inflammation. *J Exp Med* 2006;203:2473–83.
- [92] Elson CO, Cong Y, Weaver CT, Schoeb TR, McClanahan TK, Fick RB, et al. Monoclonal anti-interleukin 23 reverses active colitis in a T cell-mediated model in mice. *Gastroenterology* 2007;132:2359–70.
- [93] Uhlig HH, McKenzie BS, Hue S, Thompson C, Joyce-Shaikh B, Stepankova R, et al. Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune pathology. *Immunity* 2006;25:309–18.
- [94] Toedter GP, Blank M, Lang Y, Chen D, Sandborn WJ, de Villiers WJ. Relationship of C-reactive protein with clinical response after therapy with ustekinumab in Crohn's disease. *Am J Gastroenterol* 2009;104:2768–73.
- [95] Zheng Y, Valdez PA, Danilenko DM, Hu Y, Sa SM, Gong Q, et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat Med* 2008;14:282–9.
- [96] Brand S, Beigel F, Olszak T, Zitzmann K, Eichhorst ST, Otte JM, et al. IL-22 is increased in active Crohn's disease and promotes proinflammatory gene expression and intestinal epithelial cell migration. *Am J Physiol Gastrointest Liver Physiol* 2006;290:G827–38.
- [97] Andoh A, Zhang Z, Inatomi O, Fujino S, Deguchi Y, Araki Y, et al. Interleukin-22, a member of the IL-10 subfamily, induces inflammatory responses in colonic subepithelial myofibroblasts. *Gastroenterology* 2005;129:969–84.
- [98] Wolk K, Witte E, Hoffmann U, Doecke WD, Endesfelder S, Asadullah K, et al. IL-22 induces lipopolysaccharide-binding protein in hepatocytes: a potential systemic role of IL-22 in Crohn's disease. *J Immunol* 2007;178:5973–81.
- [99] Schmechel S, Konrad A, Diegelmann J, Glas J, Wetzke M, Paschos E, et al. Linking genetic susceptibility to Crohn's disease with Th17 cell function: IL-22 serum levels are increased in Crohn's disease and correlate with disease activity and IL23R genotype status. *Inflamm Bowel Dis* 2008;14:204–12.
- [100] Neufert C, Pickert G, Zheng Y, Wittkopf N, Wirtgen J, Nikolaev A, et al. Activation of epithelial STAT3 regulates intestinal homeostasis. *Cell Cycle* 2010;9:652–5.
- [101] Zenewicz LA, Yancopoulos GD, Valenzuela DM, Murphy AJ, Stevens S, Flavell RA. Innate and adaptive interleukin-22 protects mice from inflammatory bowel disease. *Immunity* 2008;29:947–57.
- [102] Prinz JC. Psoriasis vulgaris – a sterile antibacterial skin reaction mediated by cross-reactive T cells? An immunological view of the pathophysiology of psoriasis. *Clin Exp Dermatol* 2001;26:326–32.
- [103] Kolls JK, McCray Jr PB, Chan YR. Cytokine-mediated regulation of antimicrobial proteins. *Nat Rev Immunol* 2008;8:829–35.



Wenjun Ouyang, Ph.D., is currently a senior scientist and group leader in Immunology Department, Genentech, Inc. His group is focused on identifying and developing novel therapies for the treatment of various autoimmune diseases. His group has brought several new therapeutic candidates from early research into clinical trials. His research is interested in understanding the biology of T helper cells and their effector cytokines in autoimmune diseases. Before he joined Genentech, Inc. he had obtained his Ph.D. (1999) and postdoc training (2001) from Washington University, School of Medicine, in St. Louis.