ID: 128c75e2 Question 1

The function g is defined by $g(x) = \frac{|x|}{a} - 14$, where $a < 0$. What is the product of $g(15a)$ and $g(7a)$?

ID: 91e7ea5e Question 2

$h(x) = 2(x-4)^2 - 32$

The quadratic function *h* is defined as shown. In the *xy*-plane, the graph of $y = h(x)$ intersects the *x*-axis at the points $(0,0)$ and $(t,0)$, where *t* is a constant. What is the value of *t* ?

A. 1

B. 2

C. 4

D. 8

ID: ebed7dc6 Question 3

An auditorium has seats for $1{,}800$ people. Tickets to attend a show at the auditorium currently cost $\$4.00$. For each $\$1.00$ increase to the ticket price, 100 fewer tickets will be sold. This situation can be modeled by the equation $y = -100x^2 + 1{,}400x + 7{,}200$, where x represents the increase in ticket price, in dollars, and y represents the revenue, in dollars, from ticket sales. If this equation is graphed in the *xy*-plane, at what value of x is the maximum of the graph?

A. 4

B. 7

C. 14

D. 18

ID: a9084ca4 Question 4

$$f(x) = 9{,}000(0.66)^x$$

The given function f models the number of advertisements a company sent to its clients each year, where x represents the number of years since 1997, and $0 \leq x \leq 5$. If $y = f(x)$ is graphed in the *xy*-plane, which of the following is the best interpretation of the *y*-intercept of the graph in this context?

A. The minimum estimated number of advertisements the company sent to its clients during the 5 years was $1{,}708$.

B. The minimum estimated number of advertisements the company sent to its clients during the 5 years was $9{,}000$.

C. The estimated number of advertisements the company sent to its clients in 1997 was $1{,}708$.

D. The estimated number of advertisements the company sent to its clients in 1997 was $9{,}000$.

ID: b8f13a3a Question 5

Function f is defined by $f(x) = -a^x + b$, where a and b are constants. In the *xy*-plane, the graph of $y = f(x) - 12$ has a *y*-intercept at $\left(0, -\frac{75}{7}\right)$. The product of a and b is $\frac{320}{7}$. What is the value of a?

ID: 9654add7 Question 6

$f(x) = -500x^2 + 25{,}000x$

The revenue $f(x)$, in dollars, that a company receives from sales of a product is given by the function *f* above, where *x* is the unit price, in dollars, of the product. The graph of $y = f(x)$ in the *xy*-plane intersects the *x*-axis at 0 and *a*. What does *a* represent?

A. The revenue, in dollars, when the unit price of the product is $0

B. The unit price, in dollars, of the product that will result in maximum revenue

C. The unit price, in dollars, of the product that will result in a revenue of $0

D. The maximum revenue, in dollars, that the company can make

ID: 263f9937 Question 7

Growth of a Culture of Bacteria

Day	Number of bacteria per milliliter at end of day
1	2.5×10^5
2	5.0×10^5
3	1.0×10^6

A culture of bacteria is growing at an exponential rate, as shown in the table above. At this rate, on which day would the number of bacteria per milliliter reach 5.12×10^8?

A. Day 5

B. Day 9

C. Day 11

D. Day 12

ID: 4dd4efcf Question 8

$$f(x) = ax^2 + 4x + c$$

In the given quadratic function, a and c are constants. The graph of $y = f(x)$ in the *xy*-plane is a parabola that opens upward and has a vertex at the point (h, k), where h and k are constants. If $k < 0$ and $f(-9) = f(3)$, which of the following must be true?

I. $c < 0$

II. $a \geq 1$

A. I only

B. II only

C. I and II

D. Neither I nor II

ID: 35e05e19 Question 9

A park ranger hung squirrel houses each in the shape of a right rectangular prism for fox squirrels. Each house has a height of 11 inches. The length of each house's base is x inches, which is 1 inch more than the width of the house's base. Which function V gives the volume of each house, in cubic inches, in terms of the length of the house's base?

A. $V(x) = 11x(x - 1)$

B. $V(x) = 11x(x + 1)$

C. $V(x) = x(x + 11)(x - 1)$

D. $V(x) = x(x + 11)(x + 1)$

ID: 18e35375 Question 10

$$f(x) = (x - 14)(x + 19)$$

The function f is defined by the given equation. For what value of x does $f(x)$ reach its minimum?

A. -266

B. -19

C. $-\frac{33}{2}$

D. $-\frac{5}{2}$

ID: 8462b105 Question 11

The function f gives the product of a number, x, and a number that is 91 more than x. Which equation defines f?

A. $f(x) = x^2 + x + 91$

B. $f(x) = x^2 + 91$

C. $f(x) = x^2 + 91x$

D. $f(x) = x^2 + 91x + 91$

ID: ce579859 Question 12

A model estimates that at the end of each year from 2015 to 2020, the number of squirrels in a population was 150% more than the number of squirrels in the population at the end of the previous year. The model estimates that at the end of 2016, there were 180 squirrels in the population. Which of the following equations represents this model, where n is the estimated number of squirrels in the population t years after the end of 2015 and $t \leq 5$?

A. $n = 72(1.5)^t$

B. $n = 72(2.5)^t$

C. $n = 180(1.5)^t$

D. $n = 180(2.5)^t$

ID: d139cf4b

Question 13

$$f(t) = 55t - 2t^2$$

The function f is defined by the given equation. The function g is defined by $g(t) = f(t) + 3$. Which expression represents the maximum value of $g(t)$?

A. $3 + \left(\frac{55}{2}\right)^2$

B. $3 + 2\left(\frac{55}{4}\right)^2$

C. $3 - 2\left(\frac{55}{4}\right)^2$

D. $3 - \left(\frac{55}{2}\right)^2$

ID: 9afe2370 Question 14

The population P of a certain city y years after the last census is modeled by the equation below, where r is a constant and P_0 is the population when $y = 0$.

$$P = P_0(1 + r)^y$$

If during this time the population of the city decreases by a fixed percent each year, which of the following must be true?


A. $r < -1$

B. $-1 < r < 0$

C. $0 < r < 1$

D. $r > 1$

ID: df71424b Question 15



The graph of $y = f(x)$ is shown, where $f(x) = ab^x + c$, and a, b, and c are constants. For how many values of x does $f(x) = 0$?

A. Three

B. Two

C. One

D. Zero

ID: 0121a235 Question 16

x	$p(x)$
-2	5
-1	0
0	-3
1	-1
2	0

The table above gives selected values of a polynomial function p. Based on the values in the table, which of the following must be a factor of p ?

A. $(x-3)$

B. $(x+3)$

C. $(x-1)(x+2)$

D. $(x+1)(x-2)$

ID: bba18ecb Question 17

When the quadratic function f is graphed in the *xy*-plane, where $y = f(x)$, its vertex is $(-3, 6)$. One of the *x*-intercepts of this graph is $\left(-\frac{17}{4}, 0\right)$. What is the other *x*-intercept of the graph?

A. $\left(-\frac{29}{4}, 0\right)$

B. $\left(-\frac{7}{4}, 0\right)$

C. $\left(\frac{5}{4}, 0\right)$

D. $\left(\frac{17}{4}, 0\right)$

ID: 668f1863 Question 18

Function f is a quadratic function where $f(-20) = 0$ and $f(-4) = 0$. The graph of $y = f(x)$ in the *xy*-plane has a vertex at $(r, -64)$. What is the value of r?

ID: 70753f99 Question 19

The function f is defined by $f(x)=(x+3)(x+1)$. The graph of f in the xy-plane is a parabola. Which of the following intervals contains the x-coordinate of the vertex of the graph of f ?

A. $-4<x<-3$

B. $-3<x<1$

C. $1<x<3$

D. $3<x<4$

ID: 58dcc59f Question 20

A landscaper is designing a rectangular garden. The length of the garden is to be 5 feet longer than the width. If the area of the garden will be 104 square feet, what will be the length, in feet, of the garden?

ID: 84dd43f8 Question 21

For the function f, $f(0) = 86$, and for each increase in x by 1, the value of $f(x)$ decreases by 80%. What is the value of $f(2)$?

ID: 59d1f4b5 Question 22

$M = 1{,}800(1.02)^t$

The equation above models the number of members, M, of a gym t years after the gym opens. Of the following, which equation models the number of members of the gym q quarter years after the gym opens?

A. $M = 1{,}800(1.02)^{\frac{q}{4}}$

B. $M = 1{,}800(1.02)^{4q}$

C. $M = 1{,}800(1.005)^{4q}$

D. $M = 1{,}800(1.082)^{q}$

ID: 635f54ee Question 23

The surface area of a cube is $6\left(\frac{a}{4}\right)^2$, where a is a positive constant. Which of the following gives the perimeter of one face of the cube?

A. $\frac{a}{4}$

B. a

C. $4a$

D. $6a$

ID: de39858a Question 24

The function h is defined by $h(x) = a^x + b$, where a and b are positive constants. The graph of $y = h(x)$ in the xy-plane passes through the points $(0, 10)$ and $(-2, \frac{325}{36})$. What is the value of ab?

A. $\frac{1}{4}$

B. $\frac{1}{2}$

C. 54

D. 60

ID: 1178f2df Question 25

x	y
21	-8
23	8
25	-8

The table shows three values of x and their corresponding values of y, where $y = f(x) + 4$ and f is a quadratic function. What is the *y*-coordinate of the *y*-intercept of the graph of $y = f(x)$ in the *xy*-plane?

ID: 84e8cc72 Question 26

A quadratic function models the height, in feet, of an object above the ground in terms of the time, in seconds, after the object is launched off an elevated surface. The model indicates the object has an initial height of 10 feet above the ground and reaches its maximum height of $1{,}034$ feet above the ground 8 seconds after being launched. Based on the model, what is the height, in feet, of the object above the ground 10 seconds after being launched?

A. 234

B. 778

C. 970

D. $1{,}014$

ID: 6d9e01a2

Question 27

$$f(x) = 4x^2 - 50x + 126$$

The given equation defines the function f. For what value of x does $f(x)$ reach its minimum?

ID: 9f2ecade Question 28

$$h(x) = x^3 + ax^2 + bx + c$$

The function h is defined above, where a, b, and c are integer constants. If the zeros of the function are -5, 6, and 7, what is the value of c ?

ID: 1fe10d97 Question 29

$$p(t) = 90{,}000(1.06)^t$$

The given function p models the population of Lowell t years after a census. Which of the following functions best models the population of Lowell m months after the census?

A. $r(m) = \frac{90{,}000}{12}(1.06)^m$

B. $r(m) = 90{,}000\left(\frac{1.06}{12}\right)^m$

C. $r(m) = 90{,}000\left(\frac{1.06}{12}\right)^{\frac{m}{12}}$

D. $r(m) = 90{,}000(1.06)^{\frac{m}{12}}$

ID: b73ee6cf Question 30

The population of a town is currently 50,000, and the population is estimated to increase each year by 3% from the previous year. Which of the following equations can be used to estimate the number of years, *t*, it will take for the population of the town to reach 60,000 ?

A. $50{,}000 = 60{,}000(0.03)^t$

B. $50{,}000 = 60{,}000(3)^t$

C. $60{,}000 = 50{,}000(0.03)^t$

D. $60{,}000 = 50{,}000(1.03)^t$

ID: 7eed640d Question 31

$$h(x) = -16x^2 + 100x + 10$$

The quadratic function above models the height above the ground *h*, in feet, of a projectile *x* seconds after it had been launched vertically. If $y = h(x)$ is graphed in the *xy*-plane, which of the following represents the real-life meaning of the positive *x*-intercept of the graph?

A. The initial height of the projectile

B. The maximum height of the projectile

C. The time at which the projectile reaches its maximum height

D. The time at which the projectile hits the ground

ID: 43926bd9 Question 32

x	$f(x)$
1	a
2	a^5
3	a^9

For the exponential function f, the table above shows several values of x and their corresponding values of $f(x)$, where a is a constant greater than 1. If k is a constant and $f(k) = a^{29}$, what is the value of k ?

ID: f25a34aa Question 33

The area of a triangle is equal to x^2 square centimeters. The length of the base of the triangle is $2x + 22$ centimeters, and the height of the triangle is $x - 10$ centimeters. What is the value of x?

ID: a58232b7 Question 34

The functions g and h are defined by the given equations, where $x \geq 0$. Which of the following equations displays, as a constant or coefficient, the minimum value of the function it defines, where $x \geq 0$?

I. $g(x) = 18(1.16)(1.4)^{x+2}$

II. $h(x) = 18(1.4)^{x+4}$

A. I only

B. II only

C. I and II

D. Neither I nor II

ID: a7711fe8 Question 35

What is the minimum value of the function f defined by $f(x)=(x-2)^2-4$?

A. -4

B. -2

C. 2

D. 4

ID: 1a722d7d Question 36

Let the function p be defined as $p(x)=\dfrac{(x-c)^2+160}{2c}$, where c is a constant. If $p(c)=10$, what is the value of $p(12)$?

A. 10.00

B. 10.25

C. 10.75

D. 11.00