Question ID: deff8a2f

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

The circumference of the base of a right circular cylinder is 20π meters, and the height of the cylinder is 6 meters. What is the volume, in cubic meters, of the cylinder?

Answer

A. 60π

B. 120π

C. 600π

D. $2{,}400\pi$

Correct Answer: C

Rationale

Choice C is correct. The volume, V, of a right circular cylinder is given by the formula $V = \pi r^2 h$, where r is the radius of the base of the cylinder and h is the height of the cylinder. It's given that a right circular cylinder has a height of 6 meters. Therefore, $h = 6$. It's also given that the right circular cylinder has a base with a circumference of 20π meters. The circumference, C, of a circle is given by $C = 2\pi r$, where r is the radius of the circle. Substituting 20π for C in the formula $C = 2\pi r$ yields $20\pi = 2\pi r$. Dividing each side of this equation by 2π yields $10 = r$. Substituting 10 for r and 6 for h in the formula $V = \pi r^2 h$ yields $V = \pi(10)^2(6)$, or $V = 600\pi$. Therefore, the volume, in cubic meters, of the cylinder is 600π.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect. This is the lateral surface area, not the volume, of the cylinder.

Choice D is incorrect. This is the result of using the diameter, not the radius, for the value of r in the formula $V = \pi r^2 h$.

Question ID: 502d9690

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

Rectangle $ABCD$ is similar to rectangle $EFGH$. The area of rectangle $ABCD$ is 648 square inches, and the area of rectangle $EFGH$ is 72 square inches. The length of the longest side of rectangle $ABCD$ is 36 inches. What is the length, in inches, of the longest side of rectangle $EFGH$?

Answer

A. 4

B. 9

C. 12

D. 36

Correct Answer: C

Rationale

Choice C is correct. It's given that rectangle $ABCD$ is similar to rectangle $EFGH$. Therefore, if the length of each side of rectangle $ABCD$ is k times the length of the corresponding side of rectangle $EFGH$, then the area of rectangle $ABCD$ is k^2 times the area of rectangle $EFGH$. It's given that the area of rectangle $ABCD$ is 648 square inches and the area of rectangle $EFGH$ is 72 square inches. It follows that $k^2 = \frac{648}{72}$, or $k^2 = 9$. Taking the square root of each side of this equation yields $k = \sqrt{9}$, or $k = 3$. It follows that the length of each side of rectangle $ABCD$ is 3 times the length of the corresponding side of rectangle $EFGH$. It's given that the length of the longest side of rectangle $ABCD$ is 36 inches. Therefore, 36 inches is 3 times the length of the longest side of rectangle $EFGH$, and the longest side of rectangle $EFGH$ is equal to $\frac{36}{3}$, or 12, inches.

Choice A is incorrect. This is the length, in inches, of the longest side of a rectangle with side lengths that are $\frac{1}{9}$ the corresponding side lengths of rectangle $ABCD$, rather than a rectangle with an area that is $\frac{1}{9}$ the area of rectangle $ABCD$.


Choice B is incorrect. This is the factor by which the area of rectangle $ABCD$ is larger than the area of rectangle $EFGH$, not the length, in inches, of the longest side of rectangle $EFGH$.

Choice D is incorrect. This is the length, in inches, of the longest side of rectangle $ABCD$, not rectangle $EFGH$.

Question ID: b2528e6b

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question



The three points shown define a circle. The circumference of this circle is $k\pi$, where k is a constant. What is the value of k?

Correct Answer: 8

Rationale

The correct answer is 8. It's given that the three points shown define a circle, so the center of that circle is an equal distance from each of the three points. The point $(6, 6)$ is halfway between the points $(6, 2)$ and $(6, 10)$, and is a distance of 4 units from each of those two points. The point $(6, 6)$ is also a distance of 4 units from $(2, 6)$. Because the point $(6, 6)$ is the same distance from all three points shown, it must be the center of the circle. Since that distance is 4, it follows that the radius of the circle is 4. The circumference of a circle with radius r is equal to $2\pi r$. It follows that the circumference of the given circle is $2\pi(4)$, or 8π. It's given that the circumference of the circle is $k\pi$. Therefore, the value of k is 8.

Question ID: 5b2b8866

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

A rectangular poster has an area of 360 square inches. A copy of the poster is made in which the length and width of the original poster are each increased by 20%. What is the area of the copy, in square inches?

Correct Answer: 2592/5, 518.4

Rationale

The correct answer is 518.4. It's given that the area of the original poster is 360 square inches. Let ℓ represent the length, in inches, of the original poster, and let w represent the width, in inches, of the original poster. Since the area of a rectangle is equal to its length times its width, it follows that $360 = \ell w$. It's also given that a copy of the poster is made in which the length and width of the original poster are each increased by 20%. It follows that the length of the copy is the length of the original poster plus 20% of the length of the original poster, which is equivalent to $\ell + \frac{20}{100}\ell$ inches. This length can be rewritten as $\ell + 0.2\ell$ inches, or 1.2ℓ inches. Similarly, the width of the copy is the width of the original poster plus 20% of the width of the original poster, which is equivalent to $w + \frac{20}{100}w$ inches. This width can be rewritten as $w + 0.2w$ inches, or $1.2w$ inches. Since the area of a rectangle is equal to its length times its width, it follows that the area, in square inches, of the copy is equal to $(1.2\ell)(1.2w)$, which can be rewritten as $(1.2)(1.2)(\ell w)$. Since $360 = \ell w$, the area, in square inches, of the copy can be found by substituting 360 for ℓw in the expression $(1.2)(1.2)(\ell w)$, which yields $(1.2)(1.2)(360)$, or 518.4. Therefore, the area of the copy, in square inches, is 518.4.

Question ID: 6708546e

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

Parallelogram $ABCD$ is similar to parallelogram $PQRS$. The length of each side of parallelogram $PQRS$ is 2 times the length of its corresponding side of parallelogram $ABCD$. The area of parallelogram $ABCD$ is 5 square centimeters. What is the area, in square centimeters, of parallelogram $PQRS$?

Answer

A. 7

B. 10

C. 20

D. 25

Correct Answer: C

Rationale

Choice C is correct. It's given that parallelogram $ABCD$ is similar to parallelogram $PQRS$. When two parallelograms are similar, if the scale factor between their corresponding side lengths is k, the scale factor between their areas is k^2. It's given that the length of each side of parallelogram $PQRS$ is 2 times the length of its corresponding side of parallelogram $ABCD$, so the scale factor between their corresponding side lengths is 2. It follows that the scale factor between their areas is 2^2, or 4. It's given that the area, in square centimeters, of parallelogram $ABCD$ is 5. It follows that the area, in square centimeters, of parallelogram $PQRS$ is $5(4)$, or 20.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question ID: b0dc920d

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

A manufacturer determined that right cylindrical containers with a height that is 4 inches longer than the radius offer the optimal number of containers to be displayed on a shelf. Which of the following expresses the volume, V, in cubic inches, of such containers, where r is the radius, in inches?

Answer

A. $V = 4\pi r^3$

B. $V = \pi(2r)^3$

C. $V = \pi r^2 + 4\pi r$

D. $V = \pi r^3 + 4\pi r^2$

Correct Answer: D

Rationale

Choice D is correct. The volume, V, of a right cylinder is given by the formula $V = \pi r^2 h$, where r represents the radius of the base of the cylinder and h represents the height. Since the height is 4 inches longer than the radius, the expression $r + 4$ represents the height of each cylindrical container. It follows that the volume of each container is represented by the equation $V = \pi r^2(r + 4)$. Distributing the expression πr^2 into each term in the parentheses yields $V = \pi r^3 + 4\pi r^2$.

Choice A is incorrect and may result from representing the height as $4r$ instead of $r + 4$. Choice B is incorrect and may result from representing the height as $2r$ instead of $r + 4$. Choice C is incorrect and may result from representing the volume of a right cylinder as $V = \pi rh$ instead of $V = \pi r^2 h$.

Question ID: 310c87fe

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

A cube has a surface area of 54 square meters. What is the volume, in cubic meters, of the cube?

Answer

A. 18

B. 27

C. 36

D. 81

Correct Answer: B

Rationale

Choice B is correct. The surface area of a cube with side length s is equal to $6s^2$. Since the surface area is given as 54 square meters, the equation $54 = 6s^2$ can be used to solve for s. Dividing both sides of the equation by 6 yields $9 = s^2$. Taking the square root of both sides of this equation yields $3 = s$ and $-3 = s$. Since the side length of a cube must be a positive value, $s = -3$ can be discarded as a possible solution, leaving $s = 3$. The volume of a cube with side length s is equal to s^3. Therefore, the volume of this cube, in cubic meters, is 3^3, or 27.

Choices A, C, and D are incorrect and may result from calculation errors.

Question ID: 459dd6c5

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

Triangles ABC and DEF are similar. Each side length of triangle ABC is 4 times the corresponding side length of triangle DEF. The area of triangle ABC is 270 square inches. What is the area, in square inches, of triangle DEF?

Correct Answer: 135/8, 16.87, 16.88

Rationale

The correct answer is $\frac{135}{8}$. It's given that triangles ABC and DEF are similar and each side length of triangle ABC is 4 times the corresponding side length of triangle DEF. For two similar triangles, if each side length of the first triangle is k times the corresponding side length of the second triangle, then the area of the first triangle is k^2 times the area of the second triangle. Therefore, the area of triangle ABC is 4^2, or 16, times the area of triangle DEF. It's given that the area of triangle ABC is 270 square inches. Let a represent the area, in square inches, of triangle DEF. It follows that 270 is 16 times a, or $270 = 16a$. Dividing both sides of this equation by 16 yields $\frac{270}{16} = a$, which is equivalent to $\frac{135}{8} = a$. Thus, the area, in square inches, of triangle DEF is $\frac{135}{8}$. Note that 135/8, 16.87, and 16.88 are examples of ways to enter a correct answer.

Question ID: c984f1a5

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

A hemisphere is half of a sphere. If a hemisphere has a radius of 27 inches, which of the following is closest to the volume, in cubic inches, of this hemisphere?

Answer

A. $1{,}500$

B. $6{,}100$

C. $30{,}900$

D. $41{,}200$

Correct Answer: D

Rationale

Choice D is correct. The volume, V, of a sphere is given by $V = \frac{4}{3}\pi r^3$, where r is the radius of the sphere. Since a hemisphere is half of a sphere, it follows that the volume, V, of a hemisphere is given by $V = \left(\frac{1}{2}\right)\left(\frac{4}{3}\right)\pi r^3$, or $V = \frac{2}{3}\pi r^3$. Substituting 27 for r in this formula yields $V = \frac{2}{3}\pi(27)^3$, which gives $V = 13{,}122\pi$, or V is approximately equal to $41{,}223.98$. Therefore, the choice that is closest to the volume, in cubic inches, of this hemisphere is $41{,}200$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question ID: f7e626b2

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

The dimensions of a right rectangular prism are 4 inches by 5 inches by 6 inches. What is the surface area, in square inches, of the prism?

Answer

A. 30

B. 74

C. 120

D. 148

Rationale


Choice D is correct. The surface area is found by summing the area of each face. A right rectangular prism consists of three pairs of congruent rectangles, so the surface area is found by multiplying the areas of three adjacent rectangles by 2 and adding these products. For this prism, the surface area is equal to $2(4\cdot5)+2(5\cdot6)+2(4\cdot6)$, or $2(20)+2(30)+2(24)$, which is equal to 148.

Choice A is incorrect. This is the area of one of the faces of the prism. Choice B is incorrect and may result from adding the areas of three adjacent rectangles without multiplying by 2. Choice C is incorrect. This is the volume, in cubic inches, of the prism.

Question ID: eb70d2d0

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question



What is the area, in square units, of the triangle formed by connecting the three points shown?

Correct Answer: 24.5, 49/2

Rationale

The correct answer is 24.5. It's given that a triangle is formed by connecting the three points shown, which are $(-3, 4)$, $(5, 3)$, and $(4, -3)$. Let this triangle be triangle A. The area of triangle A can be found by calculating the area of the rectangle that circumscribes it and subtracting the areas of the three triangles that are inside the rectangle but outside triangle A. The rectangle formed by the points $(-3, 4)$, $(5, 4)$, $(5, -3)$, and $(-3, -3)$ circumscribes triangle A. The width, in units, of this rectangle can be found by calculating the distance between the points $(5, 4)$ and $(5, -3)$. This distance is $4 - (-3)$, or 7. The length, in units, of this rectangle can be found by calculating the distance between the points $(5, 4)$ and $(-3, 4)$. This distance is $5 - (-3)$, or 8. It follows that the area, in square units, of the rectangle is $(7)(8)$, or 56. One of the triangles that lies inside the rectangle but outside triangle A is formed by the points $(-3, 4)$, $(5, 4)$, and $(5, 3)$. The length, in units, of a base of this triangle can be found by calculating the distance between the points $(5, 4)$ and $(5, 3)$. This distance is $4 - 3$, or 1. The corresponding height, in units, of this triangle can be found by calculating the distance between the points $(5, 4)$ and $(-3, 4)$. This distance is $5 - (-3)$, or 8. It follows that the area, in square units, of this triangle is $\frac{1}{2}(8)(1)$, or 4. A second triangle that lies inside the rectangle but outside triangle A is formed by the points $(4, -3)$, $(5, 3)$, and $(5, -3)$. The length, in units, of a base of this triangle can be found by calculating the distance between the points $(5, 3)$ and $(5, -3)$. This distance is $3 - (-3)$, or 6. The corresponding height, in units, of this triangle can be found by calculating the distance between the points $(5, -3)$ and $(4, -3)$. This distance is $5 - 4$, or 1. It follows that the area, in square units, of this triangle is $\frac{1}{2}(1)(6)$, or 3. The third triangle that lies inside the rectangle but outside triangle A is formed by the points $(-3, 4)$, $(-3, -3)$, and $(4, -3)$. The length, in units, of a base of this triangle can be found by calculating the distance between the points $(4, -3)$ and $(-3, -3)$. This distance is $4 - (-3)$, or 7. The corresponding height, in units, of this triangle can be found by calculating the distance between the points $(-3, 4)$ and $(-3, -3)$. This distance is $4 - (-3)$, or 7. It follows that the area, in square units, of this triangle is $\frac{1}{2}(7)(7)$, or 24.5. Thus, the area, in square

units, of the triangle formed by connecting the three points shown is $56 — 4 — 3 — 24.5$, or 24.5. Note that 24.5 and 49/2 are examples of ways to enter a correct answer.

Question ID: 93de3f84

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

The volume of right circular cylinder A is 22 cubic centimeters. What is the volume, in cubic centimeters, of a right circular cylinder with twice the radius and half the height of cylinder A?

Answer

A. 11

B. 22

C. 44

D. 66

Correct Answer: C

Rationale

Choice C is correct. The volume of right circular cylinder A is given by the expression $\pi r^2 h$, where r is the radius of its circular base and h is its height. The volume of a cylinder with twice the radius and half the height of cylinder A is given by $\pi(2r)^2\left(\frac{1}{2}\right)h$, which is equivalent to $4\pi r^2\left(\frac{1}{2}\right)h = 2\pi r^2 h$. Therefore, the volume is twice the volume of cylinder A, or $2 \times 22 = 44$.

Choice A is incorrect and likely results from not multiplying the radius of cylinder A by 2. Choice B is incorrect and likely results from not squaring the 2 in 2r when applying the volume formula. Choice D is incorrect and likely results from a conceptual error.

Question ID: dc71597b

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Geometry and Trigonometry	Area and volume	Hard

Question

A right circular cone has a volume of $\frac{1}{3}\pi$ cubic feet and a height of 9 feet. What is the radius, in feet, of the base of the cone?

Answer

A. $\frac{1}{3}$

B. $\frac{1}{\sqrt{3}}$

C. $\sqrt{3}$

D. 3

Correct Answer: A

Rationale

Choice A is correct. The equation for the volume of a right circular cone is $V = \frac{1}{3}\pi r^2 h$. It's given that the volume of the right circular cone is $\frac{1}{3}\pi$ cubic feet and the height is 9 feet. Substituting these values for V and h, respectively, gives $\frac{1}{3}\pi = \frac{1}{3}\pi r^2(9)$. Dividing both sides of the equation by $\frac{1}{3}\pi$ gives $1 = r^2(9)$. Dividing both sides of the equation by 9 gives $\frac{1}{9} = r^2$. Taking the square root of both sides results in two possible values for the radius, $\sqrt{\left(\frac{1}{9}\right)}$ or $-\sqrt{\left(\frac{1}{9}\right)}$. Since the radius can't have a negative value, that leaves $\sqrt{\left(\frac{1}{9}\right)}$ as the only possibility. Applying the quotient property of square roots, $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$, results in $r = \frac{\sqrt{1}}{\sqrt{9}}$, or $r = \frac{1}{3}$.

Choices B and C are incorrect and may result from incorrectly evaluating $\sqrt{\left(\frac{1}{9}\right)}$. Choice D is incorrect and may result from solving $r^2 = 9$ instead of $r^2 = \frac{1}{9}$.