

MEMO NAME: GFAU_MATHEMATICAL_BACKGROUND

DATE: December 16, 2017

TO: EFC LaBerge

SUBJECT: Mathematical Background on the Galois Field Arithmetic Unit

1 Background

A Galois Field is a field with a finite number of elements. The nomenclature $GF(q)$ is used to indicate a Galois Field with q elements. In $GF(q)$, the parameter q must be a power of a prime. For each prime power there exists exactly one finite field. The binary field $GF(2)$ is the most frequently used Galois field. (1)

1.1 Purpose and Scope

The Galois Field Arithmetic Unit operates in fields of q^n , where $2 \leq n \leq 16$. This document will demonstrate the functionality of the unit by deriving all its functionality through mathematical approaches. The mathematical algorithms will be followed by their corresponding implementations in digital design.

1.2 Terms and Keywords

1.2.1 Input Primitive Polynomials

Input primitive polynomials in the Galois Field are represented as:

$$c_n x^n + \dots + c_2 x^2 + c_1 x^1 + c_0 x^0, \text{ where } c, x \in GF(2) = \{0, 1\}$$

For convenience and simplicity, all the examples provided will refer to the following polynomial: $x^3 + x^2 + x^0$.

1.2.2 Elements

The elements of an input polynomial refer to the $2^n - 1$ elements in the field.

1.2.3 Polynomial Form

The polynomial forms of the elements refer to the $2^n - 1$ symbolic representations of the of an input primitive polynomials in the field.

1.2.4 Example

An example of the elements and their corresponding polynomials is provided below:

Table 1: The 8 Element Vectors of $x^3 + x^2 + x^0$ in $GF(2)[x]$

Element	Symbol	Polynomial Form	Symbol
0	(1)	$0 + 0 + 0$	000
β^0	000	$0 + 0 + \beta^0$	001
β^1	001	$0 + \beta^1 + 0$	010
β^2	010	$\beta^2 + 0 + 0$	100
β^3	011	$\beta^2 + 0 + \beta^0$	101
β^4	100	$\beta^2 + \beta^1 + \beta^0$	111
β^5	101	$0 + \beta^1 + \beta^0$	011
β^6	110	$\beta^2 + \beta^1 + 0$	110
β^7	(2)	$0 + 0 + \beta^0$	001

(1) Zero is a reserved element where its binary symbol does not represent its decimal value.

(2) Elements beyond the $(2^n - 1)$ th element will be handled with special conditions since they cycle back to previous elements.

1.2.5 Notations

Several notations are used in this document to assist in linking the mathematical and digital design concepts. A list of selected notations are provided below:

Table 2: Mathematical and Logical Notations

Notation	Definition
$a + b$	Arithmetic Addition of a and b
$a - b$	Arithmetic Subtraction of a and b
$a \cdot b$	Arithmetic Multiplication of a and b
a / b	Arithmetic Division of a and b
$a \wedge b$	Logical AND of a and b
$a \vee b$	Logical OR of a and b
$a \oplus b$	Logical XOR of a and b
$a \ll b$	Logical Shift-Left a by b Bits
\bar{a}	Bitwise Inversion of a
$a_{2's}$	Two's Complement of a
$ a $	Number of Bits in a
$\langle x_n, x_{n-1}, \dots x_0 \rangle$	Ordered Array of Size n
$\{x_n, x_{n-1}, \dots x_0\}$	Unordered Set of Size n
$\langle x_m, \overleftarrow{0_{m-1}, \dots 0_n}, x_{n-1}, \dots x_0 \rangle$	Zero-Padding Between the $(m - 1)$ th and n th Bits
$A[i]$	i th Bit of Array A
$A[i : j]$	Subarray of Array A from its i th to j th index inclusive; where $i > j$

2 Algorithms

Input polynomials will be represented as 16 bit zero- base arrays. For example, the polynomial $x^3 + x^2 + x^0$ will be represented as

$$< 0000\ 0000\ 0000\ 1101 > \text{ (3rd, 2nd, and 0th bits)}$$

2.1 Determining Irreducibility

A polynomial is said to be irreducible if and only if there exists no roots for it.

2.1.1 Analytical Approach

If the sum of the coefficients of the polynomial equals 1 when $x = 0$ and $x = 1$, the polynomial is irreducible.

2.1.1.1 Example

Show that the polynomial $x^3 + x^2 + x^0$ is irreducible in $GF(2)[x]$ using the mathematical algorithm:

$$x = 0 : 0^3 + 0^2 + 0^0$$

$$= 0 + 0 + 1$$

$$= 1$$

$$x = 1 : 1^3 + 1^2 + 1^0$$

$$= 1 + 1 + 1$$

$$= 1$$

$$\therefore x^3 + x^2 + x^0 \text{ has no roots in } GF(2)[x].$$

Since the sum of the coefficients is 1 for both the cases, the polynomial is therefore determined irreducible.

2.1.2 Digital Logic

To design such algorithm, the polynomial may be checked for the following attributes:

1. the coefficient of its 0th term is 1
2. the total number of non-zero coefficients is odd

If both of these conditions are met, the polynomial is irreducible.

2.1.2.1 Example

Show that the polynomial $x^3 + x^2 + x^0$ is irreducible using the digital algorithm:

$$< 0000\ 0000\ 0000\ 1101 >$$

The 0th bit is 1 and the total number of non-zero coefficients is 3. Therefore, the polynomial is determined irreducible.

2.2 Symbols

Once a polynomial is determined irreducible, its symbols may be generated. The number of terms grow exponentially, $2^n - 1$, where n is the highest degree of the polynomial.

2.2.1 Default Symbols

Default symbols refer to terms in the field that exist for Galois Fields of all irreducible polynomials of q^n , where $2 \leq n \leq 16$. Since the number of elements cannot be smaller than 2, only zero and the 0th and 1st elements are shared among all fields.

Table 3 have all the bits of their values set to 0 except where indicated.

2.2.1.1 Analytical Approach

Using the example polynomial $x^3 + x^2 + x^0$, show that 0, the 0th element and the first element exist in $GF(2^3)$.

Table 3: Default Symbols Generated for All Irreducible Polynomials

Element	Polynomial Form	Symbol
0	$0_{15} + \dots + 0_2 + 0_1 + 0_0$	$< \overleftarrow{0_{15} \dots 0_2 0_1 0_0} >$
α^0	$0_{15} + \dots + 0_2 + 0_1 + \alpha_0^0$	$< \overleftarrow{0_{15} \dots 0_2 0_1 1_0} >$
α^1	$0_{15} + \dots + 0_2 + \alpha_1^1 + 0$	$< \overleftarrow{0_{15} \dots 0_2 1_1 0_0} >$

Let $\beta \in GF(2^3)$ be a root of $x^3 + x^2 + x^0 \implies \beta^3 + \beta^2 + \beta^0$

\therefore The coefficients are in $GF(2) \implies \beta^3 = \beta^2 + \beta^0$

\therefore a field has additive and multiplicative identities:

$$\therefore \{0, 1 = \beta^0\} \in GF(2^3)$$

$$\therefore \beta^1 \in GF(2^3) (\because \text{closure of multiplication})$$

2.2.2 Automatic Symbols

Automatic symbols refer to terms up to x^{n-1} . Automatic symbols may be generated concurrently, and consist of the following attributes:

1. The symbols for $\{x^0, x^1, \dots, x^{n-1}\}$ are generated by setting the corresponding bits to 1.
2. The symbol for x^n is generated by setting the corresponding bits for the terms in the polynomial after the highest degree term.
3. The symbol for x^{2^n-1} cycles back to x^0 , and is set to x^0 .

Automatic symbols consist of $n+1$ terms. Therefore, the maximum of 16 bits would have 17 terms generated by default.

Table 4: Automatic Symbols Generated for An Irreducible Polynomial of Degree $n < 16$. For $n = 16$, the most significant bit will be 1_{15} .

Element	Polynomial Form	Symbol
α^2	$0_{15} + \dots + 0_{n-1} + \dots + \alpha_2^2 + 0_1 + 0_0$	$< \overleftarrow{0_{15} \dots 0_{n-1}} \dots 1_2 0_1 0_0 >$
\dots	\dots	\dots
α^{n-1}	$0_{15} + \dots + \alpha_{n-1}^{n-1} + \dots + 0_2 + 0_1 + 0_0$	$< 0_{15} \dots 1_{n-1} \dots 0_2 0_1 0_0 >$
α^n	$0_{15} + \dots + \alpha_{n-1}^{n-1} + \dots + \alpha_2^2 + \alpha_1^1 + \alpha_0^0$	$< 0_{15} \dots x_{n-1} \dots x_2 x_1 x_0 >$
α^{2^n-1}	$0_{15} + \dots + 0_{n-1} + \dots + \alpha_2^2 + 0_1 + 0_0$	$< \overleftarrow{0_{15} \dots 0_2} 0_1 0_0 >$

2.2.2.1 Analytical Approach

Assuming the preceding values exist from the proof above, use the example polynomial $x^3 + x^2 + x^0$ to show that the $n - 1$ th and the n th elements exist in $GF(2^3)$.

$$\text{Let } \beta \in GF(2^3) \text{ be a root of } x^3 + x^2 + x^0 \implies \beta^3 + \beta^2 + \beta^0$$

$$\therefore \text{The coefficients are in } GF(2) \implies \beta^3 = \beta^2 + \beta^0$$

$$\therefore \beta^2 \in GF(2^3) (\because \text{closure of multiplication})$$

$$\therefore \beta^3 \in GF(2^3) (\because \beta^3 = \beta^2 + \beta^0)$$

2.2.3 Generated Symbols

The rest of the symbols for the elements x^{n+1} to x^{2^n-2} must be generated. In total, that would require $2^n - 2 - n - 1 + 1 = 2^n - 2 - n$ terms. Therefore, the maximum of 16 bits would require 65,518 terms to be generated.

2.2.3.1 Analytical Approach

Assuming the preceding values exist from the proof above, use the example polyno-

mial $x^3 + x^2 + x^0$ to show that the elements up to the (2^{n-2}) th elements exist in $GF(2^3)$.

$$\text{Let } \beta \in GF(2^3) \text{ be a root of } x^3 + x^2 + x^0 \implies \beta^3 + \beta^2 + \beta^0$$

$$\therefore \text{The coefficients are in } GF(2) \implies \beta^3 = \beta^2 + \beta^0$$

$$\begin{aligned} \therefore \beta^4 &= \beta^1 \times \beta^3 \\ &= \beta^1(\beta^2 + \beta^0) \\ &= \beta^3 + \beta^1 \\ &= \beta^2 + \beta^1 + \beta^0 \end{aligned}$$

$$\therefore \beta^4 \in GF(2^3)$$

$$\begin{aligned} \therefore \beta^5 &= \beta^1 \times \beta^4 \\ &= \beta^1(\beta^2 + \beta^1 + \beta^0) \\ &= \beta^3 + \beta^2 + \beta^1 \\ &= \beta^2 + \beta^0 + \beta^2 + \beta^1 \\ &= \beta^1 + \beta^0 \end{aligned}$$

$$\therefore \beta^5 \in GF(2^3)$$

$$\begin{aligned} \therefore \beta^6 &= \beta^1 \times \beta^5 \\ &= \beta^1(\beta^1 + \beta^0) \\ &= \beta^2 + \beta^1 \end{aligned}$$

$$\therefore \beta^6 \in GF(2^3)$$

$$\begin{aligned} \therefore \beta^7 &= \beta^1 \times \beta^6 \\ &= \beta^1(\beta^2 + \beta^1) \\ &= \beta^3 + \beta^2 \\ &= \beta^2 + \beta^0 + \beta^2 \\ &= \beta^0 = 1 \end{aligned}$$

$$\therefore \beta^7 \in GF(2^3)$$

2.2.3.2 Digital Logic

ELABORATE MORE Generating the rest of the symbols may be implemented with a linear feedback shift register (LFSR), using the following recursive equation:

$$\alpha^{n+m} = \alpha^{n+(m-1)} \times \alpha^n$$

$$= \begin{cases} (\alpha^{n+(m-1)} \ll 1)[n-2:0] & \text{if } (\alpha^{n+(m-1)} \ll 1)[n-1] = 0 \\ (\alpha^{n+(m-1)} \ll 1)[n-2:0] \oplus \alpha^n[n-2:0] & \text{if } (\alpha^{n+(m-1)} \ll 1)[n-1] = 1 \end{cases}$$

2.3 Operations

Operations in the Galois Field Arithmetic Unit consist of addition, subtraction, multiplication, division and logarithm.

2.3.1 Addition and Subtraction

Binary addition and binary subtraction are synonymous in the Galois Field. Addition and subtraction of Galois operands may be done by bitwise exclusive OR-ing the operands.

$$\begin{aligned} \alpha^i \pm \alpha^j &= \{x_{i,n-1}, \dots, x_{i,2}, x_{i,1}, x_{i,0}\} + \{x_{j,n-1}, \dots, x_{j,2}, x_{j,1}, x_{j,0}\} \\ &= \{(x_{i,n-1} \oplus x_{j,n-1}), \dots, (x_{i,2} \oplus x_{j,2}), (x_{i,1} \oplus x_{j,1}), (x_{i,0} \oplus x_{j,0})\} \\ &= \alpha^k \end{aligned}$$

2.3.1.1 Digital Design

Only polynomial forms of inputs are valid for these operations.

The implementation of Galois addition and subtraction may be computed with a single-level parallel array of XOR gates.

2.3.2 Multiplication

Binary multiplication of Galois operands is congruent to the sum of the indices of the operands. If the indices sum to greater than or equal to $2^n - 1$, then $2^n - 1$ is subtracted

from the sum to prevent overflow.

$$\begin{aligned}
\alpha^i \cdot \alpha^j &= \{x_{i,n-1}, \dots, x_{i,2}, x_{i,1}, x_{i,0}\} \cdot \{x_{j,n-1}, \dots, x_{j,2}, x_{j,1}, x_{j,0}\} \\
&= \alpha^{(i+j) \bmod (2^n-1)} \\
&= \begin{cases} \alpha^{(i+j)-(2^n-1)} & \text{if } (i+j) \geq 2^n - 1 \\ \alpha^{(i+j)} & \text{if } (i+j) < 2^n - 1 \end{cases}
\end{aligned}$$

2.3.2.1 Digital Design

Only element forms of inputs are valid for this operation.

Galois multiplication requires multiple binary additions and condition checks. To find the product of α^i and α^j , a carry-lookahead adder may be used to sum the elements $i+j$. The most significant bits of the sum and the sum +1 will then be OR-ed to compute a single-bit control signal. The control signal will be multiplexed with a binary 1 to be added to the sum.

$$\begin{aligned}
\alpha^i \cdot \alpha^j &= \{x_{i,n-1}, \dots, x_{i,2}, x_{i,1}, x_{i,0}\} \cdot \{x_{j,n-1}, \dots, x_{j,2}, x_{j,1}, x_{j,0}\} \\
&\implies \text{Let } n = |i| = |j| \\
&= \begin{cases} \alpha^{(i+j)} & \text{if } ((i+j)[n+1] \vee (i+j+1)[n+1]) = 0 \\ \alpha^{(i+j+1)} & \text{if } ((i+j)[n+1] \vee (i+j+1)[n+1]) = 1 \end{cases}
\end{aligned}$$

2.3.3 Division

Binary division of Galois operands is congruent to the difference of the indices of the operands. If the difference is negative, then the absolute value of the difference is subtracted from $2^n - 1$ to prevent underflow. If the difference is zero, then the quotient

is α^0 .

$$\begin{aligned}\alpha^i / \alpha^j &= \{x_{i,n-1}, \dots, x_{i,2}, x_{i,1}, x_{i,0}\} / \{x_{j,n-1}, \dots, x_{j,2}, x_{j,1}, x_{j,0}\} \\ &= \alpha^{(i-j) \bmod (2^n-1)} \\ &= \begin{cases} \alpha^{(2^n-1)-(j-i)} & \text{if } (i-j) < 0 \\ \alpha^{(i-j)} & \text{if } (i-j) > 0 \\ \alpha^0 & \text{if } i = j, i \neq 0 \\ ERROR & \text{if } j = 0 \end{cases}\end{aligned}$$

2.3.3.1 Digital Design

Only element forms of inputs are valid for this operation.

Galois division requires multiple binary additions and condition checks. To find the quotient of α^i and α^j , the two's complement of j has to first be summed with i . Converting to two's complement may be done with parallel inverters and a carry-lookahead adder. The overflow bit of the sum will then be used as a control signal to a multiplexer. If activated, the multiplexer will add the two's complement of a binary 1 to the sum.

$$\begin{aligned}\alpha^i / \alpha^j &= \{x_{i,n-1}, \dots, x_{i,2}, x_{i,1}, x_{i,0}\} / \{x_{j,n-1}, \dots, x_{j,2}, x_{j,1}, x_{j,0}\} \\ &\implies \text{Let } n = |i| = |j| \\ &= \begin{cases} \alpha^{(i+j_{2's})} & \text{if } (i + j_{2's})[n+1] = 1 \\ \alpha^{(i+j_{2's}+1_{2's})} & \text{if } (i + j_{2's})[n+1] = 0 \end{cases}\end{aligned}$$

2.3.4 Logarithm

Logarithm is considered a unary operation in the Galois Field, where only one operand is required. The base of the logarithm operation in the Galois Field is implicitly 2. The logarithm of a Galois operand is congruent to its index.

$$\log_{GF(2)}(\alpha^i) = i$$

2.3.4.1 Digital Design

Only element forms of inputs are valid for this operation.

2.4 Arithmetic Exceptions

2.4.1 Zero

2.4.2 “Infinity”

References

- (1) Weisstein, Eric W. "Finite Field." From *MathWorld*—A Wolfram Web Resource.
<http://mathworld.wolfram.com/FiniteField.html>