605.611 - Foundations of Computer Architecture

Assignment 02

Sabbir Ahmed

February 9, 2021

- 1. Convert the following fixed point numbers to binary fixed point. Give both the actual values, and normalize the values so that they have a binary 1 as the value for the left of the decimal point.
 - (a) 7.25

Answer: Integral part: $7 = (0111)_2$

Repeatedly multiplying the fractional part by 2:

(b) 13.5

Answer: Integral part: $13 = (1101)_2$

Repeatedly multiplying the fractional part by 2:

0.50
$$= \frac{0.50}{2} = \frac{\times 2}{1.000} = (1)_2$$

$$= (1)_2 = 3.5 = (1101.1)_2, (normalized to $(1.1011)_2 \times 2^3$)$$

(c) 0.5625

Answer: Integral part: 0

Repeatedly multiplying the fractional part by 2:

(d) 0.125

Answer: Integral part: 0

Repeatedly multiplying the fractional part by 2:

$$0.125$$

$$0.125$$

$$\times 2$$

$$0.250$$

$$= \times 2$$

$$0.500$$

$$\times 2$$

$$1.000$$

$$\times 2$$

$$1.000$$

$$\times 2$$

$$0.5625 = (0.001)_2, (normalized to $(1.0)_2 \times 2^{-3}$)$$

(e) 127.625

0.625

 $=(101)_2$

Answer: Integral part: 127 = (011111111)

Repeatedly multiplying the fractional part by 2:

(f) 51,025.025

 \Rightarrow 127.625 = (1111111.101)2, (normalized to (1.111111101)2 \times $2^6)$

Answer: Integral part: 51025 =

 $\Rightarrow 51025$

$$= 2) \frac{25512}{51025} \qquad \frac{12756}{2)25512} \qquad \frac{6378}{2)12756} \qquad \frac{3189}{6378} \qquad \frac{1594}{2)3189} \qquad \frac{797}{2)1594}$$

$$= 2) \frac{4}{51025} \qquad \frac{2}{2)25512} \qquad \frac{12}{12} \qquad \frac{6}{6} \qquad \frac{2}{2} \qquad \frac{14}{2}$$

$$= 11 \qquad 05 \qquad 07 \qquad 03 \qquad 11 \qquad 19$$

$$= 10 \qquad \frac{4}{7} \qquad \frac{6}{7} \qquad \frac{2}{7} \qquad \frac{10}{18} \qquad \frac{18}{14}$$

$$= 10 \qquad 15 \qquad 15 \qquad 17 \qquad 18 \qquad 14$$

$$= 10 \qquad 14 \qquad 14 \qquad 16 \qquad 18 \qquad 09 \qquad 0$$

$$= \frac{2}{7} \qquad \frac{10}{7} \qquad \frac{16}{7} \qquad \frac{18}{7} \qquad \frac{8}{7} \qquad 0$$

$$= \frac{2}{7} \qquad \frac{10}{7} \qquad \frac{16}{7} \qquad \frac{18}{7} \qquad \frac{8}{7} \qquad 0$$

$$= \frac{4}{7} \qquad \frac{12}{7} \qquad 0$$

$$\begin{array}{ccc}
\frac{1}{2} & 0 \\
2 & 1
\end{array}$$

1

 \Rightarrow (1100 0111 0101 0001)₂

0

Repeatedly multiplying the fractional part by 2:

0.025

The pattern $(0000011)_2$ appears to keep repeating $\Rightarrow 51,025.025 = (1100011101010001.0000011)_2,$ (normalized to $(1.1000111010100010000011)_2 \times 2^{15}$)

(g) 7.1

Answer: Integral part: $7 = (111)_2$

Repeatedly multiplying the fractional part by 2:

The pattern $(0011)_2$ appears to keep repeating

$$\Rightarrow 7.1 = (111.0\overline{00110011})_2,$$

(normalized to $(1.110\overline{00110011})_2\times 2^2)$

(h) 5.2

Answer: Integral part: $5 = (101)_2$

Repeatedly multiplying the fractional part by 2:

The pattern $(0011)_2$ appears to keep repeating

$$\Rightarrow 7.1 = (101.\overline{00110011})_2,$$

(normalized to $(1.01\overline{00110011})_2\times 2^2)$

- 5. Convert the following from decimal to excess 127 format. Write your answers as hexadecimal digits.
 - (a) -4

Answer:

$$-4 + 127$$
$$= 123 = 7B_{16}$$

(b) 4

Answer:

$$4 + 127$$

$$= 131 = 83_{16}$$

(d) 7

Answer:

$$7 + 127$$
$$= 134 = 86_{16}$$

(e) -7

Answer:

$$-7 + 127$$
$$= 120 = 78_{16}$$

- 8. Single precision floating point numbers have 7 digit decimal precision and double floating point numbers have 15 digit precision. Explain how these precision values are arrived at, and what they mean.
- 9. Convert the following numbers to IEEE 754 single precision numbers. Give your answers as hexadecimal numbers (do not give me binary, I cannot read it accurately. I WILL misread it and you WILL lose points).

(a) 7.25

Answer: Since the decimal is positive, the sign bit is 0

$$7.25 = (1.1101)_2 \times 2^2$$
 from Part 1a

Mantissa: $(1101)_2$

Exponent: +2

$$2 + 127$$

= $(1000\ 0001)_2$
 $\Rightarrow (0100\ 0000\ 1110\ 1000)_2 = (40E8\ 0000)_{16}$

(b) 13.5

Answer: Since the decimal is positive, the sign bit is 0

$$13.5 = (1.1011)_2 \times 2^3$$
 from Part 1b

Mantissa: $(1011)_2$

Exponent: +3

$$3 + 127$$

= $(1000\ 0010)_2$
 $\Rightarrow (0100\ 0001\ 0101\ 1000)_2 = (4158\ 0000)_{16}$

(c) 0.5625

Answer: Since the decimal is positive, the sign bit is 0

$$0.5625 = (1.001)_2 \times 2^{-1}$$
 from Part 1c

Mantissa: $(001)_2$

Exponent: -1

$$-1 + 127$$

= $(0111 \ 1110)_2$
 $\Rightarrow (0011 \ 1111 \ 0001 \ 0000)_2 = (3F10 \ 0000)_{16}$

(d) 0.125

Answer: Since the decimal is positive, the sign bit is 0

$$0.125 = (1.0)_2 \times 2^{-3}$$
 from Part 1d

Mantissa: 0

Exponent: -3

$$-3 + 127$$

= $(0111 \ 1100)_2$
 $\Rightarrow (0011 \ 1110 \ 0000 \ 0000)_2 = (3E00 \ 0000)_{16}$

(e) 127.625

Answer: Since the decimal is positive, the sign bit is 0

$$127.625 = (1.1111111101)_2 \times 2^6$$
 from Part 1e

Mantissa: $(1111111101)_2$

Exponent: 6

$$6 + 127$$

= $(1000 \ 0101)_2$
 $\Rightarrow (0100 \ 0010 \ 1111 \ 1111 \ 0100)_2 = (42FF \ 4000)_{16}$

(f) 51025.025

Answer: Since the decimal is positive, the sign bit is 0

$$51025.025 = (1.1000111010100010000011)_2 \times 2^{15}$$
 from Part 1f

Mantissa: $(1000111010100010000011)_2$


Exponent: 15

$$15 + 127$$

= $(1000 \ 1110)_2$
 $\Rightarrow (0100 \ 0111 \ 0100 \ 0111 \ 0101 \ 0001 \ 0000 \ 0110)_2 = (4747 \ 5106)_{16}$

- 10. For each of the following truth tables:
 - (a) Give the DNF equation for the table.
 - (b) Give the minimal equation.
 - (c) Using Boolean algebra show the two Boolean equations are equivalent.
 - (d) Draw the circuit in Logisim. Be prepared to draw the circuit by hand.

A	В	C	f(A,B,C)
0	0	0	0
0	0	1	1
0	1	0	1
0	1	1	0
1	0	0	0
1	0	1	1
1	1	0	1
1	1	1	0



A	В	C	f(A,B,C)
0	0	0	1
0	0	1	1
0	1	0	1
0	1	1	1
1	0	0	0
1	0	1	1
1	1	0	0
1	1	1	1

A	В	C	f(A,B,C)
0	0	0	1
0	0	1	0
0	1	0	0
0	1	1	1
1	0	0	1
1	0	1	1
1	1	0	1
1	1	1	1