

Latex Template

Víctor Alcázar

May 31, 2017

Contents

1	Statement	1
2	Decisional Problem	3

1 Statement

We've been given the following problem to talk about.


MINIMUM CUT LINEAR ARRANGEMENT

Given a graph $G = (V, E)$, compute a one-to-one function $f : V \rightarrow [1..|V|]$ so that the maximum number of cut edges in any integer point is minimised, i.e.

$$\max_{i \in 1..|V|} |\{ \{u, v\} \in E : f(u) \leq i < f(v) \} |$$


i. Example Instances

In order to visualize the problem and attack it using more intuition, we must consider that the cuts at integer points can be seen easily in the following manner. Given a graph G and an ordering function f , we draw the nodes in an horizontal line, in the order given by f . What we are trying to minimize is the size maximum cut among the S_i 's, where $S_i = \{v \in V : f(v) \leq i\}$, for $i = 1, 2, \dots, n-1$. But on the graph drawing mentioned above, the size of every S_i is the number of edges existing in the space between the vertical lines passing by nodes i and $i+1$. To give an example, in the following linear arrangement we easily deduce that the minimum S_i size is 3:



TODO: draw another example

The problem instances are Graphs G . As an example, we take $G = K_5$:




Here, every node is connected to every one of the rest. Therefore, every node is identical to every one else. In this case, every assignment of numbers $1 \dots n$ to the nodes yields the same graph. Therefore to solve this instance, we consider a random labeling of our nodes, and we compute the cuts at all integer points between 1 and n .

Another instance, could be this graph.



Here, the assignment matters. In order to attack the problem in a more intuitive manner, it would be more suitable to have visualize the problem as follows: Given graph G , create a drawing of G such that the nodes are placed on a straight horizontal line. The objective is to minimize the maximum number of edges that exist in the space between two consecutive nodes. For example, the graph below is a visualization of the assignment $f(1) = 1, f(2) = 4, f(3) = 2, f(4) = 3$, which gives us $\max(S_i) = 3$.



By the way, the above assignment gives us the best possible results, w.r.t. the requirements of the problems. That is, there is no assignment that gives us a lower maximum cutwidth. **explain better here**

2 Decisional Problem

Given a graph $G = (V, E)$ and $k \in \mathbb{N}$, say if there exists one ordering of the vertices so that the maximum number of cut edges in any integer point is less than or equal k , i.e.

$$\exists f : V \rightarrow [1..|V|] \max_{i \in [1..|V|]} |\{ \{u, v\} \in E : f(u) \leq i < f(v) \}| \leq k$$

Problem

iii. Parametrized Problems

Using this problem, and combining it with a computable function $\kappa : \Sigma^* \rightarrow \mathbb{N}$ we can formulate a Parametrized version of this problem.

TODO: Add Natural numbers symbol

An example:

Input	Graph G and $tw = \text{treewidth}(G)$
Parameters	tw
Question	What is the Min-Cut Linear Arrangement of G?

This problem can be solved in linear time, as the problem is expressible in Monadic Second Order Logic /citetrans.

TODO: Confirm the above statment!!

References

- [1] Maria Serna, *Class transparencies*, Fall 2017