

COMBINING NEURAL NETWORKS AND
GEOSTATISTICS FOR LANDSLIDE HAZARD
ASSESSMENT OF SAN SALVADOR
METROPOLITAN AREA, EL SALVADOR

COMBINANDO REDES NEURONALES Y
GEOESTADÍSTICA PARA EVALUACIÓN DE
DESLIZAMIENTOS DE TIERRA DEL ÁREA
METROPOLITANA DE SAN SALVADOR, EL
SALVADOR

Ricardo Ríos ^{*1}, Alexandre Ribó ^{**2}, Roberto Mejía ^{***3}, and
Giovanni Molina ^{****4}

¹Department of Mathematics, Science and Mathematics Faculty,
University of El Salvador, El Salvador.

²National Institute of Health, Ministry of Health of El Salvador, El
Salvador.

³National Institute of Health, Ministry of Health of El Salvador, El
Salvador.

⁴Ministry of Environment and Natural Resources of El Salvador.

5 de junio de 2015

Resumen

Esta contribución describe la creación de un modelo de evaluación de deslizamiento de tierra para el Área Metropolitana de San Salvador, departamento de El Salvador. El análisis inicio con la obtención de una foto área del MARN (Ministerio de Medio Ambiente y Recursos Naturales) con un total de 939407 puntos georeferenciados con el fin de producir un inventario de deslizamiento. En esta evaluación de los deslizamientos se uso 4792 eventos previamente foto-interpretados y 7 factores condicionantes incluyendo: geomorfología, geología, precipitaciones máximas, aceleraciones sísmicas, pendiente del terreno, distancia a carretera y falla geológica. Redes Neuronales Artificiales (RNA) fueron usadas para la evaluación de la susceptibilidad a deslizamiento de tierra, logrando que más del 80 % de deslizamientos fueran apropiadamente clasificados usando un criterio dentro y fuera de la muestra con la que se estimaron los parámetros del modelo. Regresión Logística fue usada como base de comparación, obteniendo este modelo un rendimiento inferior que el de RNA con un porcentaje de correcta clasificación abajo del 70 %. Para completar el análisis se realizo la interpolación de puntos usando el método kriging proveniente del enfoque geoestadístico. Finalmente, los resultados muestran que es posible obtener un mapa de riesgo a deslizamiento de tierra, haciendo uso de una combinación de RNA y técnicas geoestadísticas con lo cual la presente investigación puede ayudar a la mitigación de deslizamientos de tierra en El Salvador.

Palabras clave: deslizamiento de tierra, evaluación de riesgo, El Salvador, RNA, geoestadística.

Abstract

This contribution describes the creation of a landslide hazard assessment model for San Salvador, department in El Salvador. The analysis started with an aerial photointerpretation from Ministry of Environment and Natural Resources of El Salvador (MARN Spanish acronym) with a total amount of 939407 georeferenced points to produce a landslide inventory. In this landslide assessment we have used 4792 events previously photo-interpretated and 7 conditioning factors including: geomorphology, geology, rainfall intensity, peak ground acceleration, slope angle, road and fault distance. Artificial Neural Networks (ANN) were applied for the assessment of susceptibility to landslides, achieving more than 80 % of landslide were properly classified using in-sample and out of sample criteria. Logistic regression was used as base of comparison, obtaining this

* ricardo.sv@gmail.com

** alexandre4rt@gmail.com

*** robertomejia1685@gmail.com

**** giova.molina@gmail.com

model a performance lower than ANNs with a percentage of correct classification under 70 %. To complete the analysis we have performed interpolation of the points using kriging method from geostatistical approach. Finally, the results show that is possible to derive a landslide hazard map, making use of a combination of ANNs and geostatistical techniques wherewith the present study can help landslide mitigation in El Salvador.

Keywords: landslide, hazard assessment, El Salvador, ANN, geostatistics, Artificial Neural Networks, kriging.

Mathematics Subject Classification: 62P12.

1. Introduction

El Salvador, one of the smallest and most crowded nations in Central America, extends in Pacific coast about 240 km westward from the Gulf of Fonseca to the border with Guatemala (see Fig. 1). El Salvador borders an active subduction boundary between Cocos and Caribbean plates located 30 km offshore. Therefore El Salvador is affected with high seismicity and volcanic activity related to this active boundary. There is two main sources of seismic activity, the upper interface thrust that coincides with the location of the recent volcanoes and the Benioff-Wadaty zones of subducted Cocos plate where deeper intraplate earthquakes occurs ([1]). Due this tectonic and volcanic activity El Salvador has rugged relief. The main ranges cross the country with a rough west-east trend, parallel to the coast, these are separated from each other by faults and grabens. These ranges present several highly active volcanoes. The surface geology in El Salvador is almost entirely volcanic, dominated by upper Tertiary to Holocene volcanic rocks, only sparse outcrops of sedimentary and plutonic rocks are located in the northern ranges, in the border with Honduras ([2]). Some of the most recent volcanic layers are formed by poor consolidated ashes and tuffs highly erodible ([3]).

Throughout the year the country experiences a tropical climate with two seasons, a dry season (November to April) and wet season (May - October). The climate of El Salvador is generally warm. In the dry season there is very less rainfall but during rainy seasons heavy showers take place. Periodically El Salvador is affected by tropical storms and occasionally by hurricanes.

As in other parts of Central America, landslides in El Salvador constitute an important natural hazard due to prevailing steep terrain covered with poor consolidated volcanic materials and the frequent occurrence of extreme precipitation events and intense earthquakes. This problem is ex-

acerbated by the extreme deforestation and the consequent high level of rates of erosion. Poverty, overpopulation and uncontrolled urbanization characterize to the Salvadoran human settlements, which makes to El Salvador a country with high landslide risks. An example of this high risk was the devastating effect is Las Colinas landslides, triggered by a major earthquake (Mw 7.6) on January 13th, 2001 in Santa Tecla, a major city located close to San Salvador the Salvadoran capital([4]). A huge amount of soil mass (about 200,000 m^3) was thrown off the rim of El Balsamo range and flushed many houses and produced more than 500 deaths. Together with this event also several landslides occurred along the country, especially in the Metropolitan Area of San Salvador (MASS) ([5]). Another example is the large number of heavy-rainfall induced landslides occurred during Hurricane Mitch on October and November 1998 ([6]).

Due to the above it is necessary to implement mechanisms that allow us to quantify the hazard of a given geographic area to landslides, usually this is done with development of susceptibility maps which present in a graphical way the zones more susceptible to landslides and represent a practical tool for urban planning. It was proposed an Artificial Neural Network model (ANN) which is a family of statistical learning models inspired by biological neural networks. The model proposed is used to estimate the susceptibility to landslide. From the results obtained by the model, a map was derived using kriging which is a method of interpolation spatial from the geostatistical approach.

2. Brief State of Art

Since the pioneering work ([7]), several mathematics and statistics models have been proposed to model landslide susceptibility: deterministic models ([8], [9] and [10]), probabilistic models ([11], [12] and [13]).

It has been used popular classification models such as logistic regression ([14] and [15] and [16]), neural networks ([18],[19], [20]) and [21] and support vector machine ([22] and [23]).

According to ([24]) the magnitude of a possible slide is difficult to foresee as it depends on the magnitude of the triggering event and the environmental conditions (e.g., height of water table) at the moment of the event. Because of these complex relationships between the dependent variable and causal factors, and since that neural networks are particularly useful for detect complex non-linear relationships in large datasets, that is why this kind of model was chosen, despite the disadvantages such as greater computational burden and proneness to overfitting.




Figure 1: Location of El Salvador in Central America, map of El Salvador with location of the Metropolitan Area of San Salvador (MASS) and its topography.

3. Study area

MASS, is formed by 14 municipalities among them it is located San Salvador, the capital of the country. It has 587 km^2 and according to the most recent census ([25]) it has 1,566,629 inhabitants with a density of population of 2669 hab/km². According to geography MASS is extended over a flat erosional surface (650-760 m.a.s.l.), gently sloping to the east towards Ilopango Caldera also known as Ilopango Lake. MASS is bordered to the south by the Balsamo Range and San Jacinto Hill, to the west by San Salvador volcano and with Marionna Hills (refer to Fig. 1). Human settlement is not only restricted to the plain, but spreads up to surrounding heights and volcano flanks.

All rocks exposed in MASS have volcanic origin and consists of intercalated primary and reworked deposits from Late Tertiary to Holocene age ([26]). The younger ones are known as Tierra Blanca pyroclastic ash deposits and forms a relative continuous layer in majority of MASS with an average thickness of 4 m ([26]). These are deposits derived mainly from Ilopango caldera eruptions where they present more than 50 m of thickness ([26]). Tierra Blanca soils is highly susceptible to erosion, one of the most common dangerous effects of the erosion in this area are landslides during

heavy rainfall and earthquake ground shaking ([26]). Chávez et al. ([27]) identifies high and moderate erosion hazard in majority of MASS area. The more recent layer Tierra Blanca, known as Tierra Blanca Joven (TBJ) ([28]) present very poor geotechnical properties, especially when there is an increment of soil moisture (i.e. during rainy seasons), which results in high instability susceptibility ([29] and [30]). Several incised rivers and ravines cross MASS area, due the large erosion rates of the area, natural slopes of these watercourses are often close to vertical and can reach highs of more than 10 m. Taking into account this property of volcanic soils, usually vertical slopes are cut to urban and road construction. As Bommer & Rodriguez ([3]) identified in Central America, although such slopes may remain stable for years, they may become unstable, abruptly and totally under the action of heavy rainfall or seismic shaking. Hernández ([28]) carried out a detailed description of this process in TBJ, where these instabilities are very common. In study area fault trends are characterized by steep dip angles (65° - 90°). The main fault system is characterized by an east-west trend; this trend is responsible of the steep northern slope of the El Balsamo Range. The secondary systems are with north, northwest and northeast trends offset the main fault system and are more active in the present tectonic setting ([26]). However, all the mentioned fault families appear active. Thus the several fault systems have developed at different times, but been repeatedly activated ([31]). Finally there are also several ringlike faults that can be related to subsurface collapse due to volcanic tectonic subsidence ([26]). Last eruption of San Salvador volcano was 1917. The majority of the magma's total volume (97%) extruded from several vents in the Northwest flank of the volcano, outside of MASS. However a fraction of MASS is located under the proximal volcanic hazard zone according to Sofield ([34]). In the steep slopes San Salvador volcano, especially of Picacho, the highest peak of the volcano, located in the eastern side present a high lahar hazard that threats MASS ([33]). On the other hand one of the most destructive eruptions in the world history was 430 A.D. TBJ eruption produced by the Ilopango Caldera. According to Dull ([32]), that eruption produced total devastation in whole MASS area.

4. Landslide inventory and data sources of input variables

The Landslide inventory was developed by MARN and it was conducted in two stages. The first was carried out, based on fieldwork after the 2001 earthquakes in El Salvador. The second stage consisted of a visual inventory of places where there were landslides identified through satellite

image from IRS-C sensor (Indian) in the same year.

Once the landslide inventory was finished, a percentage of 0.5% of the total points georeferenced showed landslide occurrence. In addition to the landslide information the following data sources of input variables was provided by MARN:

1. **Geomorphology:** refers to landforms that result from lithospheric dynamics of geographic area. This variable was obtained in the project of national land use plan in El Salvador. It is unpublished geomorphologic cartography in digital format provided by MARN.
2. **Slope:** Slope gradient is an important component and a preparatory cause of landsliding ([16]). It was calculated from digital elevation model (DEM) provided by MARN with a resolution of 10 mt.
3. **Geology:** Description of the geology of the area according digitalized geological map of El Salvador (scale 1:100,000) ([2]).
4. **Rainfall intensity:** From the precipitation database compiled by the MARN's weather stations for the period 1970–2001, maximum precipitations were obtained, and then, a contour map was generated through kriging interpolation method.
5. **Peak ground acceleration:** Maximum ground acceleration expressed in Gal for a return period of 500 years. This data was obtained from the evaluation of seismic Hazard in Central America carried out under the framework of RSIS II project ([35]).
6. **Road distance:** Distance in kilometers to the nearest road. Road map was obtained from digitalized national cartography created by Centro Nacional de Registros de la República de El Salvador (CNR).
7. **Fault distance:** Distance in kilometers to the nearest fault. This information was obtained from according digitalized geological map of El Salvador (scale 1:100,000) ([2]).

5. Artificial Neural Network model for discrete choice

Logistic regression represents a neural network with a neuron in the hidden layer and whose output is dichotomous. The following adapted form of the Multilayer Perceptron (MLP) may be used for modeling binary classification problems. In the equation (1) $x_{k,i}$ are the observed values in

the i th input variables belonging to the k th case. The expressions $w_{j,k}$ and λ_j showed in the equations (1) and (3) are the parameters of the model.

$$n_{j,i} = w_{j,0} + \sum_{k=1}^{k^*} w_{j,k} x_{k,i} \quad (1)$$

$$N_{j,i} = \frac{1}{1 + \exp^{-n_{j,i}}} \quad (2)$$

$$p_i = \sum_{j=1}^{j^*} \lambda_j N_{j,i} \quad (3)$$

$$\sum_{j=1}^{j^*} \lambda_j = 1, \lambda_j \geq 0 \quad (4)$$

In the equation (3) p_i is the the predicting probability for a network with k^* input characteristics and j^* neurons. In the context of the present research, p_i represents the probability of landslide occurrence.

Before estimating the parameters of the neural network model, it is necessary standardize the input variables. In particular for classification problems is more suitable to scale inputs to $[-1, 1]$ rather than $[0, 1]$ ([36]). The following scaling was applied to each input variable:

$$z = \frac{x - \bar{x}}{\sigma_x} \quad (5)$$

In the equation (5) z is the new variable transformed, \bar{x} and σ_x are, respectively the mean and the standard deviation of the input variable to transform.

The method used for estimating the parameters of the model was an Hybrid Method ([37]): Firstly heuristic genetic algorithm using a package developed in the R statistical software ([38]) specifically developed for this purpose, was used to obtain a good estimation of the parameters of the model.

The R package can be accessed from the following web address:

<https://goo.gl/PHaaG2>

Once a good estimate was obtained, this was occupied as the initial values for the conjugate gradient method implemented in the `optim` routine of the R statistical software, to obtain a better estimation of the parameters of the model.

6. Spatial Prediction

Estimating spatial correlation

In standard statistical problems, correlation can be estimated from a scatterplot, when several data pairs x, y are available. The spatial correlation between two observations of a variable $z(s)$ at locations s_1 and s_2 cannot be estimated, as only a single pair is available. To estimate spatial correlation from observational data, it is necessary to make stationarity assumptions before we can make any progress. One commonly used form of stationarity is intrinsic stationarity, which assumes that the process that generated the samples is a random function $Z(s)$ composed of a mean and residual [39]:

$$Z(s) = \mu + \delta(s) \quad (6)$$

with a constant mean

$$E(Z(s)) = \mu \quad (7)$$

and a variogram defined as

$$\lambda(h) = \frac{1}{2} E(Z(s) - Z(s+h))^2 \quad (8)$$

Ordinary kriging in terms of the covariance function

The predictor assumption is

$$\hat{Z}(s_0) = \sum_{i=1}^n w_i Z(s_i) \quad (9)$$

It is a weighted average of the sample values, and $\sum_{i=1}^n w_i = 1$ to ensure unbiasedness. The w_i 's are the weights that will be estimated. Kriging minimizes the mean squared error of prediction [40].

$$\min \sigma_e^2 = E \left[Z(s_0) - \hat{Z}(s_0) \right]^2 \quad (10)$$

7. Results

The data were randomly divided into three sub-samples, the first was used for the estimation of the parameters of the neural network (training set), the second was used for choosing the model with more generalization

capability (validation set) and the last was used to evaluate how well the model generalize outside of the data set used for estimation (test set).

To determine the number of neurons in the hidden layer was used the method of trial and error, starting with few neurons and increasing progressively the number of neurons in the hidden layer. Table 1 summarizes how well the models fit on the training, validation and test sets.

Table 1: Summary of classification accuracy on the training, validation and test sets for neural network models

Number of neurons	Percentage score		
	Train set	Validation set	Test set
2	0.7011	0.7020	0.7124
3	0.7206	0.7072	0.7401
4	0.7425	0.7197	0.7458
5	0.7601	0.7604	0.7740
6	0.7823	0.7704	0.7604
7	0.7853	0.7704	0.7763
8	0.7942	0.7865	0.7878
9	0.8009	0.7871	0.7901
10	0.8321	0.8132	0.8009
11	0.8194	0.7792	0.7542
12	0.8230	0.8006	0.8159
13	0.8408	0.8006	0.8031
14	0.8373	0.8017	0.8127
15	0.8517	0.8210	0.8247
16	0.8246	0.8017	0.8028
17	0.8543	0.8283	0.8246
18	0.8467	0.8022	0.8418
19	0.8341	0.7991	0.8274

The model with 17 neurons in the hidden layer was chosen, after that a logistic regression model was fitted as a basis of comparison. Table 2 presents the logistic regression fits, clearly the logistic regression's performance was lower than the worst neural network model.

Table 2: Summary of classification accuracy on the training, validation and test sets for logistic regression

Model	Percentage score		
	Train set	Validation set	Test set
logistic regression	0.6710	0.6748	0.663

To generate the map of landslide hazards, the following steps were followed based on geostatistics methodology: Exploratory analysis, Variogram modelling and Spatial prediction using ordinary kriging, validation of this results and .

In order to achieve the above steps a R script was developed. This script can be downloaded in:

<https://goo.gl/3N8n99>

First of all one thousand fifty georeferenced point with their probability of occurrence which was calculated using neural network, were selected randomly for the map generation process leaving the others for validation of the kriging model. The Exploratory analysis showed the presence of spatial correlation in the data, this was done making scatter plot of pairs $Z(s_i)$ and $Z(s_j)$, grouped according to their separation distance, also the data did not show the presence of anisotropic effect.

As regards the estimation of the variogram, the spherical, gaussian and exponential models were fitted but the validation of these models on data that were not taken into account in the process of estimating the variogram showed that the best model was the exponential, since that it showed a $R^2 = 0.42$ against $R^2 = 0.40$ and $R^2 = 0.39$ of the models spherical and gaussian respectively. With the exponential model were made spatial predictions on a grid of 1500×1500 and after that, the spatial predictions were exported to Geotiff raster format. Finally Geographic Information System (ArcGIS® of ESRI) was used to design the final map (Figure 2). This map (Fig. 2) represents the landslide probability of occurrences in MASS. According to Fig. 2 higher probability of landslide occurrence, reaching probability of occurrence values around 0.8, are located in the northern slopes of Balsamo Range (western side of MASS), in the eastern slope of San Salvador volcano and Picacho, along the extension of Mariona Hills and in the southeastern side, between Ilopango lake and San Jacinto Hill. Areas with lower probability but still important, are located in southern and western side of Nejapa Hill, in north-east of the study area, in northern banks of Ilopango lake, in northeast slope of San Salvador Volcano, in the surrounding areas of San Jacinto Hill and, in general in southern side of MASS.

8. Discussion and conclusions

A landslide hazard assessment study was carried out in Metropolitan Area of San Salvador (MASS). The study started with the construction of a landslide inventory and analysis of the causal factors related to the occurrences of landslide. The problem of modelling landslide generated by

different causes is very complex and for this reason the study proved the efficacy of the neural network model with a percentage of correct classification around 80% against other models such as logistic regression with a percentage of correct classification under 70%.

In the process of estimating the weights of the model an heuristic technique was used to obtain a better solution after that a local search was used. It is better than use only backpropagation algorithm or another local search method to estimate weights since that when these are used there is a strong danger of getting stuck in a local minimum rather than a global minimum for a vector of weights.

The problem with the neural network approach is that is difficult to estimate the weights of the model due to intensive computation involved in the present study the estimation of a neural network model takes between 4 and 10 hours for this reason we could not assess the statistical significance of the input variables in the neural network processes using bootstrapping. For all of the above parallel computing must be used rather than serial computing in estimating weights of neural networks models.

The results obtained in the geostatistics methodology showed that the spatial the data satisfies the conditions for applying kriging method. Final map of landslide probability of occurrence (see Fig. 2) generated by kriging method is the result of multidisciplinary work which implied an intense geological analysis (i.e. MARN inventory of 4792 landslides) together with the use of sophisticated statistical techniques. This landslide hazard map is a user-friendly tool to evaluate landslide risk in MASS and can be useful for many purposes such as territorial planning, prevention and landslide mitigation and so on.

High probability of landslide occurrence in Picacho slope implies very high risk due the proximity of urban areas and due the large volume of mobilized materials. Major et al.([33]) identified in San Salvador volcano that lahars as voluminous as 2 million of mts^3 can affect large zones of urban areas. Southern areas of MASS also implies a high risk because large landslides, similar to Las Colinas landslide which involved a total volume of around 150,000 m^3 ([4]), can occur in northern slope of Balsamo range and San Jacinto Hill affecting very crowded urban areas.

The present hazard landslide map is consistent with former landslide hazard maps carried out in the region using approaches conceptually different. Fernández-Lavado et al.([41]) presented a landslide hazard GIS map of MASS generated through bivariate statistical method considering several conditioning factors showing similar hazard areas. The only differences are located mainly in the Nejapa Hill, San Jacinto Hill where




Figure 2: Landslide hazard map MASS where is shown the landslide probability of occurrence.

([41]) identifies a slightly higher hazard and the area between San Jacinto Hill and Ilopango Lake where the former study presents a hazard slightly lower. As landslides are an erosional phenomenon, the highest probability areas of landslide occurrence coincides with the areas with highest hazard of erosion in the erosion hazard map of MASS proposed by ([27]). There are similarities with susceptibility hazard maps at national scale generated through logistic regression method ([16]), through artificial neural network method ([17]) and through Mora-Vahrson method ([42]). However the high landslide areas are more extended in these national maps than the present study. These differences could be explained by the working scale, our map is more detailed because is at regional scale. Other possible explanation can be related to the data, present work was built using a more extended landslide catalog and more updated data (e.g. seismic hazard evaluation from Benito et al.[35]) than former works.

Rock falls were not included in the catalog; these were very common during 2001 earthquakes in El Balsamo Range ([5]). On the other hand, the MASS seismic microzonation of seismic site effects will improve the estimation of peak ground acceleration. The local geologic conditions are responsible for the modifications (amplification, frequency content and duration) experimented by seismic motions just before reaching the ground

surface, additionally, the geometry of the ground surface also may produce amplification (topographic effects) ([43]). These effects can be important in soft soils close to incised rivers and also in San Salvador volcano, Balsamo ranges and hills. Crosta et al.([44]) identified an amplification factor of 1.3-1.4 in the Las Colinas failure area (Balsamo Ranges). All these variables can improve the predictive ability of the model and surely the likelihood of landslides.

9. Acknowledgment

The authors are indebted with MARN who provided the data and Geographic Information System software such as ILWIS and ArcGIS® of ESRI.

References

- [1] Dewey J.W. White R.A., Hernández D.A. Seismicity and tectonics of El Salvador *Chapter 27 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J. 367-378. (2004)*
- [2] Weber H.S., Wieseemann G., Lorenz W., Schmidt-Thomé M. Mapa geológico de Republica de El Salvador escala 1 : 100 000 (6 sheets). *Bundesanstalt für Geowissenschaften und Rohstoffe, Hannover, Germany. (1978).*
- [3] Bommer J.J., Rodriguez C.E. Earthquake-induced landslides in Central América *Engineering Geology. (63):1889-220, (2002).*
- [4] Evans S.G. Bent A.L. Las Colinas landslide, Santa Tecla: A highly destructive flowslide triggered by January 13, 2001, El Salvador earthquake. *Chapter 3 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J. 25-37. (2004)*
- [5] Jibson R.W., Crone A.J., Harp E.L., Baum R.L., Major J.J., Pullinguer C.R., Escobar D., Martinez M., Smith M.E. Landslides triggered by the 13 January and 13 February 2001, earthquakes. *Chapter 6 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J. 69-88. (2004)*

- [6] Crone A.J., Baum R.L., Lidke D.J., Sather D.N.D., Bradley L.-A., Tarr A.C. Landslides Induced by Hurricane Mitch in El Salvador- An Inventory and Description of Selected Features *USGS Open File Repord 01-444. (2002)*
- [7] Alberto Carrara. Multivariate models for landslide hazard evaluation. *Journal of the International Association for Mathematical Geology*, 15 (3):403–426, 1983. ISSN 0020-5958. URL <http://dx.doi.org/10.1007/BF01031290>.
- [8] G. Capparelli and P. P. Versace. Landslide susceptibility from mathematical model in sarno area. *Hydrology and Earth System Sciences Discussions*, 10(10):12643–12662, 2013. URL <http://www.hydrol-earth-syst-sci-discuss.net/10/12643/2013/>.
- [9] Biswajeet Pradhan, Hyun-Joo Oh, and Manfred Buchroithner. Weights-of-evidence model applied to landslide susceptibility mapping in a tropical hilly area. *Geomatics, Natural Hazards and Risk*, 1(3):199–223, 2010. URL <http://www.tandfonline.com/doi/abs/10.1080/19475705.2010.498151>.
- [10] Bettina Neuhäuser, Bodo Damm, and Birgit Terhorst. Gis-based assessment of landslide susceptibility on the base of the weights-of-evidence model. *Landslides*, 9(4):511–528, 2012. ISSN 1612-510X. URL <http://dx.doi.org/10.1007/s10346-011-0305-5>.
- [11] Campbell R.H. Brookshire D.S. Bernknopf, R.L. and C.D Shapiro. A probabilistic approach to landslide hazard mapping in cincinnati, ohio, with application for economic evaluation. *Bulletin of the Association of Engineering Geologists*, XXV(1):39–56, 1988.
- [12] Chang-JoF. Chung and AndreaG. Fabbri. Validation of spatial prediction models for landslide hazard mapping. *Natural Hazards*, 30 (3):451–472, 2003. ISSN 0921-030X. URL <http://dx.doi.org/10.1023/B%3ANHAZ.0000007172.62651.2b>.
- [13] S. Lee, J. Choi, and K. Min. Probabilistic landslide hazard mapping using gis and remote sensing data at boun, korea. *International Journal of Remote Sensing*, 25(11):2037–2052, 2004. URL <http://www.tandfonline.com/doi/abs/10.1080/01431160310001618734>.
- [14] Akgun, Aykut A comparison of landslide susceptibility maps produced by logistic regression, multi criteria decision, and likelihood ratio methods: a case study at Izmir, Turkey *Landslides*, (11):9, 1, 93–106, 2012 Springer.

- [15] Gaskill, Jacob and Zuber, Brian and Nordman, Erik Analyzing Landslide Susceptibility in St. Vincent and the Grenadines Using Co-Kriging and Logistic Regression *The 2015 IMAGIN Award, Michigan United States*. URL http://www.imagin.org/awards/sppc/2015/papers/jacob_gaskill_paper.pdf.
- [16] García-Rodríguez M.J., Malpica J.A., Benito B., Díaz M. Susceptibility assessment of earthquake-triggered landslides in El Salvador using logistic regression. *Geomorphology* 95, 172-191, 2008.
- [17] García-Rodríguez M.J., Malpica J.A. Assessment of earthquake-triggered landslide susceptibility in El Salvador based on an Artificial Neural Network model. *Nat. Hazards Earth Syst. Sci.* 10, 1307–1315, 2010. doi:10.5194/nhess-10-1307-2010.
- [18] C. Melchiorre, E.A. Castellanos Abella, C.J. van Westen, and M. Matteucci. Evaluation of prediction capability, robustness, and sensitivity in non-linear landslide susceptibility models, guantánamo, cuba. *Computers & Geosciences*, 37(4):410 – 425, 2011. ISSN 0098-3004. URL <http://www.sciencedirect.com/science/article/pii/S0098300410003286>.
- [19] Xu Zeng-wang. Gis and ann model for landslide susceptibility mapping. *Journal of Geographical Sciences*, 11(3):374–381, 2001. ISSN 1009-637X. URL <http://dx.doi.org/10.1007/BF02892323>.
- [20] Artificial neural networks applied to landslide susceptibility assessment. *Geomorphology*, 66(1–4):327 – 343, 2005. ISSN 0169-555X URL <http://www.sciencedirect.com/science/article/pii/S0169555X04002272>.
- [21] E. Yesilnacar and T. Topal. Landslide susceptibility mapping: A comparison of logistic regression and neural networks methods in a medium scale study, Hendek region (Turkey). *Engineering Geology*, 79(11):251-266, 2005. ISSN 0013-7952, URL <http://www.sciencedirect.com/science/article/pii/S0013795205000384>.
- [22] Ballabio, Cristiano and Sterlacchini, Simone Support vector machines for landslide susceptibility mapping: the Staffora River Basin case study, Italy *Mathematical geosciences*, (11):1, 47–70, 2012. Springer.
- [23] Tien Bui, Dieu and Pradhan, Biswajeet and Lofman, Owe and Revhaug, Inge Landslide susceptibility assessment in vietnam using support vector machines, decision tree, and Naive Bayes Mod-

- els *Mathematical Problems in Engineering*, (11):2012, 2012. Hindawi Publishing Corporation.
- [24] Van Westen, CJ and Van Asch, Th WJ and Soeters, R Landslide hazard and risk zonation why is it still so difficult? *Bulletin of Engineering geology and the Environment*. (65):2, 167–184, 2006 Springer.
 - [25] Ministerio de Economía- Dirección General de Estadística y Censos VI Censo de Población y V de Vivienda 2007-2008. Available from <http://goo.gl/yvjzPy>
 - [26] Schmidt-Thomé M. The geology in San Salvador area (El Salvador, Central America), a basis for city development and planning. *Geol. Jb.* 13, 207-228, 1975.
 - [27] Chávez J.A., Sebesta J., Kopecky L., López R. Application of Geomorphologic Knowledge for Erosion Hazard Mapping *Natural Hazards*. 71,1323–1354, 2014. DOI 10.1007/s11069-013-0948-8.
 - [28] Hernández W.E. Características Geomecánicas y Vulcanológicas de las Tefras Tierra Blanca Joven, Caldera de Ilopango, El Salvador. MSc Thesis. Universidad Politécnica de Madrid – Universidad Politécnica de El Salvador. 2004
 - [29] Chávez J.A., Landaverde J.M., Ayala O.E., Mendoza L.E. (2014b). Application of Constitutive Models in the Volcanic Tephra “Tierra Blanca Joven”. *Ingeniería*. 24 (2).53-78. ISSN: 2215-2652.
 - [30] Rolo R., Bommer J.J., Houghton B.F., Vallance J.W., Berdousis P., Mavrommati C., Morphy W. Geologic and engineering characterization of Tierra Blanca pyroclastic ash deposits. *Chapter 5 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J.* 55-67, 2004.
 - [31] Rymer M.J. The San Salvador Earthquake of October 10, 1986. *Geologic Aspects Earthquake Spectra*. 3(3), 435-463, 1987.
 - [32] Dull R.A. Lessons from the mud, lessons from the Maya: Paleoeological records of the Tierra Blanca Joven eruption. *Chapter 7 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J.* 237-244., 2004.

- [33] Major J.J. Chilling S.P., Pullinguer C.R., Escobar C.D. Debris-flow hazards at San Salvador, San Vicente, and San Miguel volcanoes, El Salvador. *Chapter 7 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J.* 89-108, 2004.
- [34] Sofield D. Eruptive history and volcanic hazards of Volcan San Salvador. *Chapter 7 in Geological Society of America Special paper 375 in Natural Hazards in El Salvador of Rose W.I., Bommer J.J., López D.L., Carr M.J. and Major J.J.* 147-158, 2004.
- [35] Benito M.B., Lindholm C., Camacho E., Climent Á., Marroquín G., Molina E., Rojas W., Escobar J.J., Talavera E., Alvarado G.E. and Torres Y. A New Evaluation of Seismic Hazard for the Central America Region. *Bulletin of the Seismological Society of America* 102 (2), 504-523. 2012. doi: 10.788/0120110015
- [36] Warren S. Sarle URL <ftp://ftp.sas.com/pub/neural/FAQ.html#questions>. Accessed 31 May, 2015.
- [37] Paul D. McNelis Estimation of a Network with Evolutionary Computation *Academic Press*, (11):59-84, 2005. ISBN 9780124859678, URL <http://www.sciencedirect.com/science/article/pii/B9780124859678500038>.
- [38] R Core Team R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*. Vienna, Austria, 2014.
- [39] Bivand, Roger S and Pebesma, Edzer J and Gómez-Rubio, Virgilio and Pebesma, Edzer Jan Applied spatial data analysis with R *volume 747248717* Springer, 2014.
- [40] Nicolas Christou. Ordinary kriging in terms of the covariance function *University of California, Los Angeles* [cited 2015 June 04]. Available from <http://goo.gl/0J1RkO>
- [41] Fernandez-Lavado C., Sanchez A., Amenos M., Barrio J. Caracterización de la susceptibilidad y de la amenaza por movimientos de ladera del Área Metropolitana de San Salvador (AMSS). Scale 1: 75000 (1 Sheet). *Project IPGARAMSS framework. Geólogos del Mundo. San Salvador, El Salvador.* 2008.
- [42] SNET-MARN, Servicio Nacional de Estudios Territoriales-Ministerio de Ambiente y Recursos Naturales Memoria

Técnica Para El Mapa De Susceptibilidad de Deslizamientos de Tierra En El Salvador. *Ministerio de Ambiente y Recursos Naturales* [cited 2015 June 04]. Map available from <http://www.snet.gob.sv/ver/geologia/susceptibilidad+actual/>

- [43] Aki, K. Local site effects on weak and strong ground motion. In F. Lund (ed.) *New Horizons in Strong Motion: Seismic Studies and Engineering Practice. Tectonophysics* 218, 93-111, 1993.
- [44] Crosta G.B., Imposimato S., Rodderman, Chiesa S.m, Moia F. Small fast-moving flow-like landslides in volcanic deposits: The 2001 Las Colinas Landslide (El Salvador). *Engineering Geology*. 79, 185-214, 2005.