
Incremental Stability of Discrete-Time First-Order Projection Elements

Riccardo Bertollo

Sebastiaan van den Eijnden

W.P.M.H. (Maurice) Heemels

{r.bertollo, s.j.a.m.v.d.eijnden, w.p.m.h.heemels}@tue.nl

Incremental Stability of Discrete-Time First-Order Projection Elements

[clearly not a talk on learning]

Riccardo Bertollo


Sebastiaan van den Eijnden

W.P.M.H. (Maurice) Heemels

{r.bertollo, s.j.a.m.v.d.eijnden, w.p.m.h.heemels}@tue.nl

Motivation: Improving performance of linear controllers

Integrator-based (linear) controllers are needed to reject constant disturbances and overcome friction, but they have some **intrinsic limitations**




Motivation: Improving performance of linear controllers

Integrator-based (linear) controllers are needed to reject constant disturbances and overcome friction, but they have some **intrinsic limitations**

[Clegg, 1958] proposed the addition of a resetting mechanism

$$\mathcal{H} := \begin{cases} \dot{u} = e, & ue \geq 0 \\ u^+ = 0, & ue < 0 \end{cases}$$



Motivation: Improving performance of linear controllers


Integrator-based (linear) controllers are needed to reject constant disturbances and overcome friction, but they have some **intrinsic limitations**

[Clegg, 1958] proposed the addition of a resetting mechanism

$$\mathcal{H} := \begin{cases} \dot{u} = e, & ue \geq 0 \\ u^+ = 0, & ue < 0 \end{cases}$$

HOWEVER:

resets might be **too aggressive** to implement in practice



Motivation: Improving performance of linear controllers

Integrator-based (linear) controllers are needed to reject constant disturbances and overcome friction, but they have some **intrinsic limitations**



[Clegg, 1958] p

Can we keep the good properties of the controller, but with continuous solutions?

$$\mathcal{H} := \begin{cases} u^+ = 0, & ue < 0 \end{cases}$$

HOWEVER:

resets might be **too aggressive** to implement in practice



A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$


So, what if we **project the dynamics** instead of resetting?

A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?




A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?




A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?



A new approach: Projection-based control

IDEA:


The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?



Hybrid Integrator-Gain Systems (HIGS) [Deenen et al., 2021]

$$\dot{u} = \begin{cases} \omega_h e, & (\omega_h e, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (\omega_h e, \dot{e}) \notin T_S(u) \end{cases}$$



A new approach: Projection-based control

IDEA:


The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?



Hybrid Integrator-Gain Systems (HIGS) [Deenen et al., 2021]

$$\dot{u} = \begin{cases} \omega_h e, & (\omega_h e, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (\omega_h e, \dot{e}) \notin T_S(u) \end{cases}$$



A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?




Hybrid Integrator-Gain Systems (HIGS) [Deenen et al., 2021]

$$\dot{u} = \begin{cases} \omega_h e, & (\omega_h e, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (\omega_h e, \dot{e}) \notin T_S(u) \end{cases}$$

We can extend this to any underlying dynamics

$$\dot{u} = \begin{cases} v, & (v, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (v, \dot{e}) \notin T_S(u) \end{cases}$$



A new approach: Projection-based control

IDEA:

The “good thing” about the Clegg integrator is that $eu \geq 0$

So, what if we **project the dynamics** instead of resetting?




Hybrid Integrator-Gain Systems (HIGS) [Deenen et al., 2021]

$$\dot{u} = \begin{cases} \omega_h e, & (\omega_h e, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (\omega_h e, \dot{e}) \notin T_S(u) \end{cases}$$


First-Order Projection Element (FOPE)

$$\dot{u} = \begin{cases} au + be, & (au + be, \dot{e}) \in T_S(u) \\ k_h \dot{e}, & (au + be, \dot{e}) \notin T_S(u) \end{cases}$$



Projection-based control: the ePDS framework

Since the error derivative cannot be modified, we can
project only in **selected directions**



Projection-based control: the ePDS framework

Since the error derivative cannot be modified, we can
project only in **selected directions**




Extended Projected Dynamical Systems (ePDS)

$$\dot{x} = \Pi_{\mathcal{S}, E} f(x) = f(x) + \mu$$

where

$$\begin{aligned} \mu &:= \operatorname{argmin} ||w|| \\ \text{s. t. } & f(x) + w \in T_{\mathcal{S}}(x) \\ & w \in \operatorname{Im}(E) \end{aligned}$$



Projection-based control: the ePDS framework

Since the error derivative cannot be modified, we can
project only in **selected directions**


Extended Proj

$\dot{x} =$


What happens in the discrete-time case?

where

$$\begin{aligned} \mu &:= \operatorname{argmin} ||w|| \\ \text{s.t. } & f(x) + w \in T_{\mathcal{S}}(x) \\ & w \in \operatorname{Im}(E) \end{aligned}$$




(DT) First-Order Projection Elements



(DT) First-Order Projection Elements

Possible issues:


- **not necessarily** from the **boundary** of the set



(DT) First-Order Projection Elements

Possible issues:

- **not necessarily** from the **boundary** of the set
 - e can **change sign**



(DT) First-Order Projection Elements

Possible issues:


- **not necessarily** from the **boundary** of the set
- e can **change sign**



First-Order Projection Element: projection on an **input-dependent set**

$$u^+ = \Pi_{K(e^+)}(v) = \Pi_{K(e^+)}(au + be^+)$$

(where $|a| \leq 1, b > 0$)



(DT) First-Order Projection Elements

Possible issues:


- **not necessarily** from the **boundary** of the set
- e can **change sign**

What are we investigating specifically?


First-Order Projection Element: projection on an **input-dependent set**

$$u^+ = \Pi_{K(e^+)}(v) = \Pi_{K(e^+)}(au + be^+)$$

(where $|a| \leq 1, b > 0$)




Research question: Discrete-time Incremental Stability




DEFINITION: Incremental (asymptotic/exponential) stability

If two copies of the system are subject to the **same input**, their **trajectories converge to each other** (asymptotically/exponentially)

Literature overview



Literature overview




**Research
focus**

Corresponding article(s)

**CT incremental
stability**


Van den Eijnden et al., L-CSS, 2023

Literature overview




Research focus	Corresponding article(s)
CT incremental stability	Van den Eijnden et al., L-CSS, 2023
DT formulation and stability	Sharif et al., CDC, 2022 Sharif et al., Automatica, 2024

Literature overview




Research focus	Corresponding article(s)
CT incremental stability	Van den Eijnden et al., L-CSS, 2023
DT formulation and stability	Sharif et al., CDC, 2022 Sharif et al., Automatica, 2024
DT/SD convergence properties	Pavlov et al., TAC, 2012 Tren et al., TAC, 2019

Literature overview




Research focus	Corresponding article(s)
CT incremental stability	Van den Eijnden et al., L-CSS, 2023
DT formulation and stability	Sharif et al., CDC, 2022 Sharif et al., Automatica, 2024
DT/SD convergence properties	Pavlov et al., TAC, 2012 Tren et al., TAC, 2019
DT/SD incremental stability and convergence	Bertollo et al., ADHS, 2024


Open-loop Incremental Stability




Open-loop Incremental Stability




Open-loop Incremental Stability



Open-loop Incremental Stability



Open-loop Incremental Stability



LEMMA 1:

The projection on the same set $K(e)$ is **non-expansive**

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

$$|\delta u^+| \leq |\delta v|$$

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

$$\begin{aligned} |\delta u^+| &\leq |\delta v| \\ &\leq |au_1 + be^+ - au_2 - be^+| \end{aligned}$$

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

$$\begin{aligned} |\delta u^+| &\leq |\delta v| \\ &\leq |au_1 + be^+ - au_2 - be^+| \\ &= |a||\delta u| \end{aligned}$$

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

$$\begin{aligned} |\delta u^+| &\leq |\delta v| \\ &\leq |au_1 + be^+ - au_2 - be^+| \\ &= |a||\delta u| \end{aligned}$$

THEOREM 1:

If the underlying dynamics is GES (Globally Exponentially Stable), the FOPE is **incrementally exponentially stable**

Open-loop Incremental Stability: GES underlying dynamics

The case $|a| < 1$ is **straightforward**:

$$|\delta u^+| \leq |\delta v|$$

$$\leq |au_1 + be^+ - au_2 - be^+|$$

$$= |a||\delta u|$$

If $a=1$, this is not enough!

THEOREM 1:


If the underlying dynamics is GES (Globally Exponentially Stable), the FOPE is **incrementally exponentially stable**

Open-loop Incremental Stability: Integrator case

- **Integrator** dynamics is **not incrementally stable**


Open-loop Incremental Stability: Integrator case

- **Integrator** dynamics is **not incrementally stable**
- **When the projection is active** for (at least) one of the two trajectories, the **inequality** $|\delta u^+| < |\delta v|$ is **strict**




Open-loop Incremental Stability: Integrator case

- **Integrator** dynamics is **not incrementally stable**
- **When the projection is active** for (at least) one of the two trajectories, the **inequality** $|\delta u^+| < |\delta v|$ is **strict**
- **Without additional constraints**, it is possible that the projection is **never active**



Open-loop Incremental Stability: Integrator case

- **Integrator** dynamics is **not incrementally stable**
- **When the projection is active** for (at least) one of the two trajectories, the **inequality** $|\delta u^+| < |\delta v|$ is **strict**
- **Without** additional **constraints**, it is possible that the projection is **never active**



THEOREM 2:


If $a=1$, then the FOPE is **incrementally asymptotically stable if and only if** the parameter $b > 0$ satisfies

$$b \notin (2k_1, 2k_2)$$

Open-loop Incremental Stability: Integrator case

- **Integrator** dynamics is **not incrementally stable**
- **When the projection is active** for (at least) one of the two trajectories, the **inequality** $|\delta u^+| < |\delta v|$ is **strict**
- **Without** additional projection is **not**

What if the inputs are not the same?




THEOREM 2:

If $a=1$, then the FOPE is **incrementally asymptotically stable** if and only if the parameter $b>0$ satisfies

$$b \notin (2k_1, 2k_2)$$

Research question: DT Incremental ISS



DEFINITION: Incremental input-to-state stability (ISS)

If two copies of the system are subject to **different inputs**, the **bound on the distance between their trajectories** converges to some positive value **related to the distance between the inputs**

Open-loop Incremental ISS


- What happens if two copies of a FOPE are subject to two **different error signals**?

Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2


Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2




Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2




Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2




Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2




Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2




Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2
- However, the projection (written as a 2D map) is **Lipschitz continuous**, with constant $\bar{k} = \max\{|k_1|, |k_2|\}$



Open-loop Incremental ISS

- What happens if two copies of a FOPE are subject to two **different error signals**?
- In general, this might **increase the distance** between u_1 and u_2
- However, the projection (written as a 2D map) is **Lipschitz continuous**, with constant $\bar{k} = \max\{|k_1|, |k_2|\}$




LEMMA 2:

If the underlying dynamics is GES (i.e., $|a| < 1$), then

$$|\delta u^+| \leq |a| |\delta u| + (b + \bar{k}) |\delta e^+|$$

And the FOPE is **incrementally exponentially input-to-state stable**

Research question: Closed-loop Incremental ISS



Closed-loop incremental ISS: Incremental small gain theorem

We can use the following result

THEOREM⁽¹⁾:


When two DT systems are interconnected in feedback as shown, if the following hold

$$|y_1(j)| \leq \beta_1(j, |x_1(0)|) + \gamma_{y,1} \|y_2\|_\infty + \gamma_{w,1} \|w_1\|_\infty$$

$$|y_2(j)| \leq \beta_2(j, |x_2(0)|) + \gamma_{y,2} \|y_1\|_\infty + \gamma_{w,2} \|w_2\|_\infty$$

$$\gamma_{y,1} \cdot \gamma_{y,2} < 1$$


then the **closed-loop dynamics is ISS**.



⁽¹⁾ Z.P. Jiang, A.R. Teel, L. Praly (1994). Small-gain theorem theorem for ISS systems and applications. *Mathematics of Control, Signals and Systems*.

Consequences of incremental ISS: Incremental small gain theorem

We can easily write an **incremental version** and apply it to the **sampled-data interconnection** of the discrete-time FOPE and a continuous-time linear plant



THEOREM 3:


If the parameters of the FOPE are selected so that

$$\frac{b + \bar{k}}{1 - |a|} < \frac{1}{\|P_{dt}\|_{\ell_1}}$$

then the **sampled-data interconnection** of the FOPE with a linear plant is **incrementally ISS**.

Consequences of incremental ISS: Incremental small gain theorem

We can easily write an **incremental version** and apply it to the **sampled-data interconnection** of the discrete-time FOPE and a continuous-time linear plant



THEOREM 3:

If the parameters of the FOPE are selected so that

$$\frac{b + \bar{k}}{1 - |a|} < \frac{1}{\|P_{dt}\|_{\ell_1}}$$

then the **sampled-data interconnection** of the FOPE with a linear plant is **incrementally ISS**.

Simulations


- Plant = **precision positioning system** (2DoF)
- Closed loop is perturbed by a sinusoidal disturbance with frequency 6π rad/s
- Simulations starting from different ICs

Simulations

- Plant = **precision positioning system** (2DoF)
- Closed loop is perturbed by a sinusoidal disturbance with frequency 6π rad/s
- Simulations starting from different ICs


RESULTS:

- The states of the closed loop converge to the same trajectory
- The trajectories have the **same frequency as the input signal**



Simulations

Close-up on the previous plots, to highlight the sampled-data nature of the feedback loop



Conclusion

- We analyzed the dynamics of the **discrete-time FOPE**, a control element obtained by projecting a first order dynamics onto a sector
- We showed necessary and sufficient conditions for **incremental stability in open-loop** and sufficient conditions for **incremental ISS in closed loop**

Conclusion

- We analyzed the dynamics of the **discrete-time FOPE**, a control element obtained by projecting a first order dynamics onto a sector
- We showed necessary and sufficient conditions for **incremental stability in open-loop** and sufficient conditions for **incremental ISS in closed loop**

FUTURE WORK:

Conclusion

- We analyzed the dynamics of the **discrete-time FOPE**, a control element obtained by projecting a first order dynamics onto a sector
- We showed necessary and sufficient conditions for **incremental stability in open-loop** and sufficient conditions for **incremental ISS in closed loop**

FUTURE WORK:

- Formally prove **convergence properties** (all the ingredients are there, but we need to be careful about some details)

Conclusion

- We analyzed the dynamics of the **discrete-time FOPE**, a control element obtained by projecting a first order dynamics onto a sector
- We showed necessary and sufficient conditions for **incremental stability in open-loop** and sufficient conditions for **incremental ISS in closed loop**

FUTURE WORK:


- Formally prove **convergence properties** (all the ingredients are there, but we need to be careful about some details)
- Prove **incremental ISS** of the closed loop through **less conservative conditions**




Conclusion

- We analyzed the dynamics of the **discrete-time FOPE**, a control element obtained by projecting a first order dynamics onto a sector
- We showed necessary and sufficient conditions for **incremental stability in open-loop** and sufficient conditions for **incremental ISS in closed loop**

FUTURE WORK:

- Formally prove **convergence properties** (all the ingredients are there, but we need to be careful about some details)
- Prove **incremental ISS** of the closed loop through **less conservative conditions**
- The condition for incremental stability with $a=1$ are quite conservative: is there a **better formulation?**





Riccardo Bertollo
r.bertollo@tue.nl

