

Lecture 13 - Attention and impulsivity

The Social Brain: Critical Perspectives on Science, Society and Neurodiversity


Richard Ramsey



Today

Part 1

- Attention and impulsivity



Part 2

- Read articles and discuss



Overview

- Background & ADHD
- Performance on executive function tasks
 - Behavioural evidence
 - Neurobiological evidence



Background & ADHD



Living with ADHD

<http://www.youtube.com/watch?v=mjmiWkB1TKw>




ADHD

- Attention Deficit/Hyperactivity Disorder (ADHD) is defined by a combination of symptoms of

- Inattention
- Hyperactivity/impulsivity

- 3 types:



Diagnosis

- Six (or more) of the following symptoms of inattention or hyperactivity-impulsivity have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level. Some examples as follows:
- Inattention
 - Often does not give close attention to details or makes careless mistakes in schoolwork.
 - Often has trouble keeping attention on tasks or play activities.
 - Often does not seem to listen when spoken to directly.
- Hyperactivity-impulsivity
 - Often fidgets with hands or feet or squirms in seat.
 - Often gets up from seat when remaining in seat is expected.
 - Often runs about or climbs when and where it is not appropriate.



Features

- Peak onset between the age of 3 and 4
- Affects 3%-5% children, prevalence higher in areas of socio-economic disadvantage
- Male/Female ratio
 - 4:1 for the hyperactivity/impulsivity type
 - 2:1 for the inattention type
- Co-morbidity: 50% of the children also show another psychiatric disorder:
 - Learning difficulties, Oppositional Defiant Disorder and Conduct Disorder, Anxiety and mood disorders, Substance abuse, Tic disorders
- Medical/Physical characteristics:
 - More accident prone, Sleep problems
 - Persists in adolescence for around 50-80% of individuals and in adulthood around 30-50% of individuals



Genetics

- Disorder clusters in families
 - Increased risk for 1st and 2nd order degree relatives (Faraone et al. 1995)
 - Siblings are 3 to 5 times more at risk (Biederman et al., 1992, Faraone et al., 1993)
 - Concordance is higher in monozygotic twins (50%-80%) than dizygotic twins (33%) (Bradley & Golden, 2001)

Several candidate genes (Durstun & Konrad, 2007), some linked to dopaminergic system and others to the serotonergic system. Likely to have a complex genetic origin



Cold vs hot executive tasks

- “Cold” executive functions tasks are associated with working memory, inhibition and planning.
- “Hot” executive function tasks are associated with reward and reinforcement mechanisms



Stroop task

Baselines

GREEN **RED** **BLUE**


Cognitive control condition

BLUE **GREEN** **RED**

word is read automatically
→ cognitive control is
needed to inhibit reading
response



Stroop task processes



Delay aversion

- Greater preference for smaller-immediate over larger-delayed rewards (choice impulsivity)
- 1CHF now or 100 CHF in a year?
 - At which point do you shift your choice?
- Marshmallow test for kids. One now or two later?



Performance on executive function tasks

- Behavioural evidence
- fMRI evidence



Behavioural evidence

- Review of 8 meta-analyses:
 - Large effect: >0.50
 - Medium effect: $0.50-0.30$
 - Small effect: <0.30



(Nigg 2005)

Nigg 2005 results

Table 2. Selected Meta-analytic Findings in Neuropsychology of ADHD Versus Non-ADHD Children

Measure	Effect Size (<i>d</i>)
Spatial Working Memory (Spatial Span)	.75 ^a to .85 ^b to 1.14 ^b
Response Suppression (Stop Task SSRT/SSRT Slope)	.61 ^a to .64 ^c to .94 ^d
Signal Detection (CPT d-prime) Arousal	.72 ^e
Stroop Naming Speed	.69 ^f
Full Scale IQ	.61 ^g
Set Shifting (Trails B Time)	.55 ^a to .59 ^g to 0.75 ^d
Planning (Tower of London/Hanoi)	.51 ^a to .69 ^a
Mazes	.58 ^a
Verbal Working Memory	.51 ^a to .41 ^b
Decision Speed on Go-Task	.49 ^c
WCST Perseverations	.35 ^g /.36 ^a to .53 ^h
Fluency	.27 ^d
Stroop Interference	.25 ^f
Covert Visual Spatial Orienting	.20 ⁱ



Nigg 2005 results


Table 2. Selected Meta-analytic Findings in Neuropsychology of ADHD Versus Non-ADHD Children

Measure	Effect Size (<i>d</i>)
Spatial Working Memory (Spatial Span)	.75 ^a to .85 ^b to 1.14 ^b
Response Suppression (Stop Task SSRT/SSRT Slope)	.61 ^a to .64 ^c to .94 ^d
Signal Detection (CPT d-prime) Arousal	.72 ^e
Stroop Naming Speed	.69 ^f
Full Scale IQ	.61 ^g
Set Shifting (Trails B Time)	.55 ^a to .59 ^g to 0.75 ^d
Planning (Tower of London/Hanoi)	.51 ^a to .69 ^a
Mazes	.58 ^a
Verbal Working Memory	.51 ^a to .41 ^b
Decision Speed on Go-Task	.49 ^c
WCST Perseverations	.35 ^g /.36 ^a to .53 ^h
Fluency	.27 ^d
Stroop Interference	.25 ^f
Covert Visual Spatial Orienting	.20 ⁱ




Neurobiological evidence

- Functional MRI studies reveal atypical activity in
 - Prefrontal cortex (esp. right inferior frontal)
 - Basal ganglia (esp. putamen)
 - Cerebellum
 - Parietal areas



Response suppression task

Response suppression task



^a The interval between horizontal and vertical arrows in the stop trials becomes smaller/larger in steps of 50 msec depending on each subject's performance to ensure 50% successful and 50% unsuccessful inhibition for each subject.




ETH Zürich | 376-1309-00 | The Social Brain

(Rubia et al. 2005, Rubia 2018)

Response suppression results

FIGURE 2. Regions of Increased Activation During Successful Relative to Unsuccessful Inhibition in Healthy Subjects (N=21) Versus Medication-Naïve Adolescents With ADHD (N=16)^a



^a Talairach coordinates are indicated for slice distance (in mm) from the intercommissural line for the horizontal (A, B, C) and sagittal slices (D, E). The focus reaches from the right orbitoprefrontal (see A) to inferior prefrontal cortex (B, C), deep into the insula (B, D, E) and precentral gyrus (C, E), and bordering the temporal cortex (A, B, D, E). Cluster-wise probability of false positive activation: $p < 0.01$.



ETH Zürich | 376-1309-00 | The Social Brain

(Rubia et al. 2005; Rubia 2018)

“Cold” task summary


Individuals with ADHD show:

- Atypical performance in EF tasks
- Atypical neural responses compared to controls in “cold” EF tasks
- What about “hot” tasks?



Reward task

fMRI: Male adults with ADHD versus controls




ETH Zürich | 376-1309-00 | The Social Brain

(Ströbele et al 2008)

Reward results

Reduced striatum activation when anticipating the gain




ETH Zürich | 376-1309-00 | The Social Brain

(Ströbele et al. 2008)

Reward results

Increased orbitofrontal activation when presented with gain outcome
(over-sensitivity to reward)



(Ströbele et al. 2008)

Summary

Individuals with ADHD show:

- Atypical performance in EF tasks
- Atypical neural responses compared to controls in “cold” and “hot” EF tasks

However...



Individual differences

% of ADHD children classified as impaired
 → usually less than half!

Table 1. Illustrative Widely Used Neuropsychologic Measures Comparing ADHD (Combined Type) to Controls: Group Differences and Percent Impaired in 3 Samples

Measure	Sample	Effect Size (d)		p	% ADHD Beyond Control 90th Percentile
		d	η^2		
SSRT	MI (all)	.88	.133	<.001	51
	CO	.79	.101	<.001	45
RT Variability	MI	.75	.123	<.001	48
	CO	.77	.125	<.001	44
Stroop CW	MI	.50	.045	<.05	25
	CO	.84	.132	<.001	44
	MGH	.62	.09	<.001	25
CPT	MI	.91	.11	<.001	37
	CO	.54	.053	<.001	35
	MGH	.17	.01	.11	16
Trailmaking	MI	.35	.033	<.05	27
	CO	.35	.031	<.01	24

MI=Michigan sample
 CO=Colorado sample
 MGH=Massachusetts sample

(Nigg 2005)



Individual differences


When ADHD children tested on both inhibition and delay aversion and looking at individual performances rather than group performance:

- 23% deficit on both
- 23% inhibition deficit only
- 15% delay aversion deficit only
- 39% no apparent deficit on any of the two


So what does this mean?




Single cognitive mechanism?



Inhibition



Reward



Future research

- Is there good evidence for a single cognitive deficit hypothesis?
- How well does a deficit in inhibition or motivation/reward processing explain the clinical symptoms associated with ADHD?
- Is it plausible to assume a single core deficit? What is the alternative?



ETH Zürich | 376-1309-00 | The Social Brain

(Castellanos et al. 2006, Rubia 2018, Sonuga-Barke 2005)

Take a break



Part 2 - Read and discuss



Discussion material

- break into small groups (~ 5 per group)
- discuss aspects of the lecture
- discuss aspects of the journal article: [Rubia, 2018](#)
- There is no need to read all of the paper. Pick a section of the paper that interests you and focus on that for the discussion.



References

- Casey, B. j., & Durston, Ph. d., Sarah. (2006). From Behavior to Cognition to the Brain and Back: What Have We Learned From Functional Imaging Studies of Attention Deficit Hyperactivity Disorder? *American Journal of Psychiatry*, 163(6), 957–960. <https://doi.org/10.1176/ajp.2006.163.6.957>
- Castellanos, F. X., Sonuga-Barke, E. J. S., Milham, M. P., & Tannock, R. (2006). Characterizing cognition in ADHD: Beyond executive dysfunction. *Trends in Cognitive Sciences*, 10(3), 117–123. <https://doi.org/10.1016/j.tics.2006.01.011>
- Nigg, J. T. (2005). Neuropsychologic Theory and Findings in Attention-Deficit/Hyperactivity Disorder: The State of the Field and Salient Challenges for the Coming Decade. *Biological Psychiatry*, 57(11), 1424–1435. <https://doi.org/10.1016/j.biopsych.2004.11.011>
- Rubia, K. (2018). Cognitive Neuroscience of Attention Deficit Hyperactivity Disorder (ADHD) and Its Clinical Translation. *Frontiers in Human Neuroscience*, 12. <https://doi.org/10.3389/fnhum.2018.00100>
- Rubia, K., Smith, A. B., Brammer, M. J., Toone, B., & Taylor, E. (2005). Abnormal Brain Activation During Inhibition and Error Detection in Medication-Naive Adolescents With ADHD. *American Journal of Psychiatry*. <https://doi.org/10.1176/appi.ajp.162.6.1067>
- Sonuga-Barke, E. J. S. (2005). Causal Models of Attention-Deficit/Hyperactivity Disorder: From Common Simple Deficits to Multiple Developmental Pathways. *Biological Psychiatry*, 57(11), 1231–1238. <https://doi.org/10.1016/j.biopsych.2004.09.008>



Ströhle, A., Stoy, M., Wrase, J., Schwarzer, S., Schlagenhauf, F., Huss, M., Hein, J., Nedderhüt, A., Neumann, B., Gregor, A., Juckel, G., Knutson, B., Lehmkuhl, U., Bauer, M., & Heinz, A. (2008). Reward anticipation and outcomes in adult males with attention-deficit/hyperactivity disorder. *NeuroImage*, 39(3), 966–972. <https://doi.org/10.1016/j.neuroimage.2007.09.044>



Acknowledgements

- Unless otherwise specified, icons and images were used under license from [The Noun Project](#)
- Slides were created with [Quarto](#) and [RevealJS](#)

