

June 10, 2014

Brad Koffel, Attorney
The Koffel Law Firm
1801 Watermark Dr., Ste. 350
Columbus, OH 43215

Re: Clinical Alcohol Assessment for Robert Major

Dear Mr. Koffel:

I performed a Clinical Alcohol Assessment for Mr. Major in reference to his recently adjudicated OVI case. This assessment included the administration of: a MAST, an AUDIT, an anxiety/depression screening, and a 50 minute face-to-face clinical interview. The following are my findings and clinical recommendations:

MAST: score = 3; additional screening important for clarification of condition

AUDIT: score = 9; score suggests likelihood of mild alcohol use disorder (minimum)

DSM/Other Substance Related Concerns: meets 2 of 11 DSM 5 symptoms - mild to moderate tolerance, use of alcohol that conflicts with personal health interests

Anxiety/Depression Screening: mild to moderate symptoms including some insomnia and concentration problems; receiving medical treatment / no crisis

Substance Related Legal History Reported: denied any previous history of alcohol or substance related arrests/legal charges

Clinical Interview/Summary: Mr. Major's written screenings suggested the probability of a mild alcohol use disorder. He reported a wide variation in his normal use of alcohol; some weeks consuming no alcohol and other weeks (50% or less) drinking 4-6 drinks on several occasions per week. Mr. Major was transparent in reporting both the amounts and the effects of his consumption. He appears to have mild to moderate alcohol tolerance and some internal conflict about alcohol impacting his overall health. Recommendations (below) are provided to help moderate his alcohol use, decrease alcohol tolerance and hopefully prevent any problematic use of alcohol in the future.

DSM 5 DIAGNOSIS in reference to substances: 305.00, mild alcohol use disorder


Treatment Recommendations:

- 1) Education, Ohio Health's Drug/Alcohol Awareness Series (contact 614-566-3700);
- 2) Preventative Counseling, recommend 3-4 sessions with a substance abuse professional (once per month); if moderation of consumption is not successful or additional problem symptoms become evident during this time, additional clinical recommendations and options should be provided by the professional;
- 3) Read Controlling Your Drinking by Miller and Munoz;

- 4) If another negative consequence is incurred as a result of any alcohol use, it is recommended Mr. Major seek further evaluation and follow all professional recommendations.

Mr. Major has signed a release to communicate the results of this assessment to you. Feel free to contact me should you have any additional questions pertaining to this matter.

Sincerely,



Brian Davis, LISW-S, SAP
cc: Robert Major

Prohibition Against Re-Disclosure: This information has been disclosed to you from records protected by Federal Confidentiality rules. The Federal rules prohibit you from making any further disclosure of this information unless further disclosure is expressly permitted by the written consent of the person to whom it pertains or as otherwise permitted by 42 C.F.R Part 2. A general authorization for the release of medical or other information is not sufficient for this purpose. The Federal rules restrict any use of information to criminally investigate or prosecute any alcohol or drug abuse client. Drug abuse patient records are also protected under the Health Insurance Portability and Accountability Act of 1996 ("HIPAA"), 45 C.R.R parts 160 & 165. (These conditions apply to every page disclosed and a copy of this authorization will accompany every disclosure.)