

October 19, 2016

Wapello County Clerk of Court Wapello County Courthouse 101 West 4th Street Ottumwa, Iowa 52501

Re: Clinical Alcohol Assessment for John D. Grooms (DOB – 11/13/1993)

Dear Wapello County Clerk of Court:

I performed a Clinical Alcohol Assessment for Mr. Grooms in reference to his DUI charge in Wapello County. This assessment included a 50-minute clinical interview by telephone, and the administration of:

- A Michigan Alcohol Screening Test (MAST)
- An Alcohol Use Disorder Identification Test (AUDIT)
- A general anxiety/depression screening

The following items provide a summary of the screening results, my diagnostic opinion, and professional recommendations:

MAST: score = 2 / NEGATIVE SCREEN; no direct or indirect indication of a present alcohol use disorder (screen relative to lifetime use of alcohol)

AUDIT: score = 5 / NEGATIVE SCREEN; scores of 8 or more indicate potential hazardous use, whereas scores of 15 or more in men indicate a moderate to severe disorder may be present (screen predominantly relative to current use of alcohol)

Alcohol-Use Risk Factors:

□ negative peer influence/at-risk work environment – NA
□ current stress (relational, work, financial, etc.) – NA
☐ physical health concerns – NA
□ learning disability/mental health concerns – ADHD
□ exposure to trauma – NA
oximes family history of substance abuse or alcoholism – 1 blood-relative of significance
□ contraindicated prescription use – NA
☑ tobacco use – daily
☑ regular use of liquor – mixed drinks (monthly basis)
□ voluntary treatment history – NA

⊠ severity of incident – impaired memory, auto accident (.132 BAC)
☐ legal history – denies previous history
Summary - 5 of 12 concerns, mitigate concern with clinical recommendations
Clinical Interview/Summary: Mr. Grooms' written screenings provided no indication of an alcohol use disorder at any level (mild, moderate or severe). He reported his normal use of alcohol before this charge as 3-4 "Jack and Cokes" on 1-3 occasions per month. He also acknowledged use of up to 5-6 drinks on 3-4 special occasions per year.
Mr. Grooms and I discussed his DUI and general use of alcohol at length. I did not find him meeting DSM-5 alcohol use disorder criteria for diagnosis and his typical drinking habits seem to be within NIAAA recommendations for men's low-risk drinking. I did however, note the risks and concerns associated with his specific situation. Accordingly, I am including some preventative recommendations in the remainder of this report.
DSM-5 Alcohol Use Disorder Symptoms: ☐ Use beyond intention ☐ Persistent desire or unsuccessful efforts to reduce/quit use ☐ Substantial time spent obtaining, using, or recovering from use ☐ Strong craving/desire ☐ Failure to fulfill a major role (work, home, school) ☐ Use despite recurrent social/interpersonal problems ☐ Important social, occupational, recreational activities relinquished ☐ Physically hazardous use ☐ Continued use despite knowledge of contraindicated physical or psychological condition ☐ Tolerance ☐ Withdrawal symptoms
Summary - 0 of 11 symptoms identified (Mild 2-3 Moderate 4-5 Severe 6+)
DSM-5 DIAGNOSIS in Reference to Substances : Z03.89 (suspected substance use condition not found)
Treatment Recommendations:

1) 48 Hour Weekend OWI Program – complete program and provide documentation of completion to the court;

DIRECTIONS COUNSELING GROUP

2) <u>If another negative consequence</u> is incurred as a result of any alcohol use it is recommended Mr. Grooms seek an increased level of treatment or extended outpatient substance abuse counseling as appropriate.

If you or the court possess additional information about Mr. Grooms' use of alcohol that might impact the outcome of this assessment, I would be happy to review it and consider appropriate modifications or amendments to the present report.

Thank you for the opportunity to assist the court, Mr. Grooms, and yourself in this important matter.

Sincerely,

Trevor C. Davis, CDCA Certificate # 150427

Reviewed by Brian T. Davis, LISW-S, SAP LIC# I-7948

cc: John D. Grooms

Prohibition Against Re-Disclosure: This information has been disclosed to you from records protected by Federal Confidentiality rules. The Federal rules prohibit you from making any further disclosure of this information unless further disclosure is expressly permitted by the written consent of the person to whom it pertains or as otherwise permitted by 42 C.F.R Part 2. A general authorization for the release of medical or other information is not sufficient for this purpose. The Federal rules restrict any use of information to criminally investigate or prosecute any alcohol or drug abuse client. Drug abuse patient records are also protected under the Health Insurance Portability and Accountability Act of 1996 ("HIPAA"), 45 C.R.R parts 160 & 165. (These conditions apply to every page disclosed and a copy of this authorization will accompany every disclosure.)

Diagnostic Conclusions and Limitations: Directions Counseling Group provides Clinical Alcohol Assessments based on objective and standardized screenings and interviewing methods as well as self-reported information. Clinical conclusions are made to the best of the professional's ability with data on hand at the time of the assessment. Expanded assessments which involve additional background research, collateral information, and third party verification can be provided as an additional service upon request.

