

June 28, 2016

Attn: Susan Cox, CAC II
South Carolina Interstate ADSAP Office
Dorchester Alcohol and Drug Commission
500 North Main Street Suite 4
Summerville, SC 29483

Re: Clinical Alcohol Assessment for Joseph Jones (DOB – 11/20/1975)

Dear Ms. Cox:

I performed a Clinical Alcohol Assessment for Mr. Jones in reference to his April 2004 OVI charge in Horry County. This assessment included a 60-minute clinical interview and the administration of:

- A Substance Abuse Subtle Screening Inventory (SASSI-3)
- A Michigan Alcohol Screening Test (MAST)
- An Alcohol Use Disorder Identification Test (AUDIT)
- A general anxiety/depression screening

The following items provide a summary of the screening results, my diagnostic opinion and professional recommendations:

SASSI-3: no scores indicative of an alcohol use disorder at any level (mild, moderate, or severe); screening had a lifetime frame of reference;

MAST: score = 1 / *NEGATIVE SCREEN*; no direct or indirect indication of problem alcohol use; lifetime frame of reference

AUDIT: score = 3 / *NEGATIVE SCREEN*; scores of 8 or more indicate *hazardous use*, scores of 15 or more in men indicates *moderate to severe* disorder present

Alcohol Use Risk Factors:

- ☐ family history – NA
- ☐ prior DUI charges - NA
- ☐ early onset of alcohol use - NA
- ☐ prior alcohol/substance use treatment - NA
- ☐ peer influences - NA
- ☐ work environment - NA
- ☐ contraindicated prescription use – No Rx use
- ☐ anxiety or depression symptoms - None

- ☐ general health problems – No known diseases of allergies
- ☐ current family stress - low

Alcohol Use Risk profile = 0 of 10; low risk

Anxiety/Depression Screening: no clinically significant concerns reported

Substance Related Legal History Reported: denies any previous history of alcohol or substance related arrests/legal charges

Clinical Interview/Summary: Mr. Jones' written screenings provided no indication of an alcohol use disorder at any level (mild, moderate or severe). He reported his normal use of alcohol over the past 24 months as 2-3 drinks on 1-2 occasions per week. He stated that he will drink more than 4 drinks (e.g. up to 6) on approximately 2-4 special celebratory occasions per year.

Mr. Jones and I discussed his April 2004 alcohol-related OVI charge and his general use of alcohol at the time. Mr. Jones' clinical interview revealed no positive DSM-5 symptoms. His clinical interview did not reveal indications of recurrent use beyond intention, continued use despite interpersonal problems, continued use despite negative impact to rule functioning, nor a tolerance or withdraw condition.

In my clinical judgment, I was unable to find Mr. Jones' April 2004 incident, his general history of alcohol use nor his recent alcohol use patterns as indicative of someone experiencing DSM-5 alcohol use disorder symptoms. In summary, I do not find evidence that Mr. Jones meets criteria for an alcohol use disorder.

DSM-5 DIAGNOSIS in Reference to Substances: z03.89 (suspected substance use condition not found)

Treatment Recommendations:

- 1) There are no clinical recommendations at this time. However, it is my understanding that Mr. Jones must participate in a minimum of 16 hours of alcohol education. Therefore, I am recommending that Mr. Jones complete a 3-Driver Intervention Program (recommend Wellness Driver, 614-791-8300 or Second Chance Counseling, 614-635-9011).
- 2) If another negative consequence is incurred as a result of any alcohol use it is recommended Mr. Jones seeks an updated evaluation and follows any and all treatment recommendations.

Thank you for the opportunity to assist the state of South Carolina ADSAP office, Mr. Jones, and yourself in this important matter.

Sincerely,

Ryan P. Smith

Ryan P. Smith MSW, LISW-S, LICDC, SAP
SW Lic. #I.1000155-S, CD Lic. #101182
cc: Joseph Jones

Prohibition against Re-Disclosure: This information has been disclosed to you from records protected by Federal Confidentiality rules. The Federal rules prohibit you from making any further disclosure of this information unless further disclosure is expressly permitted by the written consent of the person to whom it pertains or as otherwise permitted by 42 C.F.R Part 2. A general authorization for the release of medical or other information is not sufficient for this purpose. The Federal rules restrict any use of information to criminally investigate or prosecute any alcohol or drug abuse client. Drug abuse patient records are also protected under the Health Insurance Portability and Accountability Act of 1996 ("HIPAA"), 45 C.R.R parts 160 & 165. (These conditions apply to every page disclosed and a copy of this authorization will accompany every disclosure.)

Diagnostic Conclusions / Limitations: Directions Counseling Group provides Clinical Alcohol Assessments based on objective and standardized screenings and interviewing methods as well as self-reported information. Clinical conclusions are made to the best of the professional's ability with data on hand at the time of the assessment. Expanded assessments which involve additional background research, collateral information and third party verification can be provided as an additional service upon request.