

June 9, 2016

Walter A. Piel Jr., Attorney
Piel Law Office
500 West Silver Spring Drive, Suite K200
Milwaukee, WI 53217-5052

Re: Clinical Alcohol Assessment for Richard L. Woida (DOB – 08/26/1985)

Dear Mr. Piel:

I performed a Clinical Alcohol Assessment for Mr. Woida in reference to his OWI charge in Waukesha County. This assessment included a 50-minute clinical interview and the administration of:

- A Michigan Alcohol Screening Test (MAST)
- An Alcohol Use Disorder Identification Test (AUDIT)
- A general anxiety/depression screening

The following items provide a summary of the screening results, my diagnostic opinion and professional recommendations:

MAST: score = 2 / *NEGATIVE SCREEN*; no direct or indirect indication of problem alcohol use, mitigate any concerns with clinical recommendations

AUDIT: score = 1 / *NEGATIVE SCREEN*; scores of 8 or more indicate *hazardous use*, scores of 15 or more in men indicate *moderate to severe* disorder present

Alcohol Use Risk Factors:

Concerns/Risks

- ☒ previous legal history – lifetime total of 2 DUI charges, 2008 = .09 BAC, 2016 = .078 BAC
- ☐ severity of incident – low breath alcohol test
- ☐ anxiety/depression/mental health – NA
- ☐ other substance use – NA
- ☐ family history – none reported
- ☐ prior treatment history – NA
- ☐ peer influences – NA
- ☐ work environment – NA
- ☐ contraindicated prescription use – NA
- ☐ general health problems – NA
- ☐ current family stress – NA

Summary - 1 of 11 concerns, mitigate concern with clinical recommendations

Clinical Interview/Summary:

ALCOHOL – Mr. Woida's written screenings provided no indication of an alcohol use disorder at any level (mild, moderate or severe). He reported his normal use of alcohol as 2-4 drinks on 0-1 occasions per week. He reported no difference in the amount consumed if he was attending a special event or celebration where he consumed alcohol.

Mr. Woida and I discussed his DUI incidents and general use of alcohol at length. While I did not find him meeting DSM-5 alcohol use disorder criteria I did note that it would be appropriate to take some step toward reducing his risk of a future charge of a similar nature, since he presently has 2 lifetime charges. Preventative education is an appropriate recommendation and further details are provided below for Mr. Woida.

DSM-5 DIAGNOSIS in Reference to Substances: Z03.89 (suspected substance use condition not found)


Treatment Recommendations:

- 1) Alcohol & Drug Education – 4 hrs. (minimum) of alcohol/drug education, recommend use of an online course since Mr. Woida travels regularly but an in-person local class would also be acceptable if one can be found (online examples include logancourtclasses.com or alcoholclass.com);
- 2) If another negative consequence is incurred as a result of any alcohol use it is recommended Mr. Woida seek further evaluation and follow all treatment recommendations.

If you or the court possess additional information about Mr. Woida's use of alcohol that might impact the outcome of this assessment, I would be happy to review it and consider appropriate modifications or amendments to the present report.

Thank you for the opportunity to assist the court, Mr. Woida, and yourself in this important matter.

Sincerely,



Brian T. Davis, LISW-S, SAP
LIC# I-7948
cc: Richard L. Woida

Prohibition Against Re-Disclosure: This information has been disclosed to you from records protected by Federal Confidentiality rules. The Federal rules prohibit you from making any further disclosure of this information unless further disclosure is expressly permitted by the written consent of the person to whom it pertains or as otherwise permitted by 42 C.F.R Part 2. A general authorization for the release of medical or other information is not sufficient for this purpose. The Federal rules restrict any use of information to

criminally investigate or prosecute any alcohol or drug abuse client. Drug abuse patient records are also protected under the Health Insurance Portability and Accountability Act of 1996 ("HIPAA"), 45 C.R.R parts 160 & 165. (These conditions apply to every page disclosed and a copy of this authorization will accompany every disclosure.)

Diagnostic Conclusions / Limitations: Directions Counseling Group provides Clinical Alcohol Assessments based on objective and standardized screenings and interviewing methods as well as self-reported information. Clinical conclusions are made to the best of the professional's ability with data on hand at the time of the assessment. Expanded assessments which involve additional background research, collateral information and third party verification can be provided as an additional service upon request.