

CO-AMOXICLAV VIAL

Only to be administered if suitably trained and competent to do so

PRESENTATION: Co-amoxiclav 1000mg/200mg (amoxicillin 1000mg, clavulanic acid 200 mg) Injection

ROUTE: IV and ORAL

INDICATIONS:

Trauma Prophylaxis...

- Penetrating injuries of the head/ neck / thorax / abdomen / limbs
- Open fractures
- Blast injuries with tissue loss

ACTIONS:

- Broad-spectrum penicillin-based antibiotic with action against beta-lactamase producing strains

CONTRAINDICATIONS:

- Penicillin allergy

CAUTIONS:

- Rash occurs commonly but they are not usually related to true penicillin allergy.
- Patient needs urgent MEDEVAC to Role2

SIDE EFFECTS:

- Dizziness, Dyspnoea, Headache, Diarrhoea

PREPARATION /ADMINISTRATION:

- Draw up 20 ml Water for Injection using a green hypodermic needle and inject into 1.2 g vial of co-amoxiclav.
- Swirl (DO NOT shake) until powder dissolves (should form colourless liquid).
- Administer over 3-4 minutes
- 1.2 g every 8 hours

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	VOLUME
Adult	1.2 g	1.2 g	8 hours	20 mL WFI