

PENTHROX

PRESENTATION: 3ml Vapour Vial Pentrox (methoxyflurane)

ROUTE: Inhalation

INDICATIONS:

- Emergency relief of pain in conscious adult patients
- Pain score of 2 or 3
- Fractures, Dislocations, Lacerations/wounds, Burns/Scalds, Chest/Abdo injuries

ACTIONS:

- Methoxyflurane, works by affecting the brain and spinal cord and making them less responsive to pain

CONTRAINDICATIONS:

- Allergy to anaesthetic drugs (or family history)- give fentanyl instead
- Liver damage or kidney impairment
- Loss of consciousness including due to head injury drugs and alcohol
- Cardiovascular instability and respiratory depression

CAUTIONS:

- Administration on consecutive days is not recommended

SIDE EFFECTS:

- Nausea, Drowsiness, Coughing, Vomiting, Dizziness, Fever, Headache

PREPARATION /ADMINISTRATION:

- Self administered by conscious adult
- Onset of pain relief within 6-10 inhalations, peak effect 15mins after treatment
- Provides analgesic relief for up to 25-30mins with continuous use or 1-hour intermittent use
- Max of 2 doses may provide analgesia for up to 2 hours

Assembly Instructions: PTO

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	DAILY MAX	MAX DOSE IN WEEK
Adult	3ml	3ml	Can be back-to-back	6ml in 24hrs	15ml