

HYPOGLYCAEMIA

Does the patient have:

- | | | |
|--|--|--------------------------------------|
| ☐ Hunger? | ☐ Dizziness? | ☐ Confusion? |
| ☐ Shaking? | ☐ Headache? | ☐ Drowsiness? |
| ☐ Sweating? | ☐ Irritability? | ☐ Seizure? |
| ☐ Blurred vision? | ☐ Weakness? | |

BG <4mmol/l

Possible hypoglycaemia
Check blood glucose (BG)

BG >4mmol/l

Confirmed hypoglycaemia
Alert MO

Not Hypoglycaemic
Consider alternative diagnosis

Patient co-operative and able to swallow?

Patient uncooperative, unable to swallow or unconscious?

Give 15-20g of fast acting carbohydrate
e.g. sweets, cake, sugary drink
OR
1-2 tubes 40% glucose oral gel

Begin immediate evacuation to

ABCDE Assessment

Gain IV access

Administer 200ml of 10% glucose IV
OR
Administer glucagon 1mg IM

Recheck BG after 10-15min

Recheck BG after 10-15min

BG <4mmol/l

BG >4mmol/l

Give up to 3 doses of fast acting carbohydrate

Give 20g long-acting carbohydrate
e.g. bread, biscuits
If patient received glucagon, give larger carbohydrate portion

BG >4mmol/l
and patient recovered

BG <4mmol/l

BG <4mmol/l
after 3 doses

Give up to 3 doses of glucose

Follow uncooperative patient algorithm

Refer to MO

Evacuate