

CEFTRIAXONE

Only to be administered if suitably trained and competent to do so

PRESENTATION: Ceftriaxone Injection 2g vial, Rocephin 2g

ROUTE: IV

INDICATIONS:

- Suspected bacterial meningitis
- Sepsis
- Intra-abdominal infections
- Treatment of human bites & animal bites
- Complicated skin and soft tissue infections
- Infections of bones and joints

ACTIONS:

Cephalosporin antibiotic – broad-spectrum used for the treatment of septicemia, meningitis etc.

CONTRAINDICATIONS:

- Hypersensitivity – up to 7% cross-sensitivity in those who are penicillin allergic.
- Avoid in patients who have severe penicillin allergy (Does not contain and Pnicillin)

CAUTIONS: Patients with suspected meningitis- urgent MEDEVAC to Role 2

SIDE EFFECTS: Bloating, Change in taste, Dizziness, Headache, Heartburn, Indigestion, Gas

PREPARATION /ADMINISTRATION:

- 2g IV
- Draw up 20 ml Water for Injection using a green hypodermic needle and inject into a 2 g vial of ceftriaxone.
- Swirl (DO NOT shake) until powder dissolves.
- Give over 5 minutes.
- Repeat at 12 hours if patient not evacuated

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	VOLUME
Adult	2 g	2 g	12 hours	20 mL WFI