

KETAMINE

PRESENTATION: Ketamine 50mg/ml (Be aware comes in different strengths)

ROUTE: IN or IM

INDICATIONS: Moderate to severe pain (2 or 3 on the DMS Pain Score) following traumatic injuries and burns.

Adults A or V on the AVPU score, pain score of 2 or more not controlled:

- At least 15 minutes after completion of maximum allowable dose of Oral Transmucosal Fentanyl Citrate (OFTC) or other opiates
- By other forms of analgesia (Penthrox, simple analgesia, splinting)
- Where other analgesia is not available

ACTIONS: Anaesthetic with analgesia properties

CONTRAINDICATIONS:

- Known ketamine allergy
- Unresponsive to voice (P on the AVPU score) Uncontrolled hypertension (SBP >200mmHg and/or DBP >110mmHg)
- Severe cardiac disease

CAUTIONS:

- Agitated or intoxicated patients
- Stroke/ head injury
- Eye injuries
- Seizures

SIDE EFFECTS:

- Apnoea
- Laryngospasm
- Cardiovascular problems
- Vomiting
- Emergence phenomena

PREPERATION/ADMINISTRATION:

- 50mg/ML by IM or IN injection
- This must only ever be administered using:
- 1ml of 50mg/ml drawn up into a 1 ml syringe or a Defence supplied fixed dose device
- Dose can be repeated ONCE after 15 minutes if the Medic Indications for Ketamine are all still met

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	VOLUME	MAX DOSE
Adult	50mg	50mg	15 minutes	1ml	100mg