

NALOXONE

Only to be administered if suitably trained and competent to do so

PRESENTATION: Naloxone Hydrochloride 400 micrograms/ml Injection
Narcan 400 micrograms 1 ml Ampoule

ROUTE: IV or IM (Glute/ Thigh)

INDICATIONS:

- Opioid overdose
- Unconsciousness with respiratory depression where opioid overdose is a possibility

ACTIONS:

- Temporarily reverses the effects of opioid overdose Will work within minutes

CONTRAINDICATIONS:

- None

CAUTIONS:

- Effects of naloxone are shorter-acting than opiates 30-60mins – monitor patient closely as may need additional doses
- IM absorption may be unpredictable
- Patients who are physically dependent on opiates may experience violent withdrawal symptoms

SIDE EFFECTS: Dizziness, Headache, Hypotension, Nausea, Vomiting

PREPARATION /ADMINISTRATION:

- Draw 1 vial into a syringe using a green hypodermic needle and then give via IV
- If no response at 2 minutes give 800 mcg for up to 2 doses

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	VOLUME	MAXIMUM DOSE
Adult	400 mcg (1 Vial)	800 mcg (2 Vials)	2 min (If no effect with first dose or effect wears off)	1 ml – 2 ml	No max but discuss with RMO

BLOCKS THE EFFECTS OF...

Diamorphine	Hydrocodone
Morphine	Codeine
Oxycodone	Hydromorphone
Methadone	Buprenorphine
Fentanyl	

Discuss with RMO:

- ASAP after administration