

PENTHROX

PRESENTATION: 3ml Vapour Vial Penthrox (methoxyflurane)

ROUTE: Inhalation

INDICATIONS:

- Emergency relief of pain in conscious adult patients
- Pain score of 2 or 3
- Fractures, Dislocations, Lacerations/wounds, Burns/Scalds, Chest/Abdo injuries

ACTIONS:

- Methoxyflurane, works by affecting the brain and spinal cord and making them less responsive to pain

CONTRAINDICATIONS:

- Allergy to anaesthetic drugs (or family history) - give fentanyl instead
- Liver damage or kidney impairment
- Loss of consciousness including due to head injury drugs and alcohol
- Cardiovascular instability and respiratory depression

CAUTIONS:

- Administration on consecutive days is not recommended

SIDE EFFECTS:

- Nausea, Drowsiness, Coughing, Vomiting, Dizziness, Fever, Headache

PREPARATION /ADMINISTRATION:

- Self administered by conscious adult
- Onset of pain relief within 6-10 inhalations, peak effect 15mins after treatment
- Provides analgesic relief for up to 25-30mins with continuous use or 1-hour intermittent use
- Max of 2 doses may provide analgesia for up to 2 hours

Assembly Instructions: PTO

AGE	INITIAL DOSE	REPEAT DOSE	DOSE INTERVAL	DAILY MAX	MAX DOSE IN WEEK
Adult	3ml	3ml	Can be back-to-back	6ml in 24hrs	15ml

PENTHROX

Ensure the Activated Carbon (AC) Chamber is inserted into the dilutor hole on the top of the PENTHROX Inhaler.



Remove the cap of the bottle by hand. Alternatively, use the base of the PENTHROX Inhaler to loosen the cap with a ½ turn. Separate the Inhaler from the bottle and remove the cap by hand.



Tilt the PENTHROX Inhaler to a 45° angle and pour the total contents of one PENTHROX bottle into the base of the Inhaler whilst rotating.



Place wrist loop over patient's wrist. Patient inhales and exhales PENTHROX through the mouthpiece to obtain analgesia. First few breaths should be gentle and then breathe normally through Inhaler.



Patient exhales into the PENTHROX Inhaler. The exhaled vapour passes through the AC Chamber to adsorb any exhaled methoxyflurane. Always use the AC chamber to adsorb exhaled methoxyflurane and reduce occupational exposure.



If stronger analgesia is required, patient can cover dilutor hole on the AC chamber with finger during use.



If further pain relief is required, after the first bottle has been used, use a second bottle if available. Alternatively, use a second bottle from a new combination pack. Use in the same way as the first bottle in step 2 and 3. No need to remove the AC Chamber. Put used bottle into the plastic bag provided.



Patient should be instructed to inhale intermittently to achieve adequate analgesia. Continuous inhalation will reduce duration of use. Minimum dose to achieve analgesia should be administered.



Replace cap onto PENTHROX bottle. Place used PENTHROX Inhaler and used bottle in sealed plastic bag and dispose of responsibly.

