

Introduction

The Zeeman effect occurs when atmospheric molecules interact with Earth's magnetic field, causing their emission lines to split into multiple components. [zeeman1897]. One of the molecules that is affected by this is molecular oxygen, measured with the Kiruna Microwave RAdiometer(KIMRA) at the Swedish Institute of Space Physics(IRF) at the 235 GHz oxygen line since 2023.

Until KIMRA's measurements, no high-latitude instrument had ever detected this effect in Earth's atmosphere, marking the first successful ground-based observation of the phenomenon from the Arctic region.

For decades, microwave measurements of molecular oxygen have been used to retrieve altitude-dependent temperature profiles in Earth's atmosphere. In this study, we investigate whether oxygen observations from KIMRA also can be used for temperature retrievals, as previously proposed in past studies [pardo].

The novelty of this work lies in characterizing the Zeeman effect in oxygen emissions at 235 GHz, something that has never been done from high latitudes measurements. Furthermore, exploring temperature retrievals from this oxygen line may unlock new research opportunities using KIMRA data, offering valuable insights into atmospheric dynamics and temperature variations in the context of a changing climate.

Method and data

Radiation from molecules affected by the Zeeman effect is polarized along the geomagnetic field lines, circularly [zeeman1897]. While the emission is intrinsically circularly polarized along magnetic field lines, it appears linearly polarized in the perpendicular plane: horizontally for the σ_{\pm} lines and vertically (parallel to the field) for π lines.

Stokes formalism can be used to mathematically describe the polarized emission, represented by the Stokes vector, $\bar{I}_{\nu} = [I, Q, U, V]^T$. Here, I , denotes the total intensity, Q and U corresponds to components of linear polarization and V is corresponds to circular polarization.

Characterization of sub lines

A central goal of this study is the characterization of sub-lines observed with KIMRA. For the 235 GHz oxygen line, the emission is split into four σ_{\pm} components and two π components due to the Zeeman effect [pardo]. To capture this, four KIMRA measurements were carried out along the cardinal directions on January 4, 2024.

Considering the polarization geometry and the fact that KIMRA can measure only linear polarization, the σ_{\pm} components are expected when pointing North or South, while the π components should appear when pointing East or West.

The characterization is performed by comparing simulations with the measurements. The linear polarization component Q in the simulations is varied to minimize the RMS error, allowing determination of the value of Q that best reproduces the measurements for each direction.

Temperature profiling investigation

A second goal in this study is to investigate whether the oxygen line measured with KIMRA would be suitable for temperature retrievals. This is evaluated through retrievals of simulated spectra of the oxygen line KIMRA observe. For comparison, the same procedure be applied to the well-established oxygen line centered at 53 GHz, which has been used for temperature retrievals for more than a decade by colleagues at the University of Bern [bern].

A key factor in temperature retrieval is the sensitivity of the spectral line to changes in temperature. This sensitivity is quantified during the retrieval process through the Jacobian matrix, which describes how variations in temperature affect the observed spectra. By comparing the Jacobian matrices derived from the 53 GHz and 235 GHz lines, it is possible to assess whether the 235 GHz line is suitable for reliable temperature retrievals.

Results

RMS minimization between measured and simulated spectra provided consistent values for the Q component, especially in the northward ($Q = 0.87$) and southward ($Q = 0.89$) directions, which align with expectations based on the line of sight and local magnetic field. Eastward and westward measurements, however, gave $Q = -0.63$ and $Q = -1$. While the sign change of Q is expected, the magnitude difference indicates additional influences that warrant further investigation.

Temperature profiling analysis, based on Jacobian matrices from the 53 GHz and 235 GHz retrievals, further highlighted key findings. The 53 GHz line showed strong sensitivity to temperature variations across most of the atmosphere, confirming its reliability for retrievals. In contrast, the 235 GHz line displayed very low sensitivity at all pressure levels, making temperature retrievals unfeasible.

Conclusion

Minimizing the RMS to estimate the linear polarization component produced reliable results, particularly for northward and southward measurements, confirming the method's robustness. The east-west discrepancies likely stem from two sources: geomagnetic disturbances caused by the world's largest iron ore deposits near Kiruna, and cross-polarization leakage from KIMRA's optics.

The most significant finding of the temperature profiling analysis is that retrievals from the 235 GHz line are unfeasible. While the cause lies outside this study's scope, the most probable explanation is that KIMRA measures a rare isotopologue of molecular oxygen, comprising less than 0.4% of atmospheric oxygen.

References