MRN	
Surname	
Given name(s)	
Date of Birth	/ / Sex Male / Female / Other

POST ANAESTHETIC CARE UNIT RECORD - FOR NON-PREGNANT PERSONS

General Instructions

- You must record all observations, including Pain, Functional Activity Scale and Sedations Scores (page 4) at a frequency appropriate to the patient's clinical condition.
- You must calculate a Total ADDS Score for each set of observations and record it in the Total ADDS Score box, even if the score is zero (Respiratory Rate + O_2 Saturation + O_2 Flow Rate + Heart Rate + Temperature + Behaviour and Consciousness).
- Refer to the table on page 1 for documenting and escalating Changing behaviour.
- When graphing observations, place a dot (•) in the appropriate box and join to the preceding dot (e.g.).
- You must write any observation outside the range of the graph as a number.

Guide for Recognising Changing Behaviour and New Confusion (page 2)

- Reported or observed change
- Distress
- Loss of touch with reality
- Loss of function
- Elevated risk to self, others, or property

Guide for Responding to Changing Behaviour and New Confusion

- Changing Behaviour or New Confusion can be a sign of deteriorating psychology or physiology
- Escalate as per local protocol (e.g. to speciality Medical or Psychiatry)
- If required, activate a Code Black

Additional Observations (assess at least every 15 minutes)

Time assessed (24hr)												
Bleeding (nil, mod, DC [dressing changed])												
Pain score (0-10)												

Post-Anaesthetic Discharge Scoring System (PADSS) (assess at every 15 minutely intervals)

Time assessed (24hr)												
Vital signs												
Activity level												
Nausea/vomiting												
Pain												
Surgical bleeding												
TOTAL												

PADDS Scoring Matrix (total score of 10 required for discharge)

Vital signs		Activity level		Nausea and vomiting		Pain		Surgical bleeding	
BP and HR within 20% baseline	2	Ambulates at pre-op level, steady gait, no dizziness	2	Minimal – treated without further medication	2	Pain tolerated	2	Minimal – no dressing change	2
BP and HR within 20–40% baseline	1	Requires assistance	1	Moderate – successful treated with medication	1	Pain not tolerated	1	Moderate – up to 1 dressing change	1
BP and HR within >40% baseline	0	Unable to ambulate	0	Severe – remain after repeated treatment	0	IV opiate administered	0	Severe – >1 dressing change	0

Discharge Record/Checklist

PADSS score (10 required for discharge)	☐	Aftercare information given	☐
Fluids tolerated	☐	Discharge letter given to client	☐
IV cannula removed	☐	Discharge letter mailed to GP (if applicable)	☐
Analgesia discussed	☐	Copy of discharge letter in record	☐
Discharge script given (if applicable)	☐	Discharge time (24hr): ..... :	
PACU Nurse (print name):		Signature:	

POST ANAESTHETIC CARE UNIT RECORD - FOR NON PREGNANT PERSONS MR440
POST ANAESTHETIC CARE UNIT RECORD - FOR NON PREGNANT PERSONS MR440

MRN	
Surname	
Given name(s)	
Date of Birth	/ / Sex Male / Female / Other

POST ANAESTHETIC CARE UNIT RECORD - FOR NON-PREGNANT PERSONS

Interventions Relating to observations

If an intervention is administered, record here and note letter in *Intervention* row on page 2 in appropriate time column.

A
B
C
D
E

Clinical Review Request

Review 1			
Date:	Time (24hr):	Sedationist/Anaesthetist:	Proceduralist:
Reason:			
Review 2			
Date:	Time (24hr):	Sedationist/Anaesthetist:	Proceduralist:
Reason:			

Additional PACU Notes

Blood Glucose Level (assess only for diabetic patients)

MR440

Adult
(for NON-pregnant persons)

				Date																				
				Time																				
Respiratory Rate (breaths / min) Measure for a full minute	E		≥36																					E
	4		31–35	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	2		25–30																					2
	1		21–24																					1
	0		17–20																					0
			13–16																					
	1		9–12																					1
	E		≤8																					E
O_2 Saturation (%)			≥98																					
	0		95–97																					0
			92–94																					
	1		90–91																					1
	2		85–89																					2
	4		≤84	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Oxygen* (L / min *or* % delivered) *If on HF / NIV use % delivered	E	15	≥60%																					E
	4	>11–14	>50–59%	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	2	>5–11	>40–50%																					2
	1	2–5	28–40%																					1
	0	<2	<28%																					0
FM Face mask **HF** High flow **HFNP** High flow nasal prongs **NIV** Non invasive — **NP** Nasal prongs **NRM** Non re-breather **RA** Room air **T** Tracheostomy — Mode																								
Actual FiO_2 on Device Screen																								
High Flow L/min on Device Screen																								
Blood Pressure (mmHg) Score systolic BP			≥200																					≥200
			190s																					190s
			180s																					180s
			170s																					170s
			160s																					160s
			150s																					150s
			140s																					140s
			130s																					130s
			120s																					120s
			110s																					110s
			100s																					100s
			90s																					90s
			80s																					80s
			70s																					70s
			60s																					60s
Systolic BP score																								
Heart Rate (beats / min)	E		≥160																					E
	4		150s	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	4		140s	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	3		130s																					3
	2		120s																					2
			110s																					
	1		100s																					1
	0		90s																					0
			80s																					
			70s																					
			60s																					
			50s																					
	2		40s																					2
	E		30s																					E
Temperature (°C)	2		≥39.5																					2
			38.5–39.4																					
	1		38–38.4																					1
	0		37.5–37.9																					0
			37–37.4																					
			36.1–36.9																					
	1		35.1–36																					1
	2		34.1–35																					2
	4		≤34	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Behaviour and Consciousness If necessary, wake patient before scoring	4		Changing behaviour or New confusion	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	0		**A**lert																					0
	1		**V**oice																					1
	E		**P**ain																					E
			Unresponsive																					
TOTAL ADDS SCORE																								
Notification (most senior position, see legend on page 3)																								
Interventions (e.g. 'A')																								
Weight (kg)																								
Bowels																								
Initials																								

Score Legend

0	Score 0
1	Score 1
2	Score 2
3	Score 3
4	Score 4
E	Emergency Call

MRN			
Surname			
Given name(s)			
Date of Birth	/ /	Sex	Male / Female / Other

FOR OFFICE USE ONLY: Complete above or use client label

Indicate which systolic BP scoring preference is in use (Usual or Default). If the Usual systolic BP is selected, write the Usual systolic BP in the space provided:

☐ **Usual** systolic BP: ..... mmHg

☐ **Default** systolic BP: 120mmHg

Target Systolic BP (SMO / Registrar ONLY): mmHg

Name:		Name:	
Signature:		Signature:	
Designation:	Date:	Designation:	Date:

Circle the column showing the patient's Usual / Default / Target systolic BP

Actual BP	180s	170s	160s	150s	140s	130s	120s	110s	100s	90s	80s
≥200	4	4	4	4	4	4	4	4	4	4	4
190s	0	0	1	1	1	2	2	3	3	4	4
180s	0	0	0	0	1	1	2	2	3	3	4
170s	0	0	0	0	1	1	2	2	3	3	3
160s	1	0	0	0	0	0	1	1	2	2	2
150s	1	1	0	0	0	0	0	1	1	2	2
140s	1	1	1	0	0	0	0	0	1	1	1
130s	2	1	1	0	0	0	0	0	0	0	1
120s	2	2	1	1	0	0	0	0	0	0	0
110s	2	2	2	1	1	0	0	0	0	0	0
100s	3	3	2	2	2	1	1	0	0	0	0
90s	3	3	3	2	2	2	2	1	1	0	0
80s			4	4	4	4	4	4	4	1	0
70s											
60s				EMERGENCY CALL							

Score current systolic BP using the circled column

Time into PACU:	Airway type:	Time removed:
Time return to theatre: ☐ N/A		
Time into PACU: ☐ N/A	Airway type:	Time removed:

Escalation and Observation Plan

Total Score 0–2
- 5 minutely until alert and normal to rouse

Total Score 3–5
- Notify sedationist and request review within 10 minutes. Note review in Clinical Review Request section (page 1)
- Increase frequency of observations to 5 minutely

Total Score 6–7
- Notify proceduralist and request review within 5 minutes. Note review in Clinical Review Request section (page 1)
- Increase frequency of observations to 5 minutely

Total Score 8
- Place Emergency Call
- Begin initial life support interventions
- Advanced life support provider to attend patient
- Call for ambulance

Sedation Score (for patients receiving potentially sedating medication)

Patient **must** be woken to assess Sedation Score

Note: **DO NOT** add the Sedation Score to the ADDS Score. Follow actions below.

Date	
Time	
0	
1	
2	
3	

0 = Awake	• Continue to monitor patient's Sedation and Pain Score in accordance with individual monitoring plan
1 = Mild (easy to rouse, able to keep eyes open for 10 secs)	• Increase monitoring of Sedation and Pain Score • Recheck Sedation Score before administering potentially sedating medication
2 = Moderate (rouseable, but unable to keep eyes open for 10 secs)	• Ensure patient receives oxygen and monitor oxygen saturation • Withhold additional sedating medication (until medical review) • Notify Team Leader • Notify Medical Officer to review within 15 minutes (remain with patient until review) • Monitor Sedation and Pain Score (minimum 15 minutely) • If concerned, initiate Emergency Response
3 = Severe (difficult to rouse or un-rouseable)	• Initiate Emergency Response • Ensure patient receives oxygen and monitor oxygen saturation

Notification Legend

Document the letter in the *Notification* row on page 2 in appropriate time column.

N Nil required **TL** Team Leader **NUM** Nurse Unit Manager **P** Proceduralist **S** Sedationist

Emergency Call if:

- Any observation is in a purple area (**E**)
- Airway threat
- Respiratory or cardiac arrest
- New drop in O_2 saturation <90%
- Sudden fall in level of consciousness
- Seizure
- You are concerned about the patient but they do not fit the above criteria