


# R Notebook

Speciation long:

```
results <- list()
params_a <- flechette::create_params(
  speciation_initiation_rate = 0.2,
  speciation_completion_rate = 5.0,
  extinction_rate = 0.1,
  crown_age = 15,
  crown_age_sigma = 0.01,
  sampling_method = "youngest",
  mutation_rate = 1.0,
  sequence_length = 100,
  mcmc_length = 300000,
  tree_sim_rng_seed = 43,
  alignment_rng_seed = 43,
  beast2_rnd_seed = 44
)
results$a <- flechette::run(parameters = params_a)
ape::plot.phylo(results$a$species_tree)
```




```
results$a$nlts_diff <- nLTT::nlts_diff(tree = results$a$species_tree, results$a$trees)
tracerer::calc_ess(results$a$estimates$posterior, sample_interval = 1000)
```

```
## [1] 300.2815
```

Short:

```
params_b <- params_a
params_b$speciation_completion_rate <- 1000000.0
results$b <- flechette::run(parameters = params_b)
ape::plot.phylo(results$b$species_tree)
```



```
results$b$nlts_diff <- nLTT::nlts_diff(tree = results$b$species_tree, results$b$trees)
tracrer::calc_ess(results$b$estimates$posterior, sample_interval = 1000)
```

```
## [1] 301
```

Plot:

```
df <- data.frame(
  sim = c(
    rep("a", times = length(results$a$nlts_diff)),
    rep("b", times = length(results$b$nlts_diff))
  ),
  nlts = c(results$a$nlts_diff, results$b$nlts_diff)
)
ggplot2::ggplot(
  data = df,
  ggplot2::aes(x = nlts, fill = sim)
) + ggplot2::geom_histogram(alpha = 0.5, position = "identity", binwidth = 0.01)
```

