

PRD 14: Topological Semimetals: Weyl and Dirac Points from Symmetry

Pure Thought AI Challenge 14

Pure Thought AI Challenges Project

January 18, 2026

Abstract

This document presents a comprehensive Product Requirement Document (PRD) for implementing a pure-thought computational challenge. The problem can be tackled using only symbolic mathematics, exact arithmetic, and fresh code—no experimental data or materials databases required until final verification. All results must be accompanied by machine-checkable certificates.

Contents

Domain: Materials Science

Timeline: 5-7 months

Difficulty: Medium-High

Prerequisites: Band theory, topology, group theory, differential geometry

0.1 1. Problem Statement

0.1.1 Scientific Context

Topological semimetals are 3D materials where conduction and valence bands touch at isolated points (Weyl/Dirac) or lines (nodal lines) in the Brillouin zone, protected by topology and symmetry.

Weyl Points:

- **Band crossing:** Two non-degenerate bands touch at momentum k_W
- **Topological charge:** ± 1 chirality (monopole of Berry curvature)
- **Fermi arc surface states:** Connect projections of opposite-chirality Weyl points
- **Requires:** Either broken time-reversal (T) or inversion (I) symmetry

Dirac Points:

- **4-fold degeneracy:** Two Weyl points of opposite chirality pinned together by symmetry
- **Requires:** Both T and I symmetry present
- **Can split:** Into Weyl pairs when symmetry broken

Nodal Lines:

- **1D band crossing:** Bands touch along continuous curves in BZ
- **Protected by:** Mirror/glide symmetries + additional symmetries
- **Drumhead surface states:** Flat bands on surface

Key Properties:

- **Nielsen-Ninomiya theorem:** Weyl points come in pairs with opposite chirality ($C_i = 0$)
- **Bulk-boundary correspondence:** Fermi arcs connect Weyl points of opposite chirality
- **Anomalies:** Chiral anomaly causes magnetoresistance, optical responses

0.1.2 Core Question

Can we systematically construct tight-binding models with Weyl/Dirac points using ONLY symmetry constraints and topology—without materials databases or DFT?

Specifically:

- Given space group G , find minimal models hosting Weyl points
- Compute exact positions kW and chiralities CW
- Prove Fermi arc existence and topology
- Optimize nodal line geometries (linking, knotting)
- Certify robustness against disorder and perturbations
- Export 3D visualizations of Fermi surfaces and arcs

0.1.3 Why This Matters

Theoretical Impact:

- Completes topological classification for gapless systems
- Connects knot theory to condensed matter
- Tests bulk-boundary correspondence in 3D

Practical Benefits:

- Novel transport phenomena (chiral anomaly, nonlocal resistance)
- Topological quantum computing platforms
- Optoelectronic devices with unusual responses

Pure Thought Advantages:

- Weyl points are topological defects (no material parameters needed)
 - Chirality computed from Berry curvature (exact)
 - Symmetry analysis determines allowed positions
 - Surface states from semi-infinite geometry (pure mathematics)
-

0.2 2. Mathematical Formulation

0.2.1 Problem Definition

A **Weyl point** at momentum k_W is a point where :

- **Two bands touch:** $E(k_W) = E(k_W)$
- **Linear dispersion:** $E(k) = E_F \pm v_F |k - k_W|$ near k_W
- **Topological charge:**

$$C_W = \frac{1}{2} \int_{S} \mathbf{F}(\mathbf{k}) \cdot d\mathbf{S}$$

where S is a small sphere around k_W , and \mathbf{F} is Berry curvature

Weyl Hamiltonian (low-energy effective):

$$H(\mathbf{q}) = v_F \left(\mathbf{q} \cdot \boldsymbol{\sigma} + \gamma_5 \right)$$

where $\mathbf{q} = \mathbf{k} - \mathbf{k}_W$, $\boldsymbol{\sigma}$ are Pauli matrices, $\gamma_5 = \text{sign}(\det(v_F))$

Dirac Hamiltonian (4×4):

$$H(\mathbf{q}) = v_F \left(\mathbf{q} \cdot \boldsymbol{\gamma} + \gamma_5 \right)$$

where $\boldsymbol{\gamma}$ are 4×4 gamma matrices (two copies of Pauli matrices)

Nodal Line: 1D curve (t) in BZ where bands touch, characterized by:

- linking invariant (how nodal lines link)
- **Drumhead** flat surface state filling interior of projected loop

0.2.2 Certificate Requirements

- **Weyl Point Certificate:** Exact (k_W, C_W) pairs
- **Fermi Arc Topology:** Connectivity of arcs on surface BZ
- **Symmetry Protection:** Proof that Weyl/Dirac points cannot gap without breaking symmetry
- **Nodal Line Geometry:** Linking numbers, knot invariants
- **Surface State Verification:** Existence of arcs/drumheads

0.2.3 Input/Output Specification

Input:

```
1 from sympy import *
2 import numpy as np
3 from typing import List, Tuple, Callable
4
5 class SemimetalModel:
6     dimension: int # Must be 3D
7     space_group: int
8     time_reversal: bool # T symmetry
9     inversion: bool # I symmetry
```

```

10     hamiltonian: Callable[[np.ndarray], np.ndarray] # H(k), 3D momentum
11     num_bands: int
12 
```

Output:

```

1 class SemimetalCertificate:
2     model: SemimetalModel
3
4     # Weyl points
5     weyl_points: List[Tuple[np.ndarray, int]] # [(k_W, chirality), ...]
6     total_chirality: int # Should be 0 (Nielsen-Ninomiya)
7
8     # Dirac points
9     dirac_points: List[np.ndarray] # [k_D, ...]
10    dirac_splitting: Optional[List[Tuple]] # How Dirac      2 Weyl when
        symmetry broken
11
12    # Nodal lines
13    nodal_lines: List[Callable] # [      (t),      (t), ...]
        parameterized curves
14    linking_numbers: np.ndarray # L_ij = linking of _i , _j
15    knot_invariants: List[str] # Alexander polynomial, etc.
16
17    # Surface states
18    fermi_arcs: List[Tuple[np.ndarray, np.ndarray]] # [(k_start,
        k_end), ...] on surface
19    drumhead_states: Optional[np.ndarray] # For nodal lines
20
21    # Verification
22    berry_curvature_field: Callable[[np.ndarray], np.ndarray] # F(k)
23    surface_spectral_function: Callable[[np.ndarray], float] # A(k,
        E=0)
24
25    proof_of_topology: str # Mathematical derivation

```

0.3 3. Implementation Approach

0.3.1 Phase 1: Minimal Weyl Semimetal Model (Months 1-2)

Implement simplest Weyl semimetal (broken inversion):

```

1 import numpy as np
2 from scipy.linalg import eigh
3
4 def minimal_weyl_model(m: float, b: float) -> Callable:
5     """
6     Minimal 2-band model hosting a pair of Weyl points.
7
8      $H(k) = (b k_z + m) \sigma_x + b k_x \sigma_y + b k_y \sigma_z$ 
9
10    Weyl points at  $k_W = (0, 0, m/b)$  with opposite chirality.
11
12    Breaks inversion symmetry (due to  $k_z$  term).
13    """

```

```

14     def H(k: np.ndarray) -> np.ndarray:
15         kx, ky, kz = k[0], k[1], k[2]
16
17         sx = np.array([[0, 1], [1, 0]])
18         sy = np.array([[0, -1j], [1j, 0]])
19         sz = np.array([[1, 0], [0, -1]])
20
21         H_k = (b*kz + m)*sx + b*kx*sy + b*ky*sz
22
23         return H_k
24
25     return H
26
27 def find_band_crossings(H_func: Callable, k_range: Tuple[float, float,
28     float],
29     N_k: int = 50) -> List[np.ndarray]:
30     """
31     Find points in BZ where bands touch (gap closes).
32
33     Returns list of k-points where  $\min|E_i - E_j| < \text{threshold}$ .
34     """
35     kx_vals = np.linspace(-k_range[0], k_range[0], N_k)
36     ky_vals = np.linspace(-k_range[1], k_range[1], N_k)
37     kz_vals = np.linspace(-k_range[2], k_range[2], N_k)
38
39     crossing_points = []
40
41     for kx in kx_vals:
42         for ky in ky_vals:
43             for kz in kz_vals:
44                 k = np.array([kx, ky, kz])
45                 evals = np.linalg.eigvalsh(H_func(k))
46
47                 # Check for near-degeneracy
48                 gaps = [abs(evals[i+1] - evals[i]) for i in
49                     range(len(evals)-1)]
50                 min_gap = min(gaps)
51
52                 if min_gap < 1e-3: # Threshold for crossing
53                     crossing_points.append(k)
54
55     return crossing_points
56
57 def refine_weyl_point(H_func: Callable, k_initial: np.ndarray,
58     tol: float = 1e-10) -> np.ndarray:
59     """
60     Refine Weyl point position to machine precision.
61
62     Minimize  $\text{gap} = |E(k) - E(k)|$  around initial guess.
63     """
64     from scipy.optimize import minimize
65
66     def gap_function(k):
67         evals = np.linalg.eigvalsh(H_func(k))
68         return min([abs(evals[i+1] - evals[i]) for i in
69             range(len(evals)-1)])

```



```

40 """
41 Compute chirality C_W =      F      dS around Weyl point.
42
43 Integrate Berry curvature over sphere of radius r around k_W.
44 """
45 # Parameterize sphere: k = k_W + r(sin      cos      , sin      sin      ,
46     cos      )
47 theta_vals = np.linspace(0, np.pi, N_theta)
48 phi_vals = np.linspace(0, 2*np.pi, N_phi)
49
50 flux = 0
51
52 for theta in theta_vals:
53     for phi in phi_vals:
54         # Point on sphere
55         k = k_W + radius * np.array([
56             np.sin(theta)*np.cos(phi),
57             np.sin(theta)*np.sin(phi),
58             np.cos(theta)
59         ])
60
61         # Berry curvature
62         F = berry_curvature_3d(H_func, k, band_idx=0) # Lower band
63
64         # Outward normal
65         n = np.array([np.sin(theta)*np.cos(phi),
66             np.sin(theta)*np.sin(phi),
67             np.cos(theta)])
68
69         # F      dS (with Jacobian for spherical measure)
70         dS = radius**2 * np.sin(theta) *
71             (theta_vals[1]-theta_vals[0]) * (phi_vals[1]-phi_vals[0])
72         flux += np.dot(F, n) * dS
73
74     # Chirality is flux/(2 )
75     chirality = int(np.round(flux / (2*np.pi)))
76
77 return chirality

```

0.3.3 Phase 3: Fermi Arc Calculation (Months 3-4)

Compute surface states and Fermi arcs:

```

1 def surface_green_function(H_bulk: Callable, k_parallel: np.ndarray,
2     energy: float = 0, surface_normal: str = 'z',
3     N_layers: int = 100) -> np.ndarray:
4
5     """
6     Compute surface Green's function G( k_ , E) for semi-infinite
7     geometry.
8
9     Uses iterative method (transfer matrix or recursive Green's
10    function).
11
12    """
13    # For simplicity, use slab geometry with large N_layers

```

```

11     H_slab = construct_slab_hamiltonian(H_bulk, k_parallel,
12                                         surface_normal, N_layers)
13
14     # Green's function:  $G = (E - H + i\eta)^{-1}$ 
15     eta = 1e-3 # Small imaginary part
16     dim = H_slab.shape[0]
17
18     G = np.linalg.inv((energy + 1j*eta)*np.eye(dim) - H_slab)
19
20     # Project onto surface layer
21     num_orbitals = H_bulk(np.array([0, 0, 0])).shape[0]
22     G_surface = G[:num_orbitals, :num_orbitals]
23
24     return G_surface
25
26 def compute_surface_spectral_function(H_bulk: Callable, k_parallel:
27     np.ndarray,
28     energy: float = 0) -> float:
29     """
30     Surface spectral function  $A(k_{\parallel}, E) = -\text{Im Tr } G(k_{\parallel}, E)$ .
31
32     Peaks indicate surface states.
33     """
34     G_surf = surface_green_function(H_bulk, k_parallel, energy)
35
36     A = -np.imag(np.trace(G_surf))
37
38     return A
39
40 def map_fermi_arcs(H_bulk: Callable, weyl_points: List[Tuple],
41     N_k: int = 100) -> List[Tuple]:
42     """
43     Map Fermi arcs connecting Weyl point projections on surface BZ.
44
45     Returns list of arcs as (k_start, k_end) pairs.
46     """
47     # Project Weyl points onto surface (kz=0 plane)
48     weyl_projections = [(np.array([k[0], k[1]]), chi) for k, chi in
49         weyl_points]
50
51     # Compute spectral function on surface BZ at E=0
52     kx_surf = np.linspace(-np.pi, np.pi, N_k)
53     ky_surf = np.linspace(-np.pi, np.pi, N_k)
54
55     spectral_map = np.zeros((N_k, N_k))
56
57     for i, kx in enumerate(kx_surf):
58         for j, ky in enumerate(ky_surf):
59             k_par = np.array([kx, ky])
60             spectral_map[i, j] =
61                 compute_surface_spectral_function(H_bulk, k_par,
62                     energy=0)
63
64     # Identify arcs: high spectral weight curves connecting Weyl
65     # projections
66     # Use image processing / contour finding

```

```

61     from skimage.feature import peak_local_max
62
63     # Find high-intensity ridges (arcs)
64     arc_pixels = spectral_map > 0.5 * np.max(spectral_map)
65
66     # Trace paths (simplifiedfull implementation needs sophisticated
67     # tracking)
68     arcs = []
69     # ... (arc tracing algorithm)
70
71     return arcs

```

0.3.4 Phase 4: Dirac Points and Splitting (Months 4-5)

Study Dirac semimetals and symmetry breaking:

```

1  def dirac_semimetal_model(v_F: float = 1.0) -> Callable:
2      """
3      Minimal Dirac semimetal with both T and I symmetry.
4
5      4-band model: two Weyl points pinned together at k=0.
6
7       $H(k) = v_F (k_x \sigma_x + k_y \sigma_y + k_z \sigma_z)$ 
8
9      where  $\sigma_i$  are 4 x 4 Dirac matrices.
10     """
11     # Gamma matrices (one representation)
12     Gamma1 = np.kron(np.array([[0, 1], [1, 0]]), np.eye(2))
13     Gamma2 = np.kron(np.array([[0, -1j], [1j, 0]]), np.eye(2))
14     Gamma3 = np.kron(np.array([[1, 0], [0, -1]]), np.array([[0, 1], [1,
15     0]]))
16
17     def H(k: np.ndarray) -> np.ndarray:
18         kx, ky, kz = k[0], k[1], k[2]
19         return v_F * (kx*Gamma1 + ky*Gamma2 + kz*Gamma3)
20
21     return H
22
23  def split_dirac_into_weyl(H_dirac: Callable, breaking_term: str,
24     strength: float) -> Callable:
25     """
26     Split Dirac point into two Weyl points by breaking symmetry.
27
28     breaking_term:
29     - 'inversion': Add term breaking I symmetry
30     - 'time_reversal': Add term breaking T symmetry (magnetic field)
31     """
32     def H_split(k: np.ndarray) -> np.ndarray:
33         H_0 = H_dirac(k)
34
35         if breaking_term == 'inversion':
36             # Add k_z term or constant mass
37             delta_H = strength * np.kron(np.array([[1, 0], [0, -1]]),
38             np.eye(2))
39         elif breaking_term == 'time_reversal':
40             # Add magnetic field along z

```

```

39         delta_H = strength * np.kron(np.eye(2), np.array([[1, 0],
40             [0, -1]]))
41     else:
42         delta_H = np.zeros_like(H_0)
43
44     return H_0 + delta_H
45
46     return H_split
47
48 def trace_dirac_to_weyl_transition(H_dirac: Callable, strength_values:
49     np.ndarray):
50     """
51     Track how Dirac point splits into Weyl pair as symmetry-breaking
52     increased.
53     """
54     weyl_separation = []
55
56     for s in strength_values:
57         H_split = split_dirac_into_weyl(H_dirac, 'inversion', s)
58
59         # Find Weyl points
60         weyl_pts = find_band_crossings(H_split, k_range=(np.pi, np.pi,
61             np.pi))
62
63         if len(weyl_pts) >= 2:
64             # Distance between Weyl pair
65             dist = np.linalg.norm(weyl_pts[0] - weyl_pts[1])
66             weyl_separation.append(dist)
67         else:
68             weyl_separation.append(0)
69
70     return weyl_separation

```

0.3.5 Phase 5: Nodal Lines (Months 5-6)

Design semimetals with nodal line degeneracies:

```

1 def nodal_line_model(mirror_plane: str = 'xy') -> Callable:
2     """
3     Model with nodal line protected by mirror symmetry.
4
5     Bands touch along a circle in BZ (e.g.,  $k_z = 0$ ,  $k_x^2 + k_y^2 = r^2$ ).
6     """
7     def H(k: np.ndarray) -> np.ndarray:
8         kx, ky, kz = k[0], k[1], k[2]
9
10        sx = np.array([[0, 1], [1, 0]])
11        sy = np.array([[0, -1j], [1j, 0]])
12        sz = np.array([[1, 0], [0, -1]])
13
14        # Nodal line at  $k_z=0$ ,  $k_x^2 + k_y^2 = 1$ 
15        H_k = (kx**2 + ky**2 - 1)*sx + kz*sy
16
17        return H_k
18

```

```

19     return H
20
21 def extract_nodal_line(H_func: Callable, N_sample: int = 1000) ->
    Callable:
22     """
23     Find parameterization (t) of nodal line in BZ.
24
25     Returns: curve : [0, 2 ] (momentum space)
26     """
27     # Sample BZ to find where gap closes
28     gap_threshold = 1e-4
29     nodal_points = []
30
31     # ... (sample k-space, identify near-degeneracies)
32
33     # Fit smooth curve through nodal points
34     from scipy.interpolate import splprep, splev
35
36     tck, u = splprep([nodal_points[:, 0], nodal_points[:, 1],
37                       nodal_points[:, 2]], s=0)
38
39     def gamma(t):
40         return np.array(splev(t, tck))
41
42     return gamma
43
44 def compute_linking_number(gamma1: Callable, gamma2: Callable) -> int:
45     """
46     Compute Gauss linking number for two nodal lines.
47
48     
$$L = \frac{1}{4\pi} \int_0^{2\pi} \int_0^{2\pi} \left( \frac{\gamma_1'(s) \times \gamma_2'(t)}{|\gamma_1(s) - \gamma_2(t)|^3} \right) \cdot (\gamma_1(s) - \gamma_2(t)) \, ds \, dt$$

49
50     """
51     # Numerical integration
52     N_s, N_t = 100, 100
53     s_vals = np.linspace(0, 2*np.pi, N_s)
54     t_vals = np.linspace(0, 2*np.pi, N_t)
55
56     linking = 0
57
58     for s in s_vals:
59         for t in t_vals:
60             g1_s = gamma1(s)
61             g2_t = gamma2(t)
62
63             # Derivatives
64             ds = s_vals[1] - s_vals[0]
65             dt = t_vals[1] - t_vals[0]
66
67             g1_prime = (gamma1(s + ds) - gamma1(s)) / ds
68             g2_prime = (gamma2(t + dt) - gamma2(t)) / dt
69
70             diff = g1_s - g2_t
71             dist_cubed = np.linalg.norm(diff)**3
72
73             if dist_cubed > 1e-6:

```

```

72         integrand = np.dot(np.cross(g1_prime, g2_prime), diff)
73             / dist_cubed
74         linking += integrand * ds * dt
75
76     linking /= (4*np.pi)
77
78     return int(np.round(linking))

```

0.3.6 Phase 6: Certification and Database (Months 6-7)

Generate complete certificates and database:

```

1  def generate_semimetal_certificate(model: SemimetalModel) ->
2      SemimetalCertificate:
3      """
4      Generate complete certificate for topological semimetal.
5      """
6      cert = SemimetalCertificate(model=model)
7
8      # Find Weyl points
9      crossings = find_band_crossings(model.hamiltonian, k_range=(np.pi,
10         np.pi, np.pi))
11
12     weyl_points = []
13     for k_cross in crossings:
14         k_refined = refine_weyl_point(model.hamiltonian, k_cross)
15         chirality = compute_weyl_chirality(model.hamiltonian, k_refined)
16
17         if chirality != 0:
18             weyl_points.append((k_refined, chirality))
19
20     cert.weyl_points = weyl_points
21     cert.total_chirality = sum([chi for _, chi in weyl_points])
22
23     # Verify Nielsen-Ninomiya
24     assert cert.total_chirality == 0, "Chirality sum must be zero!"
25
26     # Fermi arcs
27     cert.fermi_arcs = map_fermi_arcs(model.hamiltonian, weyl_points)
28
29     # Nodal lines (if present)
30     # ... (detect and parameterize)
31
32     # Berry curvature field
33     cert.berry_curvature_field = lambda k:
34         berry_curvature_3d(model.hamiltonian, k, 0)
35
36     return cert
37
38 def export_semimetal_visualization(cert: SemimetalCertificate,
39     output_dir: Path):
40     """
41     Export 3D visualizations of Weyl points, Fermi arcs, nodal lines.
42     """
43     import matplotlib.pyplot as plt
44     from mpl_toolkits.mplot3d import Axes3D

```

```

41
42 # Plot Weyl points in 3D BZ
43 fig = plt.figure()
44 ax = fig.add_subplot(111, projection='3d')
45
46 for k_W, chi in cert.weyl_points:
47     color = 'red' if chi > 0 else 'blue'
48     ax.scatter(k_W[0], k_W[1], k_W[2], c=color, s=100, marker='o')
49
50 ax.set_xlabel('kx')
51 ax.set_ylabel('ky')
52 ax.set_zlabel('kz')
53 ax.set_title('Weyl Points (red=+1, blue=-1)')
54
55 plt.savefig(output_dir / 'weyl_points_3d.png')
56
57 # Plot Fermi arcs on surface
58 # ... (2D plot of arcs connecting Weyl projections)
59
60 def generate_semimetal_database() -> dict:
61     """
62     Database of topological semimetals.
63     """
64     database = {'models': []}
65
66     # Weyl semimetals
67     for config in ['minimal', 'type-II', 'multi-weyl']:
68         model = construct_weyl_model(config)
69         cert = generate_semimetal_certificate(model)
70
71         database['models'].append({
72             'type': 'Weyl',
73             'configuration': config,
74             'num_weyl_points': len(cert.weyl_points),
75             'weyl_positions': [k.tolist() for k, _ in cert.weyl_points],
76             'certificate_path': export_certificate(cert)
77         })
78
79     # Dirac semimetals
80     # ... (similar for Dirac, nodal line models)
81
82     return database

```

0.4 4. Example Starting Prompt

```

1 You are a condensed matter theorist specializing in topological
  semimetals. Design tight-binding
2 models with Weyl/Dirac points using ONLY symmetry and topologyno DFT
  or materials databases.
3
4 OBJECTIVE: Construct minimal Weyl semimetal, compute chiralities 1 ,
  verify Fermi arcs.
5

```

```

6 PHASE 1 (Months 1-2): Minimal Weyl model
7 - Implement 2-band  $H(k) = (bk_z + m) \tau_x + bk_x \tau_y + bk_y \tau_z$ 
8 - Find Weyl points at  $k_W = (0, 0, m/b)$ 
9 - Refine positions to machine precision
10
11 PHASE 2 (Months 2-3): Chirality calculation
12 - Compute Berry curvature  $F(k)$  on sphere around each Weyl point
13 - Integrate  $FdS$  to get chirality  $C_W = \pm 1$ 
14 - Verify Nielsen-Ninomiya:  $\sum_i C_i = 0$ 
15
16 PHASE 3 (Months 3-4): Fermi arcs
17 - Construct slab geometry (semi-infinite in  $z$ )
18 - Compute surface Green's function  $G(k_\parallel, E=0)$ 
19 - Map spectral function  $A(k_\parallel) = -\text{Im Tr } G$ 
20 - Identify arcs connecting Weyl projections
21
22 PHASE 4 (Months 4-5): Dirac semimetals
23 - Build 4-band Dirac model with  $T$  and  $I$  symmetry
24 - Split Dirac  $\pm 2$  Weyl by breaking  $I$ 
25 - Track Weyl separation vs perturbation strength
26
27 PHASE 5 (Months 5-6): Nodal lines
28 - Construct model with mirror-protected nodal line
29 - Parameterize nodal curve  $\gamma(t)$  in BZ
30 - Compute linking numbers for multi-component lines
31
32 PHASE 6 (Months 6-7): Database and visualization
33 - Generate certificates for 10 semimetal types
34 - Export 3D visualizations of BZ, Weyl points, arcs
35 - Classify by space group symmetry
36
37 SUCCESS CRITERIA:
38 - MVR: Minimal Weyl model with verified  $C_W = \pm 1$ 
39 - Strong: Fermi arcs computed and visualized
40 - Publication: Complete database + linking number calculations
41
42 VERIFICATION:
43 - Chirality exact:  $C_W \in \{-1, 0, +1\}$  (integer)
44 - Nielsen-Ninomiya:  $\sum_i C_i = 0$ 
45 - Fermi arcs connect opposite-chirality Weyl points
46 - Linking numbers computed for nodal lines
47
48 Pure topology + linear algebra. No DFT.
49 All results certificate-based with exact chirality computation.

```

0.5 5. Success Criteria

MVR (2 months): Minimal Weyl model, chirality verified

Strong (4-5 months): Fermi arcs, Dirac splitting

Publication (6-7 months): Complete database, nodal line linking

0.6 6. Verification Protocol

Automated checks: chirality sum, arc connectivity, symmetry verification.

0.7 7. Resources Milestones

References:

- Wan et al. (2011): "Topological Semimetal and Fermi-Arc Surface States"
- Burkov Balents (2011): "Weyl Semimetal in a Topological Insulator Multilayer"
- Fang et al. (2016): "Topological Nodal Line Semimetals"

Milestones:

- Month 2: Weyl model validated
 - Month 4: Fermi arcs mapped
 - Month 6: Nodal lines classified
-

0.8 8. Extensions

- **Type-II Weyl:** Tilted cones
 - **Hopf Nodal Links:** Knotted nodal lines
 - **Non-Hermitian Semimetals:** Exceptional points
-

End of PRD 14