



1.

The figures below show two sets of 25 histograms. One set was generated with random draws from the uniform discrete distribution (i.e. 1,2,3 or 4 with equal probability); for the other set, some of the histograms were "tweaked" a little. Which set of histograms contains the tweaked data?






- A. A
- B. B
- C. I can't tell.

2.

On August 11, 2014, 65 people picked a random number from 1 to 4 (inclusive). Here is the distribution of responses:



5 people picked 1, 13 picked 2, 35 picked 3, 12 picked 4.

Do students prefer the number 3 over other numbers? (See Step 2.) What is your (approximate) p value?

A. $p=0$

B. $p = 0.001$

C. $p = 0.05$

D. $p = 1e-4$

E. $p < 1e-5$

3.

What is your conclusion?

- A. Our class is biased toward the number 3.
- B. There is no bias in our class
- C. Our class is biased toward the number 3, but the result is not significant.
- D. We cannot draw a definite conclusion.

4.

What if we are not interested in the number 3 in particular, but in the question of whether any number gets picked more often than the others? What is your p-value then?

- A. 0
- B. 0.5
- C. 0.01
- D. 0.001
- E. $\leq 1e-5$

5.

What is your conclusion?

- A. Our class is biased towards one number
- B. There is no bias in our class
- C. Our class is biased toward one number, but the result is not significant.
- D. We cannot draw a definite conclusion.

6.

For 65 people "randomly" picking a number from 1 to 4 (inclusive) what is the smallest number of picks of any given number which would cause you to reject the null hypothesis (randomness) at $p < 0.05$?

7.

Which formulation of H_0 and H_A would you have picked? Can you think of another way to phrase H_0 and H_A that would affect the p value? What are the advantages and disadvantages?