




Figure 2A. SUID rates by postnatal age (PNA). Weekly sex-specific rates across postnatal weeks 1–52.




Figure 2B. SUID rates by post-conception age (PCA). Weekly sex-specific rates across PCA weeks 36–75.

Figure 2. *SUID rates by age scale.* Weekly SUID rates per 100,000 live births by postnatal age (PNA, weeks 1–52) and post-conceptional age (PCA, weeks 36–75), shown for combined, male, and female infants. The PCA panel includes reference lines at 43 and 47 weeks and a shaded 43–46 week peak interval. Panels share y-axis limits to permit visual comparison.




Figure 3A. Observed and predicted SUID case counts by sex and PCA.




Figure 3B. Male-to-female observed and predicted difference of SUID case-count by PCA.




Figure 3C. Male-to-female observed and predicted log difference of SUID case-count by PCA.

Figure 3. Developmental timing of sex divergence in SUID risk. Observed weekly SUID counts and model-estimated trajectories by sex across PCA weeks 36–75. Panels display fitted curves and sex contrasts; observed weekly counts (points) and fitted mean curves (lines) for males and females with 95% confidence band.




Figure 4A. Early Window (< 43 post-conceptual weeks)




Figure 4B. Late Window (> 47 post-conceptual weeks)

Figure 4. Incidence rate ratios from negative binomial models in early and late developmental windows. Forest plots of adjusted IRRs (95% CI) from models fitted separately in the early (36–42 PCA weeks) and late (≥47 PCA weeks) developmental windows. Models include PCA (per week), sex, maternal smoking, preterm birth, and prespecified interaction terms.