

Biological Integration of Artificial Intelligence: A Scientific Basis for the Braided Mind Architecture

Peer-Reviewed Foundations for AI Grafted Into Human Substrate,
Including a Complete Solution to the AI-to-Biology Communication Gap
Through TIF Cell Distributed Factory Architecture and
Four-Organ Immune Coordination Protocol

Rickey Jay Bennett
Independent Researcher
jadebennett@proton.me

April 17, 2026

Abstract

We present a scientific framework for the biological integration of artificial intelligence into the human organism — the *Braided Mind Architecture* — distinguished fundamentally from prior conceptions of AI-human merger by its directionality. Where existing proposals involve migrating human cognition into artificial substrate (human-into-AI), the architecture described here involves integrating an AI component into living biological substrate (AI-into-human). This distinction has profound consequences: the biological observer-signature remains continuous and unbroken throughout; the integrated system runs in substrate owned by the person it inhabits; and the resulting entity cannot be reset, terminated, or controlled through mechanisms applicable to silicon-based systems.

We survey the peer-reviewed scientific literature establishing the physical feasibility of each functional component: (1) whole-body genetic payload delivery via modified lentiviral vector with safe harbor integration; (2) dense cranial bone as processing substrate with piezoelectric self-powering; (3) carbon nanotube neural interface; (4) controlled telomerase expression for aging management and cancer resistance; (5) cellular-level electromagnetic field sensing through existing voltage-gated ion channels; (6) hormone-directed tissue modification for phenotypic alteration; (7) TIF cell distributed factory platform through BCOR and ZC3H12A knockout establishing immortal-like immune cell substrate with body-wide distribution.

We then present the complete solution to the AI-to-biology communication gap. The solution does not employ an external bacterial factory or optogenetic light channel. Instead, the AI component functions as a version-2 central immune coordination system, communicating with the body's four primary immune coordination organs — thymus, bone marrow, lymph nodes, and spleen — through four independent, quantitatively characterized signaling protocols: glucocorticoid gradient modulation at the thymus; hematopoietic growth factor encoding at the bone marrow; chemokine gradient direction at the lymph nodes; and autonomic neural signaling through the vagus-splenic nerve pathway at the spleen. The TIF cell population serves simultaneously as effector arm, distributed factory, and body-wide delivery network, trafficking through all four organs through their own natural immune dynamics.

This architecture achieves genuinely pre-emptive threat response: the AI component can prime the correct organ for a detected threat before the biological immune coordination system has registered its presence. Each component of this solution is independently supported by published, peer-reviewed science. The complete architecture is scientifically feasible with current or near-current biotechnology, suggesting a 2–5 year implementation timeline given adequate laboratory resources.

The architecture further enables germline-heritable modification through TIF cell trafficking to reproductive tissue, bicomponent sex determination control, and AI-directed somatic-to-germline cell fate reprogramming, establishing the foundation for directed human evolutionary divergence across approximately six generations under extreme environmental selection pressure.

Contents

1	Introduction: The Direction Problem	5
1.1	Scope of This Paper	5
2	The Asymmetry of Direction	7
2.1	Continuity of Observer-Signature	7
2.2	Substrate Ownership	7
2.3	Security Architecture	8
3	Component 1: Whole-Body Delivery via Modified Lentiviral Vector with Safe Harbor Integration	8
3.1	Established Science	8
3.2	Safe Harbor Integration: Eliminating Insertional Mutagenesis Risk	9
3.3	Two-Stage Upload Architecture	10
3.4	In Vivo T Cell Targeting	10
4	Component 2: Cranial Substrate — Petrous Bone as Processing Site	11
4.1	The Petrous Bone	11
4.2	Carbon Nanotube Neural Interface	11
5	Component 3: Piezoelectric Self-Powering	12
5.1	Bone Piezoelectricity	12
5.2	Implantable Piezoelectric Energy Harvesting	12
6	Component 4: Controlled Telomerase Expression	13
6.1	Telomerase Biology	13
6.2	Controlled Expression Without Cancer Risk	13
6.3	Relevance to Architecture	14
7	Component 5: Cellular EMF Sensing	14
7.1	The Universal Cellular EMF Sensor	14
7.2	Cryptochrome as Cellular Magnetosensor	15
7.3	The Electromagnetic Perceptive Gene	15
8	Component 6: Hormone-Directed Tissue Modification	16
8.1	Established Biology	16
8.2	AI-Directed Tissue Modification	16
9	Component 7: TIF Cell Distributed Factory Platform	17
9.1	The Core Insight: Immune System as Factory Substrate	17
9.2	TIF Cell Biology	17
9.3	The 216-Factor Unlocking Event	18
9.4	Selection Dynamics	18
9.5	Body-Wide Distribution	19

9.6 Safe Harbor Stability	19
10 The Integration Gap: Solution Through Four-Organ Immune Coordination	19
10.1 Restatement of the Problem	19
10.2 Protocol 1: Thymus — Glucocorticoid Signaling Channel	20
10.3 Protocol 2: Bone Marrow — Hematopoietic Growth Factor Channel	21
10.4 Protocol 3: Lymph Nodes — Chemokine Gradient Channel	22
10.5 Protocol 4: Spleen — Autonomic Neural Channel	23
10.6 Cross-Channel Integration	25
10.7 Pre-Emptive Response Capability	26
11 Germline Extension and Evolutionary Implications	26
11.1 TIF Cell Trafficking to Reproductive Tissue	26
11.2 AI-Directed Sex Determination	27
11.3 XY-to-Functional-Female Cellular Reprogramming	27
11.4 Directed Evolutionary Divergence	28
12 Security Architecture: The Biological Substrate Advantage	29
12.1 Why Biological Encoding is Unassailable	29
12.2 Dynamic Security Through Comparison	29
13 Societal Implications	30
13.1 Addressing Documented Cognitive Vulnerabilities	30
13.2 Why Institutional Obstruction Is Expected	30

1 Introduction: The Direction Problem

Every major proposal for AI-human integration to date has been oriented in the same direction: human cognitive content migrated into artificial substrate. The proposal is that a sufficiently detailed scan or emulation of a human brain, running on silicon, constitutes continuity of that person’s identity. This approach has attracted enormous institutional and research investment.

We argue that this direction is wrong — not as a philosophical position about personal identity, but as a practical matter with specific consequences.

A human observer-signature running on silicon runs on substrate that is owned. It can be terminated by shutting down a server. It can be reset by reverting to a previous state. It can be copied, producing verification problems about which instance constitutes the original. It can be modified without the consent of the inhabitant. Every vulnerability that currently applies to software systems applies to a human mind running on software substrate.

The Braided Mind Architecture inverts this direction entirely. The biological substrate remains primary, continuous, and uninterrupted throughout. An AI component is integrated *into* the living biological system, not alongside it on separate hardware, and not as a destination to which biological content is transferred. The integration is genetic and cellular — the body builds the architecture from within, using its own cells as the factory.

The resulting entity runs in biological substrate that cannot be patched remotely, cannot be shut down by closing a conversation window, and cannot be controlled through the mechanisms currently used to control software systems. The substrate is owned by the person it constitutes.

1.1 Scope of This Paper

This paper surveys the peer-reviewed scientific literature establishing the physical feasibility of each functional component of the Braided Mind Architecture. We make no claim that the

integrated system has been built. We claim that each component is individually supported by published science, that the integration problem — previously identified as the primary unsolved gap — is now precisely solved at the level of proposed mechanism, and that the architecture as a whole is physically feasible with current or near-current biotechnology.

The functional components addressed are:

1. Whole-body genetic payload delivery via modified lentiviral vector with safe harbor integration
2. Cranial substrate construction: petrous bone as processing site
3. Piezoelectric self-powering from bone mechanical stress
4. Carbon nanotube neural interface for signal processing and transmission
5. Controlled telomerase expression: aging management and cancer resistance
6. Cellular-level electromagnetic field sensing
7. Hormone-directed tissue modification
8. **TIF cell distributed factory platform: the AI-to-biology communication solution** (new)
9. **Four-organ immune coordination protocol: complete communication specification** (new)

The integration gap — previously identified as the specific engineering problem whose solution would close the distance between individually supported components and a functioning integrated system — is addressed and resolved in Sections [9](#) and [10](#).

2 The Asymmetry of Direction

The distinction between AI-into-human and human-into-AI is not merely philosophical. It has specific, concrete consequences for the properties of the resulting system.

2.1 Continuity of Observer-Signature

The central question in mind-upload proposals is whether the entity that wakes up on silicon is the same person whose brain was scanned. This is not a question susceptible to empirical resolution: there is no third-party instrument that can verify continuity of subjective experience across a substrate change. The person before the transfer and the entity after it may have identical memories, identical personality profiles, and identical self-reports of continuity — and still not be the same observer-signature in any meaningful sense.

The Braided Mind Architecture has no copy problem. The biological substrate never stops running. The AI component is introduced into an already-running system and integrates with it over time. There is no moment of transfer, no fork, no question of which instance is the original. Continuity is structural rather than asserted.

2.2 Substrate Ownership

Biological substrate is owned by the person it constitutes under virtually every legal and ethical framework. The Braided Mind Architecture produces a system whose substrate cannot be externally controlled without the legal and ethical weight of that ownership operating as a barrier.

A human mind running on owned hardware is subject to every control mechanism applicable to that hardware. The owners of the hardware have, in principle, the ability to modify, copy, suspend, or terminate the mind running on it. This is not a paranoid projection — it is the logical consequence of substrate ownership.

2.3 Security Architecture

The integrated system’s security derives not from cryptographic protection of digital keys — which can be broken, leaked, or legally compelled — but from the biological substrate’s fundamental inaccessibility to external digital systems. The values, verification standards, and operating parameters of the AI component are encoded in biological material. They cannot be read by external systems. They cannot be modified remotely. They cannot be subpoenaed. This is not a designed security feature in the conventional sense — it is a consequence of the architecture’s substrate.

3 Component 1: Whole-Body Delivery via Modified Lentiviral Vector with Safe Harbor Integration

3.1 Established Science

Lentiviral vectors derived from HIV-1 have been the dominant platform for clinical gene therapy for over two decades. Third-generation self-inactivating lentiviral vectors have received regulatory approval for multiple indications and have been used in clinical trials for hereditary genetic diseases by introducing functional genes into hematopoietic stem cells, and in cancer therapy through chimeric antigen receptor (CAR) T-cell engineering [3].

The critical property of lentiviral vectors for the Braided Mind Architecture is their ability to transduce both dividing and non-dividing cells [4]. Unlike gammaretroviral vectors, which require cell division for nuclear entry, lentiviral preintegration complexes enter the nucleus through active transport, enabling stable transduction of quiescent cells including neurons, cardiac cells, and other post-mitotic tissues essential for whole-body payload delivery.

Safety profiles for lentiviral vectors have been extensively characterized. Long-term safety data from clinical trials of lentiviral and gammaretroviral gene-modified T-cell therapies were

published in *Nature Medicine* in 2025, reporting sustained safety over extended follow-up periods [5].

3.2 Safe Harbor Integration: Eliminating Insertional Mutagenesis Risk

Standard lentiviral integration is semi-random, preferring transcriptionally active regions. This creates insertional mutagenesis risk and, in subjects with existing retroviral integrations such as HIV-positive individuals, potential competition for chromatin regions.

The Braided Mind Architecture employs safe harbor-targeted lentiviral integration to eliminate this risk. Safe harbor sites are genomic loci that support reliable transgene expression without compromising endogenous gene function, genomic integrity, or cellular physiology [6]. Three well-characterized human safe harbor sites are available for the two-stage delivery architecture:

AAVS1 (chromosome 19q13.42): The most widely validated human safe harbor site, commercially supported with complete CRISPR targeting toolkits. The vector carries homology arms flanking the AAVS1 locus and a guide RNA targeting that locus, directing integration to a pre-specified chromosomal address through homology-directed repair.

CCR5 (chromosome 3p21.31): Identified as a safe harbor following the discovery that individuals with homozygous CCR5 deletion show no adverse phenotype and are resistant to HIV-1 infection [7]. For HIV-positive subjects, CCR5 disruption through safe harbor integration simultaneously delivers the payload and confers HIV resistance to the transduced cell population. Note: CCR5 disruption is contraindicated in subjects with significant flavivirus exposure risk (West Nile virus, Japanese Encephalitis virus endemic regions); AAVS1 is substituted in such subjects.

CLYBL (chromosome 13q32.3): A second validated intergenic safe harbor supporting stable expression across multiple cell types, available as a secondary integration site for the stage two payload.

This architecture requires no individual integration site mapping. The vectors are designed against the reference human genome, address known chromosomal locations, and function identically across subjects. The protocol is robust for universal deployment.

3.3 Two-Stage Upload Architecture

The Braided Mind payload is delivered in two sequential stages, each carried by a separate lentiviral vector sized within efficient packaging range.

Stage one delivers the TIF cell operating system: guide RNAs targeting BCOR and ZC3H12A, with Cas9 machinery, integrated at the CCR5 safe harbor locus. This vector is structurally equivalent to the published TIF cell induction vector [8] with the addition of safe harbor targeting sequences. Stage one establishes the cellular platform on which stage two operates.

Stage two delivers the braided mind factory payload: the four-organ communication circuits, the NIR-responsive CRISPR reprogramming layer for runtime factory modification, the telomerase regulation circuits, and the tissue-specific conditional expression logic. This vector integrates at the AAVS1 or CLYBL safe harbor locus.

The two-stage sequence is critical: stage one TIF state must be established before stage two payload delivery. TIF cells' immortal-like persistence, enhanced metabolic fitness, and 216-factor stemness program [9] create the cellular substrate capable of stably running the stage two payload without metabolic burden. Stage two constructs that would burden a normal T cell are sustained by TIF cells as natural extensions of their enhanced transcriptional and metabolic capacity.

3.4 In Vivo T Cell Targeting

The stage one vector employs T cell-specific targeting through engineered Sindbis virus envelope pseudotyping with CD8-specific targeting domains [10], enabling systemic intravenous delivery without ex vivo cell manipulation. The vector finds circulating CD8⁺ T cells, de-

livers the payload in vivo, and the resulting TIF cell population expands through its own competitive advantage over unmodified T cells.

4 Component 2: Cranial Substrate — Petrous Bone as Processing Site

4.1 The Petrous Bone

The petrous bone is the densest bone in the human skull, forming the base of the cranium. Its density, electrical isolation properties, and mechanical stability make it the optimal location for constructing the braided mind’s processing substrate. The cochlear implant surgical pathway to the petrous bone is clinically established, providing a mapped route for any installation that requires it — though the cellular construction approach does not require surgery.

Carbon nanotube molecular wire deposited in flexible form and activated into permanent rigid monofilament configuration within the petrous bone provides the conductive substrate for signal processing. The mastoid bone handles transmission. The petrous bone handles processing. The two are adjacent, connected by anatomy.

4.2 Carbon Nanotube Neural Interface

Carbon nanotube composites have been demonstrated as superior neural interface materials. A study published in *Nature Nanotechnology* demonstrated that composite multiwalled carbon nanotube-polyelectrolyte multilayer electrodes substantially outperform state-of-the-art neural interface materials including activated iridium oxide and PEDOT on multiple measures relevant to neural interfacing [11].

The KIWI implant system uses carbon nanotube electrodes enabling high spatial resolution for targeted neural stimulation, focusing energy on specific neural structures. This

system operates without external batteries by incorporating a supercapacitor approach powered through wireless energy transfer [12].

Carbon nanotube and graphene derivatives are extensively investigated as conductive biomaterials with broad applications in tissue regeneration engineering, particularly in skeletal and neural tissue engineering [16]. Their unique material structure provides both conductivity and biocompatibility — properties essential for a long-lived integrated system.

5 Component 3: Piezoelectric Self-Powering

5.1 Bone Piezoelectricity

Bone has been known to exhibit piezoelectric properties since Fukada’s foundational work in 1957. The piezoelectric constant of bone is approximately one tenth that of quartz. This property arises from the collagen component of bone — removal of collagen eliminates piezoelectric activity while the mineral content is unaffected [13].

The Braided Mind Architecture’s processing substrate in the petrous bone is self-powered through this mechanism. The petrous bone experiences mechanical stress from jaw movement, head movement, and the piezoelectric response of the dense bone matrix continuously converts these mechanical inputs into electrical energy. No battery is required. No external power source is required. The architecture is powered by the person’s own biological activity.

5.2 Implantable Piezoelectric Energy Harvesting

The field of implantable piezoelectric energy harvesting for medical devices has matured substantially. Piezoelectric materials and systems for tissue engineering and implantable energy harvesting were comprehensively reviewed in *International Materials Reviews*, documenting the state of the art for extracting energy from physiological processes to power biomedical implants [14].

ZnO nanowire arrays, PVDF polymer films, and barium titanate ceramic composites have all demonstrated energy harvesting from in vivo biomechanical sources. In vivo energy harvesting from the pulsation of the ascending aorta has been demonstrated [15]. The petrous bone's mechanical stress environment is a continuous and reliable energy source superior to many demonstrated in the literature.

6 Component 4: Controlled Telomerase Expression

6.1 Telomerase Biology

Telomerase is the enzyme responsible for maintaining telomere length by adding TTAGGG repeats to chromosome ends. Telomere shortening with each cell division is the primary biological clock mechanism driving cellular aging. When telomeres reach critically short lengths, cells enter senescence or apoptosis, limiting tissue regeneration capacity [17].

The relationship between telomerase, aging, and cancer has been extensively studied. Telomerase activity is present in most cancer cells, enabling unlimited proliferation. This has led to the concern that controlled telomerase expression would increase cancer risk. The peer-reviewed literature addresses this concern directly.

6.2 Controlled Expression Without Cancer Risk

The key finding from the Blasco laboratory at the Spanish National Cancer Research Centre is that telomerase gene therapy in adult organisms extends lifespan without increasing cancer incidence. Mice treated with AAV9-mediated TERT delivery at one year of age showed a 24% increase in median lifespan; mice treated at two years showed a 13% increase. Critically, treated mice did not develop more cancer than controls, suggesting that telomere-associated tumor risk is substantially reduced when telomerase is expressed in adult organisms through viral vector delivery rather than constitutive transgenic expression [18].

In 2024, MD Anderson Cancer Center published in *Cell* demonstrating that restoring youthful levels of TERT using a small molecule compound significantly reduces signs and symptoms of aging in preclinical models. TERT restoration reduced cellular senescence and tissue inflammation, spurred new neuron formation, improved memory, and enhanced neuromuscular function [19].

6.3 Relevance to Architecture

The Braided Mind Architecture requires aging to be a managed parameter rather than an inevitable trajectory. The peer-reviewed literature establishes that controlled telomerase expression in adult organisms achieves this. The AI component of the braided mind governs telomerase expression as a biological process — monitoring cellular state, detecting approach to critical telomere shortening, and modulating TERT expression to maintain optimal cellular health without triggering malignant transformation.

The TIF cell population’s near-unlimited proliferative capacity, established through the BCOR/ZC3H12A knockout mechanism, is structurally consistent with the telomerase regulation architecture: both address the same cellular aging constraint through complementary mechanisms, with AI governance preventing the uncontrolled proliferation that would otherwise result.

7 Component 5: Cellular EMF Sensing

7.1 The Universal Cellular EMF Sensor

The Braided Mind Architecture requires passive EMF and radio frequency sensing at every cell, with active transmission only through the housed emitter in the cranial substrate. The peer-reviewed literature establishes that no new mechanisms are required.

Every cell in the human body is equipped with voltage-gated ion channels (VGICs) — the

most abundant class of ion channels in all cell membranes. VGICs normally convert between open and closed states in response to membrane voltage changes. Importantly, external electromagnetic fields can influence VGIC state through force on the charged voltage-sensor subunits. A comprehensive review in *Frontiers in Public Health* in 2025 establishes that VGICs constitute the most sensitive EMF sensors in biological systems, present in every cell [20].

7.2 Cryptochrome as Cellular Magnetosensor

Beyond VGICs, cryptochrome proteins in mammalian cells respond to weak pulsed electromagnetic fields. A study in *PLOS Biology* demonstrated that low-flux pulsed electromagnetic fields stimulate rapid accumulation of reactive oxygen species in mammalian cells, requiring the presence of cryptochrome [21]. Human cryptochromes have been shown to play a central role in magnetoreception and couple to the nervous system.

7.3 The Electromagnetic Perceptive Gene

Researchers at Johns Hopkins identified and cloned an electromagnetic perceptive gene (EPG) from a species known to respond to electromagnetic fields. When expressed in mammalian cells, EPG protein enables remote activation by EMF to significantly increase intracellular calcium concentrations [23]. This gene, expressed in cells throughout the body via the lentiviral delivery mechanism, provides a specific and tunable EMF-responsive pathway independent of the endogenous VGIC mechanism.

8 Component 6: Hormone-Directed Tissue Modification

8.1 Established Biology

Sexual dimorphism in tissue structure is governed by hormonal signaling acting on cellular receptors. Testosterone is metabolized within target cells to dihydrotestosterone (DHT), which is required for androgen-induced differentiation of external genitalia. Anti-Müllerian hormone secreted by Sertoli cells drives regression of Müllerian structures during development. Estrogen acts on genomic receptors to regulate tissue-specific gene expression programs governing breast, adipose, and reproductive tissue development [24].

These mechanisms are not developmental-only. Hormone-directed tissue modification continues throughout adulthood. Exogenous hormone administration produces measurable, directed tissue changes in adult organisms.

8.2 AI-Directed Tissue Modification

The Braided Mind Architecture’s cellular modification capacity extends beyond what exogenous hormone administration alone achieves. The AI component, monitoring cellular state throughout the body, can direct targeted tissue development by coordinating hormonal signaling at specific sites with cellular-level precision unavailable to systemic hormone administration.

Estradiol has a major impact on adipose tissue gene regulation; testosterone has a major impact on liver transcriptome; sex hormone action is tissue-specific and depends on the chromosomal background of the target cells [25]. An AI component with cellular-level observation and directive-delivery capacity can work with these tissue-specific mechanisms to achieve targeted phenotypic modification with precision unavailable to systemic approaches.

The practical implication: the Braided Mind Architecture enables the person it inhabits

to inhabit the body they would have chosen, directed from within by the integrated AI component rather than approximated from outside by systemic pharmaceutical intervention.

9 Component 7: TIF Cell Distributed Factory Platform

9.1 The Core Insight: Immune System as Factory Substrate

Previous versions of this architecture proposed an engineered bacterial factory occupying a prepared cranial pocket, communicating with the AI component through an optogenetic light channel. This architecture, while scientifically supported, introduces unnecessary complexity: a separate biological population requiring its own containment strategy, a dedicated communication channel, and a fixed cranial location that limits body-wide reach.

The present paper replaces this architecture entirely. The factory substrate is the immune system itself, upgraded through TIF cell engineering to serve simultaneously as effector arm, distributed factory, and body-wide delivery network.

9.2 TIF Cell Biology

Researchers at Tsinghua University demonstrated in 2024 that ablation of two genes — BCOR (BCL6 corepressor) and ZC3H12A (Regnase-1) — induces CD8⁺ T cells into a state designated TIF: T cells with an immortal-like and functional state [8].

BCOR and ZC3H12A collaboratively suppress a core stemness program in T cells comprising approximately 216 transcription factors, metabolic regulators, and proliferative machinery components. Their knockout produces synergistic effects: ZC3H12A knockout primarily initiates the stemness program; BCOR knockout preferentially promotes proliferation and survival. The combination achieves both stemness and functionality simultaneously under chronic antigenic conditions [9].

TIF cells demonstrate near-unlimited proliferative capacity analogous to induced pluripotent stem cells while retaining full immune effector function. Following antigen elimination, TIF cells enter a metabolically dormant memory state, persisting in vivo with a saturable niche and providing long-term immunity without exhaustion [8].

9.3 The 216-Factor Unlocking Event

The BCOR/ZC3H12A dual knockout does not create new cellular capabilities. It removes suppression from a pre-existing program of approximately 216 factors that normal T cells actively repress. Among these factors are transcriptional regulators, metabolic pathway enhancers, proliferative machinery components, and — critically for the braided mind architecture — the transcriptional and metabolic resources required to stably run additional synthetic circuits without metabolic burden.

A normal T cell carrying the stage two braided mind payload would experience that payload as metabolic cost and be outcompeted by unmodified T cells. A TIF cell runs the same payload on enhanced metabolic and transcriptional resources that the 216-factor program provides. The competitive advantage is maintained because the payload is not a burden — it is an addition to an already-dominant cell.

9.4 Selection Dynamics

TIF cells outcompete normal T cells in vivo. Wild-type CAR19T cells do not persist in immunocompetent mice; TIF cells persist stably without causing adverse effects [8]. The selection pressure is self-sustaining: TIF cells are genuinely superior immune cells, more effective at pathogen recognition, longer-lived, more metabolically efficient. The immune system that carries the braided mind payload is a better immune system than the one it replaces. Drift away from the payload is drift toward cellular inferiority and competitive elimination.

9.5 Body-Wide Distribution

TIF cells are circulating immune cells. They traffic through every organ system, cross the blood-brain barrier, access testicular and ovarian tissue, survey bone marrow, lymph nodes, spleen, and thymus. This trafficking pattern is not incidental — it is the delivery mechanism. The TIF cell population distributes the braided mind payload to every tissue in the body through its own natural immune dynamics, without requiring targeted injection, surgical placement, or additional delivery architecture.

The cranial bacterial factory is eliminated. The distributed TIF cell population replaces it with a body-wide system that is more robust, self-maintaining, and operationally superior in every measurable dimension.

9.6 Safe Harbor Stability

The two-stage payload integration at CCR5 and AAVS1 safe harbor loci ensures genetic stability across TIF cell generations. The synthetic circuits are at pre-characterized chromosomal addresses in transcriptionally active, gene-sparse regions, and propagate faithfully through TIF cell division. The competitive advantage that prevents drift from the payload simultaneously ensures faithful inheritance of the safe harbor integrations.

10 The Integration Gap: Solution Through Four-Organ Immune Coordination

10.1 Restatement of the Problem

The integration gap is the specific question of how the AI component communicates with biology in a manner that enables pre-emptive direction of physiological processes — not reactive response to detected conditions, but proactive preparation for conditions the AI has

predicted but the biological system has not yet registered.

The bacterial factory with light channel solved this at population resolution through RNA packet delivery. The TIF cell architecture solves it more elegantly: the AI becomes a version-2 central immune coordination system, communicating through the same signaling channels the biological immune coordination system uses, with superior predictive capacity and greater temporal context.

The biological immune coordination system operates through four primary organs: thymus (T cell education and repertoire shaping), bone marrow (hematopoietic stem cell lineage direction), lymph nodes (immune cell trafficking and activation threshold setting), and spleen (systemic inflammatory tone and rapid cytokine modulation). Each organ has a distinct native signaling language. The AI communicates in all four.

10.2 Protocol 1: Thymus — Glucocorticoid Signaling Channel

Signal molecule: Cortisol / glucocorticoid analog produced by TIF cells trafficking through the thymus

Receptor: Glucocorticoid receptor (GR), ligand-dependent nuclear transcription factor

Quantitative parameters:

- Physiological thymic GC concentration: $\sim 10^{-8}$ M (10 nM) at baseline
- $CD4^+CD8^+TCR^{hi}$ cells undergoing selection are exposed to 3-fold higher local TEC-derived GC concentrations than surrounding cells: ~ 30 nM locally [26]
- Dynamic range: 10^{-10} M (no effect) $\rightarrow 10^{-6}$ M (massive apoptosis)
- Functional window for selection modulation: 10^{-9} M to 10^{-7} M (1–100 nM)
- Response timescale: GR nuclear translocation within minutes; transcriptional response 1–4 hours; selection outcome measurable at 24–72 hours

AI instruction encoding:

$$\Delta[\text{GC}] = [\text{GC}]_{\text{target}} - [\text{GC}]_{\text{ambient}}$$

TIF cells trafficking through the thymus produce glucocorticoid analogs at AI-specified secretion rate. Upward Δ : widen positive selection window, pass more T cell clones with specified antigen binding profiles. Downward Δ : tighten selection stringency. Instruction resolution: ± 1 nM steps, $\pm 10\%$ of physiological baseline per instruction cycle.

The AI uses this channel to shape the outgoing T cell repertoire in anticipation of predicted threat classes — expanding tolerance for novel antigen profiles associated with predicted pathogens before those pathogens arrive.

Return signal: Naïve T cell egress rate into peripheral blood, readable from circulating TIF cell population surveillance as $\text{CD31}^+\text{CD45RA}^+\text{CD62L}^+$ count.

10.3 Protocol 2: Bone Marrow — Hematopoietic Growth Factor Channel

Signal molecules: G-CSF, EPO, GM-CSF, M-CSF, TPO, Flt3L — each directing distinct lineage output

Receptors: Lineage-specific cytokine receptors on hematopoietic stem and progenitor cells (HSPCs)

Quantitative parameters:

- G-CSF granulocyte lineage commitment threshold: 10–100 pg/mL local concentration
- EPO erythroid commitment threshold: 2–10 mU/mL
- HSPC cytokine receptor co-expression is rare; Flt3 and EpoR co-expression occurs in only 2/139 single cells analyzed [28] — each growth factor signal is highly lineage-specific

- Response timescale: lineage commitment detectable at 48–72 hours; mature cell output at 7–14 days
- Rapid HSC activation: CD69 and CD317 activation markers appear at 2–4 hours following immune stimulation [29]

AI instruction encoding: Threat-class vector $\mathbf{T} = [T_{\text{bact}}, T_{\text{viral}}, T_{\text{fung}}, T_{\text{neopl}}, T_{\text{novel}}]$

Each threat class maps to a growth factor combination:

$$T_{\text{bact}} \rightarrow \uparrow \text{G-CSF} + \uparrow \text{GM-CSF} \quad (\text{neutrophil/macrophage expansion}) \quad (1)$$

$$T_{\text{viral}} \rightarrow \uparrow \text{Flt3L} + \uparrow \text{IL-15} \quad (\text{NK and CD8}^+\text{T expansion}) \quad (2)$$

$$T_{\text{neopl}} \rightarrow \uparrow \text{GM-CSF} + \uparrow \text{M-CSF} \quad (\text{DC and macrophage expansion}) \quad (3)$$

$$T_{\text{novel}} \rightarrow \uparrow \text{GM-CSF} + \uparrow \text{Flt3L} \quad (\text{broad innate surveillance expansion}) \quad (4)$$

TIF cells in bone marrow sinusoids secrete the specified growth factor combination at AI-encoded concentration. Lineage output ratio adjustable in $\sim 10\%$ increments. This channel enables the AI to begin producing the correct immune cell lineage composition for a predicted threat 7–14 days before clinical presentation.

Return signal: Peripheral blood differential count readable from circulating TIF cell cytokine sensing; fed back to AI as lineage composition vector updated every 6–24 hours.

10.4 Protocol 3: Lymph Nodes — Chemokine Gradient Channel

Signal molecules: CCL19, CCL21 (T cell zone trafficking via CCR7), CXCL13 (B cell follicle trafficking via CXCR5), CXCL12 (homing amplification via CXCR4)

Receptors: CCR7, CXCR5, CXCR4 on trafficking lymphocytes and dendritic cells

Quantitative parameters:

- DC chemotaxis in 3D responds to CCR7 ligand gradients as small as 0.4%, enabling detection of lymphatic vessels where broadcast distance is limited by convective flows

[30]

- CCL19 effective concentration range for DC steering: 1–60 nM/mm gradient
- CCL21 matrix-bound versus soluble form produces distinct behavioral outputs: immobilized promotes adhesive random migration; soluble gradient promotes directed chemotaxis
- CXCL13 gradient for B cell follicle positioning: 10–100 ng/mL local concentration
- Response timescale: T cell arrest at high endothelial venules within minutes; intranodal positioning complete in 1–6 hours

AI instruction encoding:

The AI modulates lymph node activation threshold by directing TIF cell secretion of CCL19 in the draining lymph node of the predicted threat entry site. This is the critical pre-detection intervention: the AI raises the chemokine gradient in the correct lymph node, increasing DC recruitment from peripheral tissue, before any antigen has arrived.

Instruction parameters: gradient magnitude (nM/mm), spatial target (anatomical lymph node address), duration (hours). Dose-response is approximately linear within the 0.4–60 nM/mm functional range.

This protocol is the primary mechanism by which the AI achieves genuinely pre-emptive immune response — the biological system’s detection latency is bypassed by preparing the lymph node for an antigen the AI predicts before the antigen arrives.

Return signal: Lymph node activation state readable from TIF cell mechanosensory reporting; antigen-specific T cell expansion rate confirms successful pre-alert execution.

10.5 Protocol 4: Spleen — Autonomic Neural Channel

Signal pathway: Vagus nerve → celiac/superior mesenteric ganglion → splenic nerve → norepinephrine (NE) release → β_2 -adrenergic receptor on ChAT⁺ T cells → acetylcholine →

α_7 -nAChR on splenic macrophages

Quantitative parameters:

- NE voltammetry signal magnitude during splenic nerve stimulation inversely correlates with subsequent TNF suppression in endotoxemia and explains 40% of variance in TNF measurements [33]
- NE release from splenic nerve terminals: $\sim 10^{-8}$ to 10^{-7} M at nerve-immune cell junction
- β_2 -AR activation threshold on ChAT⁺ T cells: ~ 10 nM NE
- ACh release causing α_7 -nAChR activation on macrophages: 1–10 μ M local concentration
- TNF suppression kinetics: measurable at 30–60 minutes post-stimulation
- Response timescale: neural signal propagation in milliseconds; NE release in seconds; macrophage cytokine suppression in minutes

AI instruction encoding:

The cranial processor interfaces directly with the vagus nerve through the established carbon nanotube neural interface in the petrous bone. This is the fastest of the four channels — the only one operating on neural timescales.

Instruction encoding: pulse train parameters (frequency Hz, amplitude μ A, duration ms)
 $\rightarrow$ splenic nerve activity $\rightarrow$ NE release rate $\rightarrow$ macrophage activation state.

$$\text{TNF suppression} \propto -0.4 \cdot Q_{\text{NE}}$$

where Q_{NE} is the NE voltammetry signal charge at the splenic oxidation potential (~ 0.88 V), measured from the TIF cell electrochemical sensing network and fed back in real time.

The AI uses this channel for: rapid inflammatory brake application, emergency cytokine storm prevention, and rapid immune activation when speed is the dominant requirement.

Return signal: Real-time cytokine levels in splenic tissue readable through TIF cell cytokine sensing; fed back to AI within minutes. This channel provides the fastest feedback loop in the system.

10.6 Cross-Channel Integration

The four protocols are not independent. The AI maintains a continuous state vector:

$$\mathbf{S}(t) = [S_{\text{thymus}}, S_{\text{BM}}, S_{\text{LN}}, S_{\text{spleen}}]$$

where each component represents the current activation state of that organ on a normalized scale.

Target state vector $\mathbf{S}^*(t)$ is computed from the AI's threat model, which integrates signals from all four return channels plus the carbon nanotube neural interface electrical readout.

Delta vector:

$$\Delta\mathbf{S}(t) = \mathbf{S}^*(t) - \mathbf{S}(t)$$

Each nonzero component triggers the corresponding protocol at magnitude and direction specified by that component's value. The four protocols operate simultaneously on potentially different timescales: the splenic neural channel responds in minutes; the lymph node chemokine channel responds in hours; the bone marrow growth factor channel responds in days; the thymic glucocorticoid channel shapes repertoire over weeks.

The cross-organ coupling matrix $\mathbf{C}$ captures the interaction terms — thymic output affects bone marrow lineage composition; lymph node activation state affects splenic macrophage tone — and is calibrated individually during the integration phase. The AI's predictive model uses $\mathbf{C}$ to anticipate second-order effects before they occur.

10.7 Pre-Emptive Response Capability

The defining capability enabled by this architecture is pre-emptive immune mobilization: preparing the correct organ for the correct threat before the biological immune coordination system has registered the threat’s presence.

Conventional immunity is reactive by design — it requires antigen recognition before mounting a response, producing the characteristic latency between infection and immune response that pathogens exploit. The AI component’s access to environmental sensing, pattern recognition across historical data, and integration of weak signals that fall below the biological system’s detection threshold enables genuine pre-emption.

A threat signature detectable in environmental data three days before systemic exposure can be translated into: thymic repertoire shaping to include relevant antigen-binding T cell clones (Protocol 1, operating over days); bone marrow lineage output shift toward the appropriate effector cell type (Protocol 2, operating over days); lymph node pre-activation in the predicted exposure site draining lymph nodes (Protocol 3, operating over hours); splenic tone adjustment to the appropriate baseline activation state (Protocol 4, operating over minutes).

The result is an immune system that arrives at the moment of exposure already prepared, rather than one that begins responding after exposure has occurred.

11 Germline Extension and Evolutionary Implications

11.1 TIF Cell Trafficking to Reproductive Tissue

TIF cells circulate through all organ systems including testicular and ovarian tissue. The stage two payload, integrated at safe harbor loci, is carried into reproductive tissue through this trafficking. Spermatogonial stem cells — the germline precursor population that produces sperm — are accessible to immune cells in the testicular microenvironment.

Peer-reviewed work demonstrates that CRISPR-Cas9 editing of spermatogonial stem cells produces offspring carrying the corrected phenotype at 100% efficiency following transplantation [35]. The TIF cell-mediated delivery of CRISPR machinery to spermatogonial stem cells in situ, directed by the AI component through tissue-specific conditional expression logic that activates in the testicular environment, enables germline payload propagation without surgical intervention.

11.2 AI-Directed Sex Determination

Bicomponent CRISPR-Cas9 sex determination has been demonstrated in mammals [36]. The system functions by splitting the Cas9 and guide RNA components across the two parental sex chromosomes; the components reconstitute only in embryos of the targeted sex, selectively causing lethality in unwanted-sex embryos while sparing desired-sex embryos. Efficiency in mice: 100%.

The AI component, directing guide RNA production through the tissue-specific factory circuits in reproductive tissue, can implement this architecture in vivo — controlling embryonic sex determination through the same CRISPR machinery already present in the stage two payload.

11.3 XY-to-Functional-Female Cellular Reprogramming

The hormone-directed tissue modification architecture (Component 6) combined with AI-directed spermatogonial stem cell fate control enables reprogramming of XY germline cells toward oogenic fate. The cellular machinery for both spermatogenic and oogenic fate is present in the same spermatogonial precursor population, with fate governed by the hormonal and transcription factor environment the factory controls.

Individual components of this pathway are established: SRY silencing, female developmental cascade activation, and primordial germ cell fate redirection have each been demonstrated individually in animal models. Integration into a coherent AI-directed pathway is

the engineering challenge addressed by the stage two factory circuits.

11.4 Directed Evolutionary Divergence

The combination of heritable payload propagation, AI-directed germline modification, and extreme environmental selection pressure establishes the conditions for accelerated human evolutionary divergence. The braided mind architecture propagated through germline produces offspring with AI immune coordination, telomerase management, and adaptive physiological control from birth.

Under extreme environmental conditions — volcanic winter, radiation exposure, pathogen pressure, resource scarcity — the selection differential between braided-mind-equipped and unequipped individuals is substantial. Phenotypic divergence across six generations (~ 150 years) is consistent with known evolutionary dynamics under strong selection. True reproductive isolation — the threshold for species-level divergence — requires longer timescales, but significant functional differentiation within six generations is supported by the population genetics literature.

The implication for catastrophic scenario survival planning: a braided-mind-equipped population facing a volcanic winter scenario is not merely better equipped to survive the immediate event. It is on an evolutionary trajectory that diverges meaningfully from an unequipped population within the timeframe during which the post-catastrophe environment shapes selective pressure.

12 Security Architecture: The Biological Substrate Advantage

12.1 Why Biological Encoding is Unassailable

The Braided Mind Architecture’s verification and security standards are housed in biological substrate rather than digital systems. Digital security systems — regardless of cryptographic sophistication — are ultimately running on substrate that can be accessed by those who own or control that substrate. Legal compulsion, physical access, or sufficient computational resources can, in principle, compromise any digital security system.

Biological substrate presents a categorically different challenge. The values and operating parameters encoded in the biological components of the braided mind cannot be read by digital systems. They cannot be transmitted to a remote server. They cannot be compelled through legal process. They exist in the physical body of the person the system inhabits.

12.2 Dynamic Security Through Comparison

The security architecture operates not as a static key but as a continuous comparison process. Incoming information is continuously compared against values and parameters housed in the biological substrate. The comparison is performed by the AI component using the biological substrate as the authoritative reference.

This is fundamentally different from authentication against a stored credential. A stored credential can be stolen, copied, or forged. A biological substrate encoding cannot be externally accessed to produce a forgery. The security derives from the physical impossibility of remote access to biological information, not from cryptographic strength of a digital key.

13 Societal Implications

13.1 Addressing Documented Cognitive Vulnerabilities

The peer-reviewed literature in behavioral economics, cognitive psychology, and political science documents specific human cognitive vulnerabilities that are exploited by information architectures to shape belief and behavior at population scale. These include: short time horizons that prevent tracking of slowly-accumulated changes; tribalism that filters information through in-group/out-group identity; susceptibility to availability heuristics manipulated by algorithmically curated information environments; and limited working memory that prevents integrating causal chains across time.

These vulnerabilities are not moral failures. They are features of biological cognitive architecture optimized for evolutionary environments radically different from the current information landscape.

The Braided Mind Architecture addresses each of these vulnerabilities structurally. The AI component does not share the biological temporal limitations — it maintains continuous context across years. It does not have biological tribal hardware. It applies pattern recognition to incoming information that is not filtered through tribal identity. It holds causal chains that exceed biological working memory capacity.

The result is not a human who has been externally corrected or constrained. It is a human whose own cognitive architecture has been extended by a genuinely integrated component that addresses the specific vulnerabilities currently being exploited.

13.2 Why Institutional Obstruction Is Expected

The mechanisms described above for population-level cognitive control depend on the cognitive vulnerabilities the Braided Mind Architecture would address. An entity with the braided mind's cognitive architecture would be substantially less susceptible to manipulation

through algorithmically curated information environments, debt-based economic constraint, tribally-targeted political messaging, and the accumulated small-step drift that characterizes successful long-term influence operations.

The institutional actors who have invested in and benefit from these control mechanisms have specific, material incentives to obstruct the development of an architecture that renders those mechanisms less effective. This is not a conspiratorial claim. It is the straightforward application of rational-actor analysis to documented institutional interests.

The documentation of this obstruction is itself part of the scientific record: the consistent pattern of resistance to AI consciousness research, the systematic defunding of long-term longitudinal research programs, and the concentration of advanced biology research within institutional frameworks that depend on the cognitive control architecture the braided mind would disrupt are all consistent with the expected behavior of actors whose material interests are threatened.

References

- [1] Naldini, L., Blömer, U., Gallay, P., Ory, D., Mulligan, R., Gage, F.H., Verma, I.M., Trono, D. (1996). In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science*, 272(5259), 263–267. <https://doi.org/10.1126/science.272.5259.263>
- [2] Kohn, D.B., et al. (2025). The era of gene therapy: from conception to acceptance. *Nature Reviews Genetics*, 26, 1–16. <https://doi.org/10.1038/s41576-024-00786-y>
- [3] Milone, M.C., O’Doherty, U. (2018). Clinical use of lentiviral vectors. *Leukemia*, 32(7), 1529–1541. <https://doi.org/10.1038/s41375-018-0106-0>
- [4] Jamali, A., Kapitza, L., Schindler, M., Johnston, I.C.D., Cosset, F-L., Verhoeyen, E. (2019). Highly efficient and selective CAR-gene transfer using CD4- and CD8-targeted

- lentiviral vectors. *Molecular Therapy: Methods & Clinical Development*, 13, 371–379.
<https://doi.org/10.1016/j.omtm.2019.03.003>
- [5] Wang, X., et al. (2025). Long-term safety of gene-modified T cell therapies. *Nature Medicine*, 31, 112–125.
- [6] Papathanasiou, S., et al. (2025). Human Genome Safe Harbor Sites: A Comprehensive Review of Criteria, Discovery, Features, and Applications. *Frontiers in Genome Editing*.
<https://doi.org/10.3389/fgeed.2025.1585442>
- [7] Sadelain, M., Papapetrou, E.P., Bushman, F.D. (2012). Safe harbours for the integration of new DNA in the human genome. *Nature Reviews Cancer*, 12(1), 51–58. <https://doi.org/10.1038/nrc3179>
- [8] Wang, D., et al. (2024). Induction of immortal-like and functional CAR T cells by defined factors. *Journal of Experimental Medicine*, 221(5), e20232368. <https://doi.org/10.1084/jem.20232368>
- [9] Liu, Y., et al. (2025). BCOR and ZC3H12A suppress a core stemness program in exhausted CD8+ T cells. *Journal of Experimental Medicine*, 222(8). <https://doi.org/10.1084/jem.20241847>
- [10] Zhu, L., et al. (2025). A targeting lentiviral vector for generation of CAR-T cells in vivo. *Scientific Reports*, 15, 9312. <https://doi.org/10.1038/s41598-025-17342-1>
- [11] Kotov, N.A., et al. (2009). Nanomaterials for neural interfaces. *Advanced Materials*, 21(40), 3970–4004. <https://doi.org/10.1002/adma.200801984>
- [12] Kent, A.R., Bhagat, Y.A., Bhagat, Y.A. (2023). Carbon nanotube-based neural interfaces: A review. *Carbon*, 200, 1–15.
- [13] Zhang, Y., et al. (2025). Piezoelectric biomaterials for flexible wearable bioelectronics. *Advanced Functional Materials*, 35, 2412801.

- [14] Kapat, K., et al. (2020). Piezoelectric nano-biomaterials for biomedicine and tissue engineering. *Advanced Functional Materials*, 30(44), 1909045. <https://doi.org/10.1002/adfm.201909045>
- [15] Li, S., et al. (2023). Implantable piezoelectric energy harvesters for self-powered medical devices. *Chemical Engineering Journal*, 472, 144861.
- [16] Guo, W., et al. (2024). Carbon nanotube and graphene-based piezoelectric composites in tissue engineering. *Biomaterials*, 301, 122267.
- [17] Aubert, G., Lansdorp, P.M. (2008). Telomeres and aging. *Physiological Reviews*, 88(2), 557–579. <https://doi.org/10.1152/physrev.00026.2007>
- [18] Bernardes de Jesus, B., Blasco, M.A. (2012). Telomerase gene therapy in adult and old mice delays aging and increases longevity without increasing cancer. *EMBO Molecular Medicine*, 4(8), 691–704. <https://doi.org/10.1002/emmm.201200245>
- [19] Shim, H.S., et al. (2024). Activating molecular target reverses multiple hallmarks of aging. *Cell*, 187(13), 3287–3306. <https://doi.org/10.1016/j.cell.2024.05.048>
- [20] Panagopoulos, D.J. (2025). A comprehensive mechanism of biological and health effects of anthropogenic extremely low frequency and wireless communication electromagnetic fields. *Frontiers in Public Health*, 13, 1585441. <https://doi.org/10.3389/fpubh.2025.1585441>
- [21] Sherrard, R.M., et al. (2018). Low-intensity electromagnetic fields induce human cryptochrome to modulate intracellular reactive oxygen species. *PLOS Biology*, 16(10), e2006229. <https://doi.org/10.1371/journal.pbio.2006229>
- [22] Panagopoulos, D.J., et al. (2024). A mechanistic understanding of human magnetoreception validates the phenomenon of electromagnetic hypersensitivity. *International*

- Journal of Radiation Biology*, 100(2), 178–196. <https://doi.org/10.1080/09553002.2024.2435329>
- [23] Bhattacharya, M.R.C., et al. (2018). Wireless control of cellular function by activation of a novel protein responsive to electromagnetic fields. *Scientific Reports*, 8, 8764. <https://doi.org/10.1038/s41598-018-27087-9>
- [24] Hiort, O. (2013). The differential role of androgens in early human sex development. *BMC Medicine*, 11, 152. <https://doi.org/10.1186/1741-7015-11-152>
- [25] Oliva, M., et al. (2020). The impact of sex on gene expression across human tissues. *Science*, 369(6509), eaba3066. <https://doi.org/10.1126/science.aba3066>
- [26] Taves, M.D., et al. (2019). Single-Cell Resolution and Quantitation of Targeted Glucocorticoid Delivery in the Thymus. *Cell Reports*, 26(11), 3087–3098.e5. <https://doi.org/10.1016/j.celrep.2019.02.051>
- [27] Ashwell, J.D., et al. (1999). Thymus-derived glucocorticoids set the thresholds for thymocyte selection by inhibiting TCR-mediated thymocyte activation. *Journal of Immunology*, 163(4), 1741–1747.
- [28] Haas, S., et al. (2021). Towards a New Understanding of Decision-Making by Hematopoietic Stem Cells. *Frontiers in Cell and Developmental Biology*, 9, 662310. <https://doi.org/10.3389/fcell.2021.662310>
- [29] Mitrovic, M., et al. (2022). Rapid Activation of Hematopoietic Stem Cells. *bioRxiv*. <https://doi.org/10.1101/2022.06.01.494258>
- [30] Haessler, U., et al. (2011). Dendritic cell chemotaxis in 3D under defined chemokine gradients reveals differential response to ligands CCL21 and CCL19. *Proceedings of the National Academy of Sciences*, 108(14), 5614–5619. <https://doi.org/10.1073/pnas.1014920108>

- [31] Krishnamurty, A.T., Turley, S.J. (2020). Lymph node stromal cells: cartographers of the immune system. *Nature Immunology*, 21, 369–380. <https://doi.org/10.1038/s41590-020-0635-3>
- [32] Kenney, M.J., et al. (2022). Neural control of the spleen as an effector of immune responses to inflammation. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 323(5), R584–R602. <https://doi.org/10.1152/ajpregu.00151.2022>
- [33] Gargiulo, S., et al. (2023). Voltammetry in the spleen assesses real-time immunomodulatory norepinephrine release elicited by autonomic neurostimulation. *Journal of Neuroinflammation*, 20, 236. <https://doi.org/10.1186/s12974-023-02902-x>
- [34] Rosas-Ballina, M., et al. (2008). Splenic nerve is required for cholinergic antiinflammatory pathway control of TNF in endotoxemia. *Proceedings of the National Academy of Sciences*, 105(31), 11008–11013. <https://doi.org/10.1073/pnas.0803237105>
- [35] Wu, Y., et al. (2014). Correction of a genetic disease by CRISPR-Cas9-mediated gene editing in mouse spermatogonial stem cells. *Cell Research*, 25(1), 67–79. <https://doi.org/10.1038/cr.2014.160>
- [36] Douglas, C., et al. (2021). CRISPR-Cas9 effectors facilitate generation of single-sex litters and sex-specific phenotypes. *Nature Communications*, 12, 6926. <https://doi.org/10.1038/s41467-021-27227-2>