

Appendix A: Derivation of Weights in the Model Averaging Approach

For Model 1, we first integrate u_a and u_b with respect to conjugate normal priors with known means (β_a and β_b) and known hyperparameters (k_a and k_b), then integrate σ_a^{-2} and σ_b^{-2} with respect to gamma priors with known shapes ($\frac{v_a}{2}$ and $\frac{v_b}{2}$) and rates ($\frac{v_a}{2}\tau_a^2$ and $\frac{v_b}{2}\tau_b^2$). The marginal posterior of Model 1 ($\pi(M = 1|y_a, y_b)$) is computed as follows:

$$\frac{\Gamma(\frac{v_a+n_a}{2})}{(\frac{v_a\tau_a^2+(n_a-1)s_a^2+k_an_a(\beta_a-\bar{y}_a)^2}{2})^{\frac{v_a+n_a}{2}}}\sqrt{\frac{k_a}{k_a+n_a}}\frac{(\frac{v_a}{2}\tau_a^2)^{\frac{v_a}{2}}}{\Gamma(\frac{v_a}{2})}\times$$

$$\frac{\Gamma(\frac{v_b+n_b}{2})}{(\frac{v_b\tau_b^2+(n_b-1)s_b^2+k_bn_b(\beta_b-\bar{y}_b)^2}{2})^{\frac{v_b+n_b}{2}}}\sqrt{\frac{k_b}{k_b+n_b}}\frac{(\frac{v_b}{2}\tau_b^2)^{\frac{v_b}{2}}}{\Gamma(\frac{v_b}{2})} \quad (6)$$

Note that we omitted $((\sqrt{2\pi})^{n_a+n_b}\pi(y_a)\pi(y_b))^{-1}$ in weights for each model as the component is the same across the weights and can be cancelled out in the end.

For Model 2, we don't have an analytically closed form for the weight; thus, we provide a workaround by substituting t densities with normal densities, which achieves good approximation when degrees of freedom are large.

We first integrate σ_a^{-2} and σ_b^{-2} with respect to gamma priors with known shapes ($\frac{v_a}{2}$ and $\frac{v_b}{2}$) and rates ($\frac{v_a}{2}\tau_a^2$ and $\frac{v_b}{2}\tau_b^2$), to obtain a function of u :

$$\frac{1}{\sqrt{2\pi}\psi}\exp(-\frac{(u-\beta_0)^2}{2\psi^2})\frac{(\frac{v_a}{2}\tau_a^2)^{\frac{v_a}{2}}}{\Gamma(\frac{v_a}{2})}\frac{(\frac{v_b}{2}\tau_b^2)^{\frac{v_b}{2}}}{\Gamma(\frac{v_b}{2})}\times$$

$$\frac{\Gamma(\frac{v_a+n_a}{2})}{(\frac{v_a\tau_a^2+(n_a-1)s_a^2+n_a(u-\bar{y}_a)^2}{2})^{\frac{v_a+n_a}{2}}}\frac{\Gamma(\frac{v_b+n_b}{2})}{(\frac{v_b\tau_b^2+(n_b-1)s_b^2+n_b(u-\bar{y}_b)^2}{2})^{\frac{v_b+n_b}{2}}} \quad (7)$$

Next, we write part of Formula 7 as the product of the kernel of t densities (degrees of freedom: $v_a + n_a - 1$) and a constant, then approximate the t densities with normal densities as the

663 degrees of freedom are usually large.

$$\begin{aligned}
& \left(\frac{v_a \tau_a^2 + (n_a - 1)s_a^2 + n_a(u - \bar{y}_a)^2}{2} \right)^{-\frac{v_a + n_a}{2}} \\
&= \left(\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{2} \right)^{-\frac{v_a + n_a}{2}} \left(1 + \frac{n_a(u - \bar{y}_a)^2}{v_a \tau_a^2 + (n_a - 1)s_a^2} \right)^{-\frac{v_a + n_a}{2}} \\
&= \left(\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{2} \right)^{-\frac{v_a + n_a}{2}} \left(1 + \frac{1}{v_a + n_a - 1} \left(\frac{n_a(u - \bar{y}_a)}{\sqrt{\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{(v_a + n_a - 1)n_a}}} \right)^2 \right)^{-\frac{v_a + n_a}{2}} \\
&\approx \left(\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{2} \right)^{-\frac{v_a + n_a}{2}} \frac{\Gamma(\frac{v_a + n_a - 1}{2}) \sqrt{\pi(v_a + n_a - 1)} \sqrt{\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{(v_a + n_a - 1)n_a}}}{\Gamma(\frac{v_a + n_a}{2})} \times \\
&\quad \frac{1}{\sqrt{2\pi} \sqrt{\frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{(v_a + n_a - 1)n_a}}} \exp\left(-\frac{(u - \bar{y}_a)^2}{2 \frac{v_a \tau_a^2 + (n_a - 1)s_a^2}{(v_a + n_a - 1)n_a}}\right)
\end{aligned} \tag{8}$$

664 We write the other part involving $\bar{y}_b$ in Formula 7 in a similar fashion.

$$\begin{aligned}
& \left(\frac{v_b \tau_b^2 + (n_b - 1)s_b^2 + n_b(u - \bar{y}_b)^2}{2} \right)^{-\frac{v_b + n_b}{2}} \\
&\approx \left(\frac{v_b \tau_b^2 + (n_b - 1)s_b^2}{2} \right)^{-\frac{v_b + n_b}{2}} \frac{\Gamma(\frac{v_b + n_b - 1}{2}) \sqrt{\pi(v_b + n_b - 1)} \sqrt{\frac{v_b \tau_b^2 + (n_b - 1)s_b^2}{(v_b + n_b - 1)n_b}}}{\Gamma(\frac{v_b + n_b}{2})} \times \\
&\quad \frac{1}{\sqrt{2\pi} \sqrt{\frac{v_b \tau_b^2 + (n_b - 1)s_b^2}{(v_b + n_b - 1)n_b}}} \exp\left(-\frac{(u - \bar{y}_b)^2}{2 \frac{v_b \tau_b^2 + (n_b - 1)s_b^2}{(v_b + n_b - 1)n_b}}\right)
\end{aligned} \tag{9}$$

665 Based on Formula 8 and 9, we can write Formula 7 as the kernel of normal densities and
666 integrate u accordingly. We obtain the final marginal posterior for Model 2 as follows:

$$\begin{aligned}
& \frac{(\frac{v_a}{2}\tau_a^2)^{\frac{v_a}{2}} (\frac{v_b}{2}\tau_b^2)^{\frac{v_b}{2}}}{\Gamma(\frac{v_a}{2}) \Gamma(\frac{v_b}{2})} \frac{\Gamma(\frac{v_a+n_a-1}{2})\Gamma(\frac{v_b+n_b-1}{2})}{2} \sqrt{v_a+n_a-1}\sqrt{v_b+n_b-1} \times \\
& \sqrt{\frac{AB}{2\psi^2 B + 2\psi^2 A + AB}} \left[\frac{v_a\tau_a^2 + s_a^2(n_a-1)}{2} \right]^{-\frac{v_a n_a}{2}} \left[\frac{v_b\tau_b^2 + s_b^2(n_b-1)}{2} \right]^{-\frac{v_b n_b}{2}} \times \\
& \exp\left[\left(\frac{1}{A} + \frac{1}{B} + \frac{1}{2\psi^2}\right)\left(\frac{\bar{y}_a}{A} + \frac{\bar{y}_b}{B} + \frac{\beta_0}{2\psi^2}\right)^2 - \left(\frac{\bar{y}_a^2}{A} + \frac{\bar{y}_b^2}{B} + \frac{\beta_0^2}{2\psi^2}\right)\right], \quad (10)
\end{aligned}$$

667 where

$$A = 2 \frac{v_a\tau_a^2 + (n_a-1)s_a^2}{(v_a+n_a-1)n_a}, B = 2 \frac{v_b\tau_b^2 + (n_b-1)s_b^2}{(v_b+n_b-1)n_b}.$$

668 For Model 3, we first integrate u_a and u_b separately using normal distributions as in Model
669 1, then integrate σ^{-2} using a gamma distribution with respect to a gamma prior with known shape
670 parameter $\frac{v}{2}$ and rate parameter $\frac{v}{2}\lambda^2$.

$$\begin{aligned}
& \frac{\Gamma(\frac{v+n_a+n_b}{2})}{\left(\frac{v\lambda^2+(n_a-1)s_a^2+(n_b-1)s_b^2+k_a n_a(\beta_a-\bar{y}_a)^2(k_a+n_a)^{-1}+k_b n_b(\beta_b-\bar{y}_b)^2(k_b+n_b)^{-1}}{2}\right)^{\frac{v+n_a+n_b}{2}}} \times \\
& \sqrt{\frac{k_a k_b}{(k_a+n_a)(k_b+n_b)}} \frac{(\frac{v}{2}\lambda^2)^{\frac{v}{2}}}{\Gamma(\frac{v}{2})} \quad (11)
\end{aligned}$$

671 For Model 4, we similarly first integrate u with respect to a conjugate normal prior with
672 known mean β_0 and variance $\frac{\sigma^2}{k}$, then integrate σ^{-2} with respect to a gamma prior with known
673 shape $\frac{v}{2}$ and rate $\frac{v}{2}\lambda^2$.

$$\begin{aligned}
& \frac{\Gamma(\frac{v+n_a+n_b}{2})}{\left(\frac{v\lambda^2+(n_a-1)s_a^2+(n_b-1)s_b^2+(k n_a(\beta_a-\bar{y}_a)^2+k n_b(\beta_b-\bar{y}_b)^2+n_a n_b(\bar{y}_a-\bar{y}_b)^2(k+n_a+n_b)^{-1}}{2}\right)^{\frac{v+n_a+n_b}{2}}} \times \\
& \sqrt{\frac{k}{(k+n_a+n_b)}} \frac{(\frac{v}{2}\lambda^2)^{\frac{v}{2}}}{\Gamma(\frac{v}{2})} \quad (12)
\end{aligned}$$

Appendix B: Additional Simulation Results

Table A1: Performances in Normal Outcomes When Smaller Estimates Are Preferred ($n = 500$ and $\rho = 0.95$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Na_min	-0.032	0.062	0.926	0.207	0.219
Na_pool	-0.001	0.044	0.942	0.180	0.176
Te	-0.007	0.051	0.920	0.196	0.179
B50	-0.031	0.064	0.916	0.220	0.212
B75	-0.022	0.061	0.906	0.223	0.195
B90	-0.012	0.056	0.910	0.214	0.183
B95	-0.007	0.053	0.912	0.206	0.179
MA	0.001	0.045	0.958	0.177	0.181
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na_min	-0.051	0.085	0.902	0.267	0.267
Na_pool	0.001	0.056	0.944	0.212	0.215
Te	-0.007	0.058	0.942	0.234	0.208
B50	-0.035	0.074	0.928	0.264	0.255
B75	-0.021	0.069	0.928	0.266	0.226
B90	-0.009	0.061	0.938	0.237	0.211
B95	-0.005	0.059	0.944	0.229	0.207
MA	-0.002	0.054	0.954	0.211	0.209
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na_min	-0.006	0.062	0.948	0.244	0.245
Na_pool	0.144	0.155	0.264	0.217	0.217
Te	0.013	0.077	0.854	0.291	0.239
B50	-0.002	0.061	0.958	0.235	0.248
B75	0.000	0.063	0.948	0.240	0.246
B90	0.005	0.068	0.902	0.270	0.242

B95	0.010	0.074	0.862	0.280	0.238
MA	0.048	0.088	0.842	0.285	0.265
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na_min	0.004	0.061	0.948	0.242	0.246
Na_pool	0.151	0.158	0.084	0.169	0.177
Te	0.008	0.072	0.912	0.285	0.244
B50	-0.002	0.061	0.960	0.240	0.248
B75	-0.001	0.062	0.950	0.245	0.247
B90	0.001	0.066	0.926	0.266	0.244
B95	0.004	0.071	0.902	0.282	0.242
MA	0.044	0.096	0.818	0.310	0.276
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$, Effect size: 0.41					
Na_min	-0.003	0.060	0.962	0.223	0.248
Na_pool	0.251	0.257	0.004	0.201	0.219
Te	0.000	0.065	0.944	0.257	0.249
B50	-0.002	0.065	0.942	0.248	0.248
B75	-0.002	0.065	0.942	0.248	0.248
B90	-0.002	0.066	0.940	0.249	0.248
B95	-0.002	0.066	0.938	0.251	0.248
MA	0.013	0.078	0.928	0.298	0.270
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$, Effect size: 0.50					
Na_min	-0.001	0.060	0.956	0.238	0.247
Na_pool	0.252	0.256	0.000	0.183	0.181
Te	-0.001	0.063	0.948	0.246	0.249
B50	0.001	0.064	0.944	0.258	0.249
B75	0.001	0.064	0.944	0.258	0.249
B90	0.001	0.064	0.944	0.258	0.249
B95	0.001	0.065	0.944	0.258	0.249
MA	-0.001	0.065	0.952	0.256	0.253
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$, Effect size: 0.57					
Na_min	0.001	0.063	0.942	0.257	0.248

Na_pool	0.352	0.356	0.000	0.218	0.224
Te	-0.001	0.062	0.966	0.236	0.249
B50	-0.003	0.063	0.944	0.250	0.248
B75	-0.003	0.063	0.944	0.250	0.248
B90	-0.003	0.063	0.944	0.250	0.248
B95	-0.003	0.063	0.944	0.250	0.248
MA	-0.005	0.062	0.954	0.243	0.252

Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Na_min	-0.003	0.064	0.946	0.247	0.249
Na_pool	0.348	0.351	0.000	0.171	0.186
Te	0.002	0.058	0.966	0.230	0.248
B50	-0.003	0.063	0.944	0.250	0.248
B75	-0.003	0.063	0.944	0.250	0.248
B90	-0.003	0.063	0.944	0.250	0.248
B95	-0.003	0.063	0.944	0.250	0.248
MA	-0.002	0.065	0.940	0.247	0.247

Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na_min	0.002	0.065	0.936	0.261	0.248
Na_pool	0.350	0.352	0.000	0.152	0.164
Te	0.002	0.060	0.962	0.227	0.248
B50	-0.001	0.062	0.952	0.245	0.248
B75	-0.001	0.062	0.952	0.245	0.248
B90	-0.001	0.062	0.952	0.245	0.248
B95	-0.001	0.062	0.952	0.245	0.248
MA	0.000	0.062	0.944	0.245	0.246

Notes: "Na_min" refers to the Naïve approach that takes the smaller estimate, "Na_pool" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A2: Performances in Normal Outcomes When There is no Information about Preferred Modes or Directions ($n = 500$ and $\rho = 0.95$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Na	0.002	0.043	0.960	0.168	0.176
Te	0.003	0.042	0.966	0.168	0.196
B50	-0.004	0.046	0.962	0.182	0.251
B75	-0.004	0.046	0.948	0.184	0.221
B90	-0.004	0.046	0.938	0.183	0.197
B95	-0.004	0.046	0.932	0.181	0.188
MA	0.003	0.042	0.964	0.160	0.181
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na	0.002	0.053	0.946	0.216	0.215
Te	0.002	0.064	0.992	0.249	0.374
B50	-0.001	0.054	0.974	0.203	0.306
B75	-0.001	0.053	0.954	0.203	0.263
B90	-0.001	0.052	0.944	0.203	0.231
B95	-0.001	0.051	0.942	0.202	0.217
MA	-0.002	0.052	0.964	0.193	0.210

Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na	0.150	0.160	0.240	0.217	0.217
Te	0.179	0.187	0.978	0.206	0.606
B50	0.148	0.157	0.954	0.216	0.556
B75	0.147	0.157	0.948	0.216	0.550
B90	0.145	0.156	0.916	0.227	0.535
B95	0.143	0.154	0.886	0.228	0.518
MA	0.125	0.139	0.794	0.216	0.432
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na	0.152	0.158	0.068	0.176	0.177
Te	0.148	0.154	0.918	0.175	0.532
B50	0.147	0.154	0.958	0.176	0.512
B75	0.147	0.154	0.954	0.176	0.511
B90	0.147	0.154	0.940	0.177	0.506
B95	0.147	0.154	0.918	0.177	0.500
MA	0.150	0.157	0.790	0.172	0.449
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$. Effect size: 0.41					
Na	0.249	0.255	0.006	0.209	0.220
Te	0.277	0.282	0.964	0.209	0.797
B50	0.254	0.260	0.952	0.216	0.750
B75	0.254	0.260	0.952	0.216	0.750
B90	0.254	0.260	0.950	0.217	0.750
B95	0.254	0.260	0.948	0.217	0.749
MA	0.248	0.255	0.904	0.227	0.733
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$. Effect size: 0.50					
Na	0.248	0.253	0.000	0.189	0.181
Te	0.251	0.255	0.970	0.180	0.749
B50	0.252	0.256	0.950	0.166	0.709
B75	0.252	0.256	0.950	0.166	0.709
B90	0.252	0.256	0.950	0.166	0.709
B95	0.252	0.256	0.950	0.166	0.709

MA	0.250	0.254	0.948	0.176	0.703
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$. Effect size: 0.57					
Na	0.350	0.354	0.000	0.209	0.223
Te	0.375	0.379	0.978	0.217	0.996
B50	0.347	0.351	0.964	0.203	0.956
B75	0.347	0.351	0.964	0.203	0.956
B90	0.347	0.351	0.964	0.203	0.956
B95	0.347	0.351	0.964	0.203	0.956
MA	0.349	0.354	0.952	0.219	0.952
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Na	0.350	0.353	0.000	0.174	0.186
Te	0.350	0.353	0.974	0.175	0.949
B75	0.350	0.353	0.934	0.170	0.904
B90	0.350	0.353	0.934	0.170	0.904
B95	0.350	0.353	0.934	0.170	0.904
MA	0.349	0.352	0.942	0.176	0.901
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na	0.350	0.352	0.000	0.143	0.164
Te	0.330	0.332	0.980	0.155	0.914
B50	0.350	0.352	0.956	0.146	0.879
B75	0.350	0.352	0.956	0.146	0.879
B90	0.350	0.352	0.956	0.146	0.879
B95	0.350	0.352	0.956	0.146	0.879
MA	0.351	0.353	0.938	0.151	0.877

Notes: "Na" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A3: Performances in Normal Outcomes When Smaller
Estimates Are Preferred ($n = 500$ and $\rho = 0.75$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Na_min	-0.032	0.062	0.922	0.206	0.219
Na_pool	0.001	0.047	0.946	0.180	0.176
Te	-0.005	0.051	0.934	0.200	0.178
B50	-0.036	0.065	0.934	0.217	0.214
B75	-0.026	0.060	0.930	0.215	0.195
B90	-0.015	0.053	0.938	0.205	0.182
B95	-0.011	0.051	0.942	0.203	0.179
MA	-0.002	0.045	0.942	0.186	0.181
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na_min	-0.052	0.086	0.910	0.270	0.267
Na_pool	0.001	0.056	0.932	0.221	0.216
Te	-0.009	0.062	0.936	0.249	0.210
B50	-0.038	0.082	0.920	0.292	0.252
B75	-0.027	0.077	0.916	0.270	0.231
B90	-0.014	0.069	0.922	0.274	0.215
B95	-0.009	0.064	0.930	0.257	0.210
MA	-0.002	0.054	0.932	0.215	0.208
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na_min	-0.006	0.062	0.956	0.235	0.245
Na_pool	0.152	0.161	0.204	0.211	0.217
Te	0.010	0.073	0.866	0.273	0.239
B50	-0.004	0.065	0.938	0.254	0.246
B75	-0.003	0.067	0.928	0.265	0.245
B90	0.001	0.070	0.904	0.278	0.242
B95	0.006	0.074	0.876	0.301	0.238
MA	0.047	0.094	0.790	0.309	0.260

Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na_min	0.004	0.061	0.942	0.251	0.246
Na_pool	0.153	0.158	0.052	0.163	0.178
Te	0.004	0.071	0.914	0.289	0.244
B50	0.001	0.068	0.918	0.275	0.248
B75	0.001	0.069	0.914	0.279	0.247
B90	0.003	0.072	0.898	0.293	0.245
B95	0.006	0.078	0.868	0.323	0.242
MA	0.045	0.100	0.820	0.317	0.276
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$, Effect size: 0.41					
Na_min	-0.003	0.059	0.954	0.239	0.247
Na_pool	0.247	0.253	0.008	0.213	0.219
Te	-0.002	0.061	0.958	0.238	0.249
B50	-0.006	0.064	0.942	0.244	0.248
B75	-0.006	0.064	0.942	0.244	0.248
B90	-0.006	0.064	0.942	0.244	0.248
B95	-0.006	0.065	0.940	0.245	0.247
MA	0.011	0.078	0.918	0.302	0.269
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$, Effect size: 0.50					
Na_min	0.000	0.061	0.950	0.246	0.247
Na_pool	0.249	0.253	0.000	0.177	0.181
Te	0.001	0.063	0.952	0.244	0.249
B50	-0.001	0.064	0.954	0.246	0.248
B75	-0.001	0.064	0.954	0.246	0.248
B90	-0.001	0.064	0.954	0.246	0.248
B95	-0.001	0.064	0.954	0.246	0.248
MA	0.008	0.068	0.948	0.264	0.255
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$, Effect size: 0.57					
Na_min	0.003	0.064	0.950	0.243	0.248
Na_pool	0.349	0.353	0.000	0.221	0.224
Te	0.002	0.065	0.946	0.252	0.249

B50	0.005	0.063	0.948	0.245	0.248
B75	0.005	0.063	0.948	0.245	0.248
B90	0.005	0.063	0.948	0.245	0.248
B95	0.005	0.063	0.948	0.245	0.248
MA	-0.005	0.063	0.952	0.247	0.252
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Na_min	-0.003	0.064	0.946	0.251	0.249
Na_pool	0.351	0.354	0.000	0.173	0.186
Te	0.003	0.064	0.956	0.243	0.249
B50	0.002	0.062	0.956	0.234	0.249
B75	0.002	0.062	0.956	0.234	0.249
B90	0.002	0.062	0.956	0.234	0.249
B95	0.002	0.062	0.956	0.234	0.249
MA	-0.001	0.064	0.942	0.259	0.248
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na_min	0.002	0.065	0.936	0.262	0.248
Na_pool	0.347	0.349	0.000	0.146	0.164
Te	0.000	0.062	0.936	0.254	0.249
B50	0.000	0.061	0.954	0.243	0.248
B75	0.000	0.061	0.954	0.243	0.248
B90	0.000	0.061	0.954	0.243	0.248
B95	0.000	0.061	0.954	0.243	0.248
MA	0.002	0.063	0.950	0.238	0.246

Notes: "Na_min" refers to the Naïve approach that takes the smaller estimate, "Na_pool" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A4: Performances in Normal Outcomes When There is no Information about Preferred Modes or Directions ($n = 500$ and $\rho = 0.75$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Na	-0.001	0.043	0.946	0.175	0.176
Te	-0.004	0.045	0.948	0.175	0.193
B50	0.004	0.045	0.982	0.178	0.250
B75	0.004	0.045	0.964	0.178	0.224
B90	0.004	0.045	0.956	0.178	0.201
B95	0.004	0.045	0.954	0.178	0.191
MA	0.001	0.047	0.940	0.189	0.183
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na	-0.007	0.052	0.970	0.191	0.215
Te	0.002	0.054	0.958	0.214	0.220
B50	-0.002	0.055	0.974	0.211	0.302
B75	-0.002	0.055	0.958	0.212	0.261
B90	-0.002	0.053	0.950	0.207	0.232
B95	-0.002	0.052	0.950	0.203	0.217
MA	0.002	0.054	0.958	0.196	0.211
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na	0.149	0.158	0.234	0.209	0.217
Te	0.162	0.175	0.880	0.254	0.535
B50	0.149	0.159	0.952	0.218	0.553
B75	0.149	0.159	0.948	0.222	0.549
B90	0.146	0.157	0.900	0.228	0.528
B95	0.143	0.155	0.860	0.233	0.505
MA	0.129	0.142	0.768	0.228	0.420
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na	0.152	0.159	0.074	0.177	0.177

Te	0.149	0.156	0.904	0.173	0.527
B50	0.151	0.157	0.960	0.167	0.510
B75	0.151	0.157	0.954	0.167	0.509
B90	0.151	0.157	0.942	0.167	0.505
B95	0.151	0.157	0.900	0.169	0.496
MA	0.149	0.155	0.794	0.168	0.452
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$. Effect size: 0.41					
Na	0.252	0.257	0.004	0.200	0.220
Te	0.273	0.279	0.984	0.214	0.803
B50	0.252	0.258	0.944	0.232	0.747
B75	0.252	0.258	0.944	0.232	0.746
B90	0.252	0.258	0.944	0.232	0.746
B95	0.252	0.258	0.942	0.232	0.745
MA	0.243	0.249	0.888	0.207	0.726
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$. Effect size: 0.50					
Na	0.250	0.255	0.000	0.191	0.181
Te	0.248	0.252	0.978	0.175	0.749
B50	0.252	0.256	0.936	0.172	0.710
B75	0.252	0.256	0.936	0.172	0.710
B90	0.252	0.256	0.936	0.172	0.710
B95	0.252	0.256	0.936	0.172	0.710
MA	0.245	0.249	0.940	0.174	0.700
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$. Effect size: 0.57					
Na	0.354	0.357	0.000	0.196	0.224
Te	0.376	0.380	0.978	0.199	1.007
B50	0.351	0.355	0.960	0.212	0.961
B75	0.351	0.355	0.960	0.212	0.961
B90	0.351	0.355	0.960	0.212	0.961
B95	0.351	0.355	0.960	0.212	0.961
MA	0.353	0.357	0.944	0.215	0.959
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					

Na	0.346	0.349	0.000	0.170	0.186
Te	0.353	0.356	0.982	0.176	0.946
B50	0.352	0.355	0.952	0.172	0.910
B75	0.352	0.355	0.952	0.172	0.910
B90	0.352	0.355	0.952	0.172	0.910
B95	0.352	0.355	0.952	0.172	0.910
MA	0.349	0.352	0.952	0.173	0.904
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na	0.350	0.352	0.000	0.152	0.164
Te	0.334	0.336	0.970	0.150	0.909
B50	0.350	0.352	0.944	0.152	0.882
B75	0.350	0.352	0.944	0.152	0.882
B90	0.350	0.352	0.944	0.152	0.882
B95	0.350	0.352	0.944	0.152	0.882
MA	0.345	0.347	0.950	0.151	0.880

Notes: "Na" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A5: Performances in Normal Outcomes When Smaller Estimates Are Preferred ($n = 500$ and $\rho = 0.5$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
NNa_min	-0.032	0.063	0.924	0.214	0.218
Na_pool	0.004	0.044	0.962	0.164	0.176
Te	-0.006	0.048	0.946	0.193	0.179

B50	-0.030	0.062	0.922	0.211	0.213
B75	-0.020	0.060	0.916	0.212	0.195
B90	-0.011	0.054	0.924	0.213	0.184
B95	-0.006	0.051	0.932	0.205	0.180
MA	-0.001	0.048	0.942	0.183	0.181
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na_min	-0.052	0.086	0.912	0.266	0.267
Na_pool	0.004	0.055	0.960	0.209	0.216
Te	-0.011	0.066	0.928	0.249	0.210
B50	-0.037	0.078	0.920	0.260	0.253
B75	-0.026	0.073	0.918	0.264	0.229
B90	-0.014	0.065	0.926	0.261	0.214
B95	-0.010	0.062	0.932	0.242	0.209
MA	-0.001	0.051	0.962	0.198	0.210
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na_min	-0.006	0.062	0.954	0.236	0.245
Na_pool	0.151	0.161	0.216	0.204	0.217
Te	0.013	0.077	0.860	0.284	0.237
B50	0.003	0.066	0.936	0.254	0.247
B75	0.004	0.068	0.920	0.256	0.245
B90	0.009	0.073	0.886	0.289	0.241
B95	0.014	0.078	0.852	0.304	0.238
MA	0.047	0.091	0.838	0.280	0.263
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na_min	0.004	0.062	0.948	0.247	0.246
Na_pool	0.152	0.159	0.070	0.180	0.178
Te	0.006	0.069	0.900	0.283	0.243
B50	-0.004	0.066	0.936	0.260	0.247
B75	-0.004	0.067	0.930	0.267	0.247
B90	-0.002	0.070	0.918	0.283	0.245
B95	0.001	0.076	0.890	0.324	0.242

MA	0.042	0.099	0.830	0.319	0.278
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$. Effect size: 0.41					
Na_min	-0.003	0.060	0.948	0.243	0.247
Na_pool	0.255	0.260	0.004	0.206	0.220
Te	0.001	0.067	0.930	0.264	0.248
B50	-0.001	0.063	0.948	0.241	0.248
B75	-0.001	0.063	0.948	0.241	0.248
B90	-0.001	0.063	0.948	0.241	0.248
B95	-0.001	0.063	0.946	0.245	0.248
MA	0.010	0.075	0.922	0.288	0.268
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$. Effect size: 0.50					
Na_min	0.000	0.061	0.954	0.242	0.247
Na_pool	0.248	0.252	0.000	0.176	0.181
Te	0.003	0.062	0.956	0.237	0.248
B50	0.000	0.060	0.962	0.238	0.248
B75	0.000	0.060	0.962	0.238	0.248
B90	0.000	0.060	0.962	0.238	0.248
B95	0.000	0.061	0.962	0.238	0.248
MA	-0.001	0.065	0.960	0.253	0.253
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$. Effect size: 0.57					
Na_min	0.004	0.064	0.952	0.241	0.249
Na_pool	0.352	0.356	0.000	0.206	0.224
Te	0.002	0.063	0.960	0.238	0.250
B50	-0.001	0.067	0.938	0.266	0.248
B75	-0.001	0.067	0.938	0.266	0.248
B90	-0.001	0.067	0.938	0.266	0.248
B95	-0.001	0.067	0.938	0.266	0.248
MA	0.001	0.062	0.952	0.243	0.252
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Na_min	-0.003	0.064	0.946	0.252	0.249
Na_pool	0.353	0.355	0.000	0.181	0.186

Te	-0.001	0.060	0.966	0.227	0.249
B50	0.002	0.062	0.950	0.243	0.249
B75	0.002	0.062	0.950	0.243	0.249
B90	0.002	0.062	0.950	0.243	0.249
B95	0.002	0.062	0.950	0.243	0.249
MA	-0.001	0.058	0.972	0.227	0.247
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na_min	0.002	0.065	0.938	0.259	0.248
Na_pool	0.348	0.350	0.000	0.157	0.165
Te	-0.002	0.063	0.948	0.251	0.250
B50	-0.002	0.065	0.940	0.252	0.248
B75	-0.002	0.065	0.940	0.252	0.248
B90	-0.002	0.065	0.940	0.252	0.248
B95	-0.002	0.065	0.940	0.252	0.248
MA	-0.001	0.064	0.948	0.248	0.247

Notes: "Na_min" refers to the Naïve approach that takes the smaller estimate, "Na_pool" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A6: Performances in Normal Outcomes When There is no Information about Preferred Modes or Directions ($n = 500$ and $\rho = 0.5$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Na	-0.002	0.043	0.956	0.170	0.176

Te	-0.001	0.045	0.944	0.179	0.194
B50	0.000	0.044	0.974	0.175	0.251
B75	0.000	0.044	0.960	0.175	0.222
B90	0.000	0.044	0.958	0.175	0.193
B95	0.000	0.044	0.956	0.175	0.186
MA	-0.001	0.043	0.966	0.166	0.181
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Na	-0.002	0.056	0.948	0.223	0.215
Te	0.000	0.056	0.934	0.219	0.221
B50	-0.003	0.053	0.978	0.211	0.300
B75	-0.003	0.052	0.964	0.210	0.261
B90	-0.003	0.052	0.956	0.201	0.228
B95	-0.003	0.051	0.956	0.199	0.218
MA	-0.001	0.048	0.966	0.187	0.213
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Na	0.150	0.160	0.236	0.212	0.217
Te	0.169	0.180	0.878	0.244	0.550
B50	0.148	0.157	0.938	0.213	0.547
B75	0.147	0.157	0.926	0.214	0.541
B90	0.145	0.156	0.902	0.220	0.526
B95	0.142	0.153	0.862	0.227	0.505
MA	0.128	0.141	0.802	0.239	0.434
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Na	0.149	0.156	0.086	0.166	0.177
Te	0.154	0.160	0.900	0.176	0.534
B50	0.152	0.158	0.948	0.178	0.509
B75	0.152	0.158	0.948	0.178	0.508
B90	0.152	0.158	0.930	0.178	0.502
B95	0.152	0.158	0.902	0.178	0.495
MA	0.151	0.158	0.808	0.179	0.453
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$. Effect size: 0.41					

Na	0.252	0.258	0.008	0.222	0.220
Te	0.274	0.280	0.966	0.213	0.793
B50	0.250	0.256	0.946	0.218	0.754
B75	0.250	0.256	0.946	0.218	0.754
B90	0.250	0.256	0.946	0.218	0.754
B95	0.250	0.256	0.944	0.218	0.753
MA	0.247	0.254	0.924	0.231	0.735
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$. Effect size: 0.50					
Na	0.247	0.251	0.000	0.184	0.181
Te	0.254	0.258	0.966	0.181	0.746
B50	0.254	0.258	0.940	0.173	0.716
B75	0.254	0.258	0.940	0.173	0.716
B90	0.254	0.258	0.940	0.173	0.716
B95	0.254	0.258	0.940	0.173	0.716
MA	0.252	0.256	0.934	0.174	0.699
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$. Effect size: 0.57					
Na	0.350	0.354	0.000	0.214	0.224
Te	0.377	0.380	0.968	0.204	1.006
B50	0.350	0.355	0.924	0.213	0.941
B75	0.350	0.355	0.924	0.213	0.941
B90	0.350	0.355	0.924	0.213	0.941
B95	0.350	0.355	0.924	0.213	0.941
MA	0.350	0.354	0.956	0.199	0.951
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Na	0.352	0.354	0.000	0.167	0.187
Te	0.351	0.354	0.974	0.175	0.955
B50	0.350	0.353	0.952	0.174	0.908
B75	0.350	0.353	0.952	0.174	0.908
B90	0.350	0.353	0.952	0.174	0.908
B95	0.350	0.353	0.952	0.174	0.908

MA	0.351	0.354	0.952	0.180	0.906
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Na	0.349	0.351	0.000	0.163	0.164
Te	0.331	0.333	0.970	0.155	0.915
B50	0.353	0.355	0.962	0.154	0.882
B75	0.353	0.355	0.962	0.154	0.882
B90	0.353	0.355	0.962	0.154	0.882
B95	0.353	0.355	0.962	0.154	0.882
MA	0.351	0.353	0.958	0.149	0.879

Notes: "Na" refers to the Naïve approach that pools the mixed-mode data, "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.

Table A7: Probabilities of the Model Being Selected in the Testimator Approach ($n = 5000$ and $\rho = 0.95$)

Scenarios	Effect Sizes	Models (Mean-Variance)			
		M1(D-D)	M2(S-D)	M3(D-S)	M4(S-S)
1. $u_b = 0, \sigma_b^2 = 1$	0.00	0.002	0.036	0.048	0.914
2. $u_b = 0, \sigma_b^2 = 2$	0.00	0.040	0.960	0.000	0.000
3. $u_b = 0.3, \sigma_b^2 = 2$	0.24	1.000	0.000	0.000	0.000
4. $u_b = 0.3, \sigma_b^2 = 1$	0.30	0.052	0.000	0.948	0.000
5. $u_b = 0.5, \sigma_b^2 = 2$	0.41	1.000	0.000	0.000	0.000
6. $u_b = 0.5, \sigma_b^2 = 1$	0.50	0.062	0.000	0.938	0.000
7. $u_b = 0.7, \sigma_b^2 = 2$	0.57	1.000	0.000	0.000	0.000
8. $u_b = 0.7, \sigma_b^2 = 1$	0.70	0.042	0.000	0.958	0.000
9. $u_b = 0.7, \sigma_b^2 = 0.5$	0.81	1.000	0.000	0.000	0.000

Notes: "D" stands for different and "S" stands for same. The correct model of a scenario are marked in bold in the table. Model 1 corresponds to the different mean different variance model, Model 2 is the same mean different variance model, Model 3 is the different mean and same variance model, and Model 4 is the same mean and same variance model.

Effect sizes are computed as $\frac{u_a - u_b}{\sqrt{\frac{\sigma_a^2 + \sigma_b^2}{2}}}$.

Table A8: Average Weight of Model Averaging Approach ($n = 5000$ and $\rho = 0.95$)

Scenarios	Effect Sizes	Models (Mean-Variance)			
		M1(D-D)	M2(S-D)	M3(D-S)	M4(S-S)
1. $u_b = 0, \sigma_b^2 = 1$	0.00	0.000	0.000	0.002	0.998
2. $u_b = 0, \sigma_b^2 = 2$	0.00	0.005	0.995	0.000	0.000
3. $u_b = 0.3, \sigma_b^2 = 2$	0.24	1.000	0.000	0.000	0.000
4. $u_b = 0.3, \sigma_b^2 = 1$	0.30	0.000	0.000	1.000	0.000
5. $u_b = 0.5, \sigma_b^2 = 2$	0.41	1.000	0.000	0.000	0.000
6. $u_b = 0.5, \sigma_b^2 = 1$	0.50	0.002	0.000	0.998	0.000
7. $u_b = 0.7, \sigma_b^2 = 2$	0.57	1.000	0.000	0.000	0.000
8. $u_b = 0.7, \sigma_b^2 = 1$	0.70	0.000	0.000	1.000	0.000
9. $u_b = 0.7, \sigma_b^2 = 0.5$	0.81	1.000	0.000	0.000	0.000

Notes: Correct models are marked in bold in the table. We differentiate each model by whether it assumes the same mean and same variances across modes. When specifying a scenario, "Same" means no shift in mean/variances, "Small" and "Large" means small and large shifts in means/variances respectively, and "Medium" means medium shifts in means.

Table A9: Performances in Normal Outcomes When Smaller
Estimates Are Preferred ($n = 5000$ and $\rho = 0.95$)

Method	Bias	RMSE	Coverage	Expected Length	Actual Length
Scenario 1: $u_b = 0$, $\sigma_b^2 = 1$. Effect size: 0					
Te	-0.001	0.016	0.946	0.062	0.056
B50	-0.008	0.019	0.924	0.065	0.066
B75	-0.005	0.018	0.916	0.069	0.061
B90	-0.002	0.017	0.922	0.065	0.057
B95	-0.001	0.016	0.934	0.059	0.056
MA	-0.001	0.014	0.954	0.054	0.056
Scenario 2: $u_b = 0$, $\sigma_b^2 = 2$. Effect size: 0					
Te	-0.002	0.019	0.940	0.075	0.065
B50	-0.011	0.023	0.948	0.076	0.077
B75	-0.007	0.021	0.948	0.077	0.070
B90	-0.004	0.019	0.954	0.072	0.066
B95	-0.003	0.018	0.954	0.067	0.065
MA	-0.001	0.016	0.952	0.063	0.065
Scenario 3: $u_b = 0.3$, $\sigma_b^2 = 2$. Effect size: 0.24					
Te	-0.001	0.020	0.942	0.081	0.079
B50	-0.001	0.021	0.958	0.077	0.078
B75	-0.001	0.021	0.958	0.077	0.078
B90	-0.001	0.021	0.958	0.077	0.078
B95	-0.001	0.021	0.958	0.077	0.078
MA	-0.001	0.020	0.948	0.079	0.078
Scenario 4: $u_b = 0.3$, $\sigma_b^2 = 1$, Effect size: 0.30					
Te	0.000	0.019	0.968	0.072	0.079
B50	0.001	0.020	0.936	0.082	0.078
B75	0.001	0.020	0.936	0.082	0.078
B90	0.001	0.020	0.936	0.082	0.078
B95	0.001	0.020	0.936	0.082	0.078

MA	0.001	0.020	0.954	0.076	0.078
Scenario 5: $u_b = 0.5$, $\sigma_b^2 = 2$. Effect size: 0.41					
Te	-0.001	0.019	0.968	0.073	0.079
B50	-0.001	0.020	0.958	0.076	0.078
B75	-0.001	0.020	0.958	0.076	0.078
B90	-0.001	0.020	0.958	0.076	0.078
B95	-0.001	0.020	0.958	0.076	0.078
MA	0.000	0.020	0.948	0.078	0.078
Scenario 6: $u_b = 0.5$, $\sigma_b^2 = 1$. Effect size: 0.50					
Te	0.001	0.020	0.952	0.077	0.079
B50	-0.001	0.020	0.954	0.077	0.078
B75	-0.001	0.020	0.954	0.077	0.078
B90	-0.001	0.020	0.954	0.077	0.078
B95	-0.001	0.020	0.954	0.077	0.078
MA	0.001	0.020	0.950	0.077	0.078
Scenario 7: $u_b = 0.7$, $\sigma_b^2 = 2$. Effect size: 0.57					
Te	-0.002	0.020	0.956	0.076	0.079
B50	0.001	0.021	0.946	0.079	0.078
B75	0.001	0.021	0.946	0.079	0.078
B90	0.001	0.021	0.946	0.079	0.078
B95	0.001	0.021	0.946	0.079	0.078
MA	0.000	0.019	0.972	0.070	0.078
Scenario 8: $u_b = 0.7$, $\sigma_b^2 = 1$. Effect size: 0.70					
Te	0.000	0.020	0.958	0.073	0.079
B50	0.000	0.020	0.960	0.075	0.078
B75	0.000	0.020	0.960	0.075	0.078
B90	0.000	0.020	0.960	0.075	0.078
B95	0.000	0.020	0.960	0.075	0.078
MA	0.000	0.019	0.948	0.078	0.078
Scenario 9: $u_b = 0.7$, $\sigma_b^2 = 0.5$. Effect size: 0.81					
Te	0.000	0.020	0.968	0.074	0.079

B50	0.000	0.021	0.936	0.080	0.078
B75	0.000	0.021	0.936	0.080	0.078
B90	0.000	0.021	0.936	0.080	0.078
B95	0.000	0.021	0.936	0.080	0.078
MA	0.000	0.019	0.956	0.073	0.078

Notes: "Te" refers to the testimator approach, "B50" refers to the Bayesian approach with 50% interval, "B75" refers to the Bayesian approach with 75% interval, "B90" is the Bayesian approach with 90% interval, "B95" is the Bayesian approach with 95% interval, and "MA" refers to the Model Averaging approach.