

Title: Protected area planning to conserve biodiversity in an uncertain world

Authors: Richard Schuster^{1,2*}, Rachel Buxton¹, Jeffrey O. Hanson¹, Allison D. Binley¹, Jeremy Pittman³, Vivitskaia Tulloch⁴, Frank A. La Sorte⁵, Patrick R. Roehrdanz⁶, Peter H. Verburg⁷, Amanda D. Rodewald^{5,8}, Scott Wilson^{1,9}, Hugh P. Possingham¹⁰, Joseph R. Bennett¹

Affiliations:

¹ Department of Biology, 1125 Colonel By Drive, Carleton University, Ottawa ON, K1S 5B6 Canada.

² Nature Conservancy of Canada, 245 Eglinton Ave East, Suite 410, Toronto, Ontario, M4P 3J1, Canada.

³ School of Planning, University of Waterloo, 200 University Ave W, Waterloo, ON, N2T 3G1, Canada

⁴ Conservation Decisions Lab, Department of Forest and Conservation Sciences, 2424 Main Mall, Vancouver, BC V6T 1Z4, Canada.

⁵ Cornell Lab of Ornithology, Cornell University, Ithaca, NY 14850, USA

⁶ Conservation International, 2100 Crystal Drive #600, Arlington, VA 22202, USA

⁷ Institute for Environmental Studies, VU University Amsterdam, Amsterdam, The Netherlands

⁸ Department of Natural Resources and the Environment, Cornell University, Ithaca, NY 14853, USA.

⁹ Wildlife Research Division, Environment and Climate Change Canada, 1125 Colonel By Drive, Ottawa, Ontario, Canada, K1S 5B6

¹⁰ Centre for Biodiversity and Conservation Science, University of Queensland, St Lucia, Queensland 4067, Australia

*Corresponding author. Email: richard.schuster@glel.carleton.ca

- 24 Running title: Protected area planning in an uncertain world
- 25 Keywords: biodiversity, protected areas, optimization, risk, governance, land use, climate change
- 26 Number of words (Abstract): 150
- 27 Number of words (Manuscript): 2863
- 28 Number of figures: 4
- 29 Number of tables: 0
- 30 Number of references: 40

Abstract:

Protected areas are a key instrument for conservation. Despite this, they are vulnerable to risks associated with weak governance, land use intensification, and climate change. Using a novel hierarchical optimization approach, we identified priority areas for expanding the global protected area system to explicitly account for such risks whilst maximizing protection of all known terrestrial vertebrate species. We illustrate how reducing exposure to these risks only requires expanding the area of the global protected area system by 1.6% while still meeting conservation targets. Incorporating risks from weak governance drove the greatest changes in spatial priorities for protection, while incorporating risks from climate change required the largest increase in global protected area. Conserving wide-ranging species required countries with relatively strong governance to protect more land when bordering nations with comparatively weak governance. Our results underscore the need for cross-jurisdictional coordination and demonstrate how risk can be efficiently incorporated into conservation planning.

Introduction

Protecting land is one of the best strategies to stem the biodiversity crisis (Watson et al. 2014) and a cornerstone of international agreements to safeguard biodiversity (Convention on Biological Diversity (CBD) 2017; CBD 2020). The effectiveness of a global network of protected areas depends upon the identification of areas that will both meet the needs of species and provide the greatest return on conservation investments. Yet most spatial planning efforts base decisions heavily upon the estimated ecological value of land (Brooks et al. 2006; Venter et al. 2014; CBD 2017; CBD 2020) and carry the tacit, but often incorrect, assumption that protection will be enforced, effective, and permanent. Despite evidence that protected areas are subject to risks associated with weak governance (e.g., political instability and corruption; Schulze et al. 2018), land use intensification (e.g., deforestation and degazetting of parks; Tesfaw et al. 2018), and climate change (e.g., extreme weather events; Maxwell et al. 2019), few plans have explicitly considered these risks (McBride et al. 2007; Alagador et al. 2014). Here we demonstrate how we can expand the planning lens to include risks as well as ecological value, with the goal of improving the resilience and performance of protected areas.

We considered the following three broad categories of risk, which we defined as factors likely to diminish the long-term effectiveness of protected areas: (i) governance, (ii) land use, and (iii) climate. We then generated plans for establishing protected areas (“prioritizations”) based on scenarios for different risk factors. Our framework provides a flexible approach for incorporating multiple risk metrics into conservation decision making.

Methods

We considered the influence of risk categories on allocating protection decisions at a global scale in suitable habitat for all 29,350 vertebrate species from the IUCN Red List of

Threatened Species (IUCN 2019) using a multi-objective optimization approach. To incorporate risk categories, we built on the minimum set problem, where the objective is to reach species distribution protection targets while accounting for one constraint such as land cost or area (Margules & Pressey 2000; Moilanen et al. 2009). We expanded this approach to include multiple objectives accounting for risk in the problem formulation, by treating each risk layer as a separate objective in the problem formulation (Deb 2014).

Biodiversity Data

We produced area of habitat (AOH) estimates for 10,774 species of birds, 5,219 mammals, 4,462 reptiles and 6,254 amphibians with available IUCN range polygon data following the procedure outlined in (Brooks et al. 2019). Species range polygons obtained from the IUCN Red List spatial data (<https://www.iucnredlist.org/>) and Birdlife International (<http://datazone.birdlife.org/species/requestdis>) were first filtered for ‘extant’ range then rasterized to a global 1 km grid in the Eckert IV equal area projection. Individual species range rasters were then modified to only include land cover classes that match the habitat associations for each species. Habitat associations were obtained from the IUCN Red List species habitat classification scheme and were matched to ESA land cover classes for the year 2018 following Santini et al. 2019. ESA land cover classification data was aggregated from 300 m resolution to match the global 1 km grid using a majority rule. Species ranges were additionally filtered so that only areas within a species’ accepted elevational range were included. Global elevation data derived from SRTM was obtained from WorldClim v. 2 (Fick & Hijmans 2017). For bird species with multiple seasonal distributions, data for resident, breeding, and non-breeding ranges were processed separately.

Basic administrative delineations

National boundaries were derived from the Global Administrative Areas database (<http://gadm.org>). We obtained protected area boundaries from the World Database on Protected Areas (<https://www.protectedplanet.net>). Following standard procedures for cleaning the protected area dataset (Butchart et al. 2015), we (i) projected the data to an equal-area coordinate system (World Behrman), (ii) excluded reserves with unknown or proposed designations, (iii) excluded UNESCO Biosphere Reserves (Coetzer et al. 2014), (iv) buffered sites represented as point localities to their reported area, (v) dissolved boundaries to prevent issues with overlapping areas, and (vi) removed slivers (code available at <https://github.com/jeffreyhanson/global-protected-areas>). After the protected area data were modified as described above, we overlaid the protected area boundaries with a 10 x 10 km grid covering the Earth. These spatial data procedures were implemented using ArcMap (version 10.3.1) and python (version 2.7.8).

Governance risk

We used worldwide governance indicators from the World Bank to capture governance risk (Kaufmann et al. 2011). The indicators include six scaled measures: voice and accountability; political stability and absence of violence; government effectiveness; regulatory quality; rule of law; and control of corruption (see Table S4 for definitions). We chose these indicators because evidence suggests that they reliably predict protected area effectiveness (Barnes et al. 2016) and state investment and efforts for biodiversity conservation (Coetzer et al. 2014). For each country, we used a mean of annual averages of all six measures (Baynham-Herd et al. 2018) (Figure S1).

Land use risk

We used a global land systems map produced by Kehoe et al. (2017) to incorporate the risk of land-use change. This map is based on a global land systems map for the year 2000 (Asselen & Verburg 2012) at a 9.25 km² spatial resolution but is refined based on recent land-cover and land-use datasets to a spatial resolution of 1 km². Kehoe et al. (2017) further estimated the impact of land use and land use intensity on biodiversity, with data originating from the PREDICTS project (Hudson et al. 2014). They first matched their land-systems classes to varying intensity levels for each land use type (for detailed conversion table, see Asselen & Verburg 2012). This allowed Kehoe et al. (2017) to calculate average biodiversity loss per land system (relative to an unimpacted baseline) by taking the mean model estimates of biodiversity loss per land use intensity class from previous work (Hudson et al. 2014). The result gives average relative biodiversity gain or loss per land-system class. Here, we used their modelled mean estimates (following Newbold et al. (2015)) of relative percent biodiversity change for each land-system class for species abundance as a measure of the land use pressure (Figure S2).

Climate risk

We used, velocity of climate change, which is an instantaneous measurement of how projected temperature increases translate to horizontal climate velocity on the landscape (Loarie et al. 2009). It is an integration of both the rate of change in average climate and landscape properties that govern how bands of similar temperature redistribute spatially as climate changes. For example, in a region with high topographic diversity, a species may be able to track its climatic niche through relatively small dispersal distances (e.g. 10s or 100s of meters) upslope or downslope. By contrast, keeping pace with preferred climate under the same magnitude of temperature rise in the plains may require much larger dispersal distances – 100s or 1000s of kilometers. Velocity of future temperature change used here follows the method of Loarie et al.

(2009), and is essentially the ratio of the projected temporal rate of change (C/year) to the spatial rate of change (C/km). Projected temporal rate of change was based on the 20-year mean (2040-2060) projection for mean annual temperature from the HadGEM2-ES model (CMIP5) and the baseline (1960-1990) temperature available from Worldclim v1.4. Spatial rate of change was derived from 30 arc second elevation data and calculated with the ‘terrain’ function from the R ‘raster’ package.

We also explored an alternative measure of climate risk: exposure to extreme events. Detailed methods and results for this alternative measure are provided in the online Supporting Information.

Multi-objective optimization of risk reduction

We created 16 planning scenarios, such that solutions accounted for all possible combinations of risk categories within each hierarchical level (Table S1). We then compared these risk-based solutions to those produced with a null scenario that adopted the traditional area-minimizing approach to optimization without considering risk.

We processed all data described previously to a 10 x 10 km resolution and clipped data to the extent of land based on the global administrative areas database. Our multi-objective approach uses a hierarchical (lexicographic) framework that assigns a priority to each objective, and sequentially optimizes for the objectives in order of decreasing priority. At each step, it finds the best solution for the current objective, but only from among those that would not degrade the solution quality for higher-priority objectives. We considered up to three objectives in our prioritization scenarios, i) governance risk, ii) land use risk, and iii) climate risk. To compare different scenarios, we calculated solutions for each unique objective combination ($n = 15$), as well as one where we use a constant objective function as the null scenario, as the order of the

hierarchy can influence the results. For a detailed mathematical exploration of this method, see the online Supporting Information.

For all scenarios we locked in current protected areas. Following Hanson et al. (2020), we used flexible targets for suitable habitat based on species' ranges. Species with less than 1,000 km² of suitable habitat were assigned a 100% target (1,802 amphibians, 893 avian and 645 mammalian species), species with more than 250,000 km² of suitable habitat were assigned a 10% target (712 amphibians, 4,518 avian and 1,868 mammalian species) and species with an intermediate amount of suitable habitat were assigned a target by log-linearly interpolating values between the previous two thresholds (2,683 amphibians, 5,190 avian and 2,557 mammalian species). Migratory bird species were assigned targets for each seasonal distribution separately. Additionally, to prevent species with very large suitable habitats from requiring excessively large amounts of area to be protected, the targets for species' distributions larger than 10,000,000 km² were capped at 1,000,000 km². This upper limit affected only 206 (1%) species, and sensitivity analyses showed that it had negligible effect on our results. We acknowledge that these targets are arbitrary; however, they are more precise than previous targets based on species' ranges (which can contain a large amount of unsuitable habitat), and account for the increased vulnerability of species with smaller range sizes (Pimm & Raven 2000), as well as the difficulty in conserving all habitat for species that occur over large areas.

Results

Surprisingly, scenarios that incorporated combinations of the three risk categories increased the priority area by only 1.6% on average (0.08 – 2.52%) compared to the null scenario based solely on ecological value to species. Among single-risk scenarios, accommodating risks

due to climate change velocity required the greatest increase in global protected area, compared to scenarios including only governance or land use intensification risks (Table S1).

Scenarios shared many overlapping spatial priorities, which can be considered as reliably good investments in terms of both ecological value and risk management. Most notably, all 15 scenarios prioritized the same 8.5 million km² (5.8% of global land area) (“no regrets” areas, Fig. 2), much of which was located in Canada, Kenya and Peru (Figure S4). There also was substantial overlap among the priorities across scenarios within Conservation International’s global biodiversity hotspots (Myers et al. 2000), but many high overlap areas lie either outside of (53.3%) or in small areas within hotspots (Figure S5).

At the same time, risk scenarios elicited several prominent shifts in spatial priorities among areas varying in risk exposure (Fig. 3; Table S2). In some cases, high risks to protected areas in weakly-governed countries could be compensated by expanding protected areas in well-governed neighboring nations (Fig S6). For example, challenges to the transborder conservation of the wide-ranging and IUCN-vulnerable caribou (*Rangifer tarandus*) due to weak governance in Russia (Table S3) were mitigated by increasing the land area protected by Finland from 16.2% to 36.4% (Fig. 4). High exposure to risks from land use change could be offset in a similar fashion, such as by protecting more land in Liberia (32% rather than 22.5% in null scenario) than in the agriculturally intensifying nation of Sierra Leone (Fig. 4). Likewise, climate-associated risks in Hungary and Serbia (Figure S3) might be tempered by protecting twice as much land (20.4% from 10.2% in null) in nearby Kosovo, which has lower predicted climate velocity (Fig. 4). Addressing risks from extreme weather events (La Sorte et al. 2021) (Figures S7 – S9) also required shifting some priority areas to less climatically volatile locations.

Discussion

Although a growing body of literature shows that protected areas are seldom effective if subject to unmitigated risks from land use change, weak governance and climate change (Schulze et al. 2018; Tesfaw et al. 2018; Maxwell et al. 2019), most systematic conservation planning efforts prioritize land based on ecological value and some measure of cost. We show how only small (1.6%) increases in land area, but importantly a change in the spatial configuration of where protected areas are places, may reduce the vulnerability of protected areas to future threats if risk are explicitly considered during planning stages (Fig. 1). Across all planning scenarios, we identified 8.5 million km² of priority lands that either uniquely contributed to conservation targets (e.g., high endemism) or were resilient to the risks we modeled. Countries with large proportions of land already in protection (e.g., Brazil with >30%) also had similar priorities for risk vs. null scenarios. Such shared priority areas across planning scenarios are likely to provide good return on conservation investments.

Despite modest differences across scenarios in the global land area required to conserve terrestrial vertebrates, shifts in the locations of priority areas sometimes resulted in substantial increases within any given nation. These asymmetries highlight the importance of cross-jurisdictional coordination to promote collaboration and improve the effectiveness of protected area systems (Dallimer & Strange 2015). In regions where nations vary widely in exposure to risks, coordination may provide opportunity to offset or otherwise mitigate risks by adjusting the geographic locations and/or boundaries of protected areas. Cooperative governance frameworks (Miller et al. 2019) are especially important for countries supporting wide-ranging species that are expected to be impacted by climate, land use, and governance risk across borders (Fig. 3). These governance frameworks, both within and among countries, would need to be developed in an environmentally just and equitable way to deliver benefits to biodiversity and local communities (Martin et al. 2013).

A novel contribution of our framework is that it explicitly incorporates multiple risk factors at the same time. Previous work has incorporated single risk factors analogous to those we used, including governance (Mascia & Pailler 2011; Eklund & Cabeza-Jaimejuan 2017), climate change (Hoffmann et al. 2019) and land use change (Pouzols et al. 2014; Di Minin et al. 2016) demonstrating the importance of each type of risk in protected area planning. Our results similarly demonstrate that protected area expansion decisions can be profoundly influenced by all three risk factors combined, yet they also show that relatively little additional protected area is required to account for these risks.

Our flexible framework and methods can also allow conservation agencies to set their own priorities from local to global scales and incorporate different metrics to assess the relevance of different forms and levels of risk. Indeed, our risk metrics were chosen as reasonable examples, rather than definitive recommendations. Our alternative climate risk framework (presented in the Supporting Information) illustrates the importance of metric choice. The difference between our climate risk scenarios highlights the need for agencies to carefully consider their choices of risk metrics and suggests that smaller-scale planning exercises should choose metrics that are most relevant for each region.

The conservation community has traditionally neglected to estimate how future changes in climate (Kelly et al. 2020), land use (Di Minin et al. 2016), and governance risk might compromise the effectiveness of protected areas. Yet, as we work towards an ambitious new plan to curb biodiversity loss (CBD 2020) in a rapidly changing world, we show that incorporating future risk has profound implications for the spatial distribution of protected areas. The risk of weak governance was particularly influential. Surprisingly, incorporating risk into decision-making adds <2% to the total global area required to meet biodiversity targets. Thus, accounting for risk comes at limited extra cost which is likely outweighed by increased likelihood of

achieving global biodiversity targets. Our results also emphasize the importance of cross-jurisdictional conservation initiatives, especially in adjacent countries sharing wide-ranging species where risk varies considerably from country to country. Considering risk in conservation decision-making will result in more resilient and effective conservation plans into the future to help safeguard our planet's biodiversity in the face of the current extinction crisis.

References

- Alagador, D., Cerdeira, J.O. & Araújo, M.B. (2014). Shifting protected areas: scheduling spatial priorities under climate change. *Journal of applied ecology*, 51, 703–713.
- Asselen, S. van & Verburg, P.H. (2012). A Land System representation for global assessments and land-use modeling. *Global Change Biology*, 18, 3125–3148.
- Barnes, M.D., Craigie, I.D., Harrison, L.B., Geldmann, J., Collen, B., Whitmee, S., Balmford, A., Burgess, N.D., Brooks, T., Hockings, M. & Woodley, S. (2016). Wildlife population trends in protected areas predicted by national socio-economic metrics and body size. *Nature Communications*, 7, 12747.
- Baynham-Herd, Z., Amano, T., Sutherland, W.J. & Donald, P.F. (2018). Governance explains variation in national responses to the biodiversity crisis. *Environmental Conservation*, 45, 407–418.
- Brooks, T.M., Mittermeier, R.A., da Fonseca, G.A., Gerlach, J., Hoffmann, M., Lamoreux, J.F., Mittermeier, C.G., Pilgrim, J.D. & Rodrigues, A.S. (2006). Global biodiversity conservation priorities. *science*, 313, 58–61.
- Brooks, T.M., Pimm, S.L., Akçakaya, H.R., Buchanan, G.M., Butchart, S.H.M., Foden, W., Hilton-Taylor, C., Hoffmann, M., Jenkins, C.N., Joppa, L., Li, B.V., Menon, V., Ocampo-Peñuela, N. & Rondinini, C. (2019). Measuring Terrestrial Area of Habitat

284 (AOH) and Its Utility for the IUCN Red List. *Trends in Ecology & Evolution*, 34, 977–
 285 986.

286 Butchart, S.H.M., Clarke, M., Smith, R.J., Sykes, R.E., Scharlemann, J.P.W., Harfoot, M.,
 287 Buchanan, G.M., Angulo, A., Balmford, A., Bertzky, B., Brooks, T.M., Carpenter, K.E.,
 288 Comeros-Raynal, M.T., Cornell, J., Ficetola, G.F., Fishpool, L.D.C., Fuller, R.A.,
 289 Geldmann, J., Harwell, H., Hilton-Taylor, C., Hoffmann, M., Joolia, A., Joppa, L.,
 290 Kingston, N., May, I., Milam, A., Polidoro, B., Ralph, G., Richman, N., Rondinini, C.,
 291 Segan, D.B., Skolnik, B., Spalding, M.D., Stuart, S.N., Symes, A., Taylor, J., Visconti,
 292 P., Watson, J.E.M., Wood, L. & Burgess, N.D. (2015). Shortfalls and Solutions for
 293 Meeting National and Global Conservation Area Targets. *Conservation Letters*, 8, 329–
 294 337.

295 CBD. (2020). Zero draft of the post-2020 global biodiversity framework [WWW Document].
 296 URL [https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-](https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-en.pdf)
 297 [en.pdf](https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-en.pdf)

298 Coetzer, K.L., Witkowski, E.T.F. & Erasmus, B.F.N. (2014). Reviewing Biosphere Reserves
 299 globally: effective conservation action or bureaucratic label? *Biological Reviews*, 89, 82–
 300 104.

301 Convention on Biological Diversity. (2017). Aichi Biodiversity Targets [WWW Document].
 302 URL <https://www.cbd.int/sp/targets/>

303 Dallimer, M. & Strange, N. (2015). Why socio-political borders and boundaries matter in
 304 conservation. *Trends in Ecology & Evolution*, 30, 132–139.

305 Deb, K. (2014). Multi-objective optimization. In: *Search methodologies*. Springer, pp. 403–449.

Di Minin, E., Slotow, R., Hunter, L.T.B., Montesino Pouzols, F., Toivonen, T., Verburg, P.H.,
 Leader-Williams, N., Petracca, L. & Moilanen, A. (2016). Global priorities for national
 carnivore conservation under land use change. *Scientific Reports*, 6, 23814.

Eklund, J.F. & Cabeza-Jaimejuan, M.D.M. (2017). Quality of governance and effectiveness of
 protected areas: crucial concepts for conservation planning. *Annals of the New York
 Academy of Sciences*.

Fick, S.E. & Hijmans, R.J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces
 for global land areas. *International Journal of Climatology*, 37, 4302–4315.

Hanson, J.O., Rhodes, J.R., Butchart, S.H.M., Buchanan, G.M., Rondinini, C., Ficetola, G.F. &
 Fuller, R.A. (2020). Global conservation of species' niches. *Nature*, 580, 232–234.

Hoffmann, S., Irl, S.D. & Beierkuhnlein, C. (2019). Predicted climate shifts within terrestrial
 protected areas worldwide. *Nature communications*, 10, 1–10.

Hudson, L.N., Newbold, T., Contu, S., Hill, S.L.L., Lysenko, I., De Palma, A., Phillips, H.R.P.,
 Senior, R.A., Bennett, D.J., Booth, H., Choimes, A., Correia, D.L.P., Day, J., Echeverría-
 Londoño, S., Garon, M., Harrison, M.L.K., Ingram, D.J., Jung, M., Kemp, V.,
 Kirkpatrick, L., Martin, C.D., Pan, Y., White, H.J., Aben, J., Abrahamczyk, S., Adum,
 G.B., Aguilar-Barquero, V., Aizen, M.A., Ancrenaz, M., Arbeláez-Cortés, E., Armbrecht,
 I., Azhar, B., Azpiroz, A.B., Baeten, L., Báldi, A., Banks, J.E., Barlow, J., Batáry, P.,
 Bates, A.J., Bayne, E.M., Beja, P., Berg, Å., Berry, N.J., Bicknell, J.E., Bihn, J.H.,
 Böhning-Gaese, K., Boekhout, T., Boutin, C., Bouyer, J., Brearley, F.Q., Brito, I., Brunet,
 J., Buczkowski, G., Buscardo, E., Cabra-García, J., Calviño-Cancela, M., Cameron, S.A.,
 Cancellato, E.M., Carrijo, T.F., Carvalho, A.L., Castro, H., Castro-Luna, A.A., Cerda, R.,
 Cerezo, A., Chauvat, M., Clarke, F.M., Cleary, D.F.R., Connop, S.P., D'Aniello, B., da
 Silva, P.G., Darvill, B., Dauber, J., Dejean, A., Diekötter, T., Dominguez-Haydar, Y.,

330 Dormann, C.F., Dumont, B., Dures, S.G., Dynesius, M., Edenius, L., Elek, Z., Entling,
 331 M.H., Farwig, N., Fayle, T.M., Felicioli, A., Felton, A.M., Ficetola, G.F., Filgueiras,
 332 B.K.C., Fonte, S.J., Fraser, L.H., Fukuda, D., Furlani, D., Ganzhorn, J.U., Garden, J.G.,
 333 Gheler-Costa, C., Giordani, P., Giordano, S., Gottschalk, M.S., Goulson, D., Gove, A.D.,
 334 Grogan, J., Hanley, M.E., Hanson, T., Hashim, N.R., Hawes, J.E., Hébert, C., Helden,
 335 A.J., Henden, J.-A., Hernández, L., Herzog, F., Higuera-Diaz, D., Hilje, B., Horgan, F.G.,
 336 Horváth, R., Hylander, K., Isaacs-Cubides, P., Ishitani, M., Jacobs, C.T., Jaramillo, V.J.,
 337 Jauker, B., Jonsell, M., Jung, T.S., Kapoor, V., Kati, V., Katovai, E., Kessler, M., Knop,
 338 E., Kolb, A., Kőrösi, Á., Lachat, T., Lantschner, V., Le Féon, V., LeBuhn, G., Légaré, J.-
 339 P., Letcher, S.G., Littlewood, N.A., López-Quintero, C.A., Louhaichi, M., Lövei, G.L.,
 340 Lucas-Borja, M.E., Luja, V.H., Maeto, K., Magura, T., Mallari, N.A., Marin-Spiotta, E.,
 341 Marshall, E.J.P., Martínez, E., Mayfield, M.M., Mikusinski, G., Milder, J.C., Miller, J.R.,
 342 Morales, C.L., Muchane, M.N., Muchane, M., Naidoo, R., Nakamura, A., Naoe, S.,
 343 Nates-Parra, G., Navarrete Gutierrez, D.A., Neuschulz, E.L., Noreika, N., Norfolk, O.,
 344 Noriega, J.A., Nöske, N.M., O’Dea, N., Oduro, W., Ofori-Boateng, C., Oke, C.O.,
 345 Osgathorpe, L.M., Paritsis, J., Parra-H, A., Pelegrin, N., Peres, C.A., Persson, A.S.,
 346 Petanidou, T., Phalan, B., Philips, T.K., Poveda, K., Power, E.F., Presley, S.J., Proença,
 347 V., Quaranta, M., Quintero, C., Redpath-Downing, N.A., Reid, J.L., Reis, Y.T., Ribeiro,
 348 D.B., Richardson, B.A., Richardson, M.J., Robles, C.A., Römbke, J., Romero-Duque,
 349 L.P., Rosselli, L., Rossiter, S.J., Roulston, T.H., Rousseau, L., Sadler, J.P., Sáfián, S.,
 350 Saldaña-Vázquez, R.A., Samnegård, U., Schüepp, C., Schweiger, O., Sedlock, J.L.,
 351 Shahabuddin, G., Sheil, D., Silva, F.A.B., Slade, E.M., Smith-Pardo, A.H., Sodhi, N.S.,
 352 Somarriba, E.J., Sosa, R.A., Stout, J.C., Struebig, M.J., Sung, Y.-H., Threlfall, C.G.,
 353 Tonietto, R., Tóthmérés, B., Tschantke, T., Turner, E.C., Tylianakis, J.M., Vanbergen,

A.J., Vassilev, K., Verboven, H.A.F., Vergara, C.H., Vergara, P.M., Verhulst, J., Walker, T.R., Wang, Y., Watling, J.I., Wells, K., Williams, C.D., Willig, M.R., Woinarski, J.C.Z., Wolf, J.H.D., Woodcock, B.A., Yu, D.W., Zaitsev, A.S., Collen, B., Ewers, R.M., Mace, G.M., Purves, D.W., Scharlemann, J.P.W. & Purvis, A. (2014). The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution*, 4, 4701–4735.

IUCN. (2019). The IUCN Red List of Threatened Species. version 1.18.

Kaufmann, D., Kraay, A. & Mastruzzi, M. (2011). The Worldwide Governance Indicators: Methodology and Analytical Issues1. *Hague Journal on the Rule of Law*, 3, 220–246.

Kehoe, L., Romero-Muñoz, A., Polaina, E., Estes, L., Kreft, H. & Kuemmerle, T. (2017). Biodiversity at risk under future cropland expansion and intensification. *Nature Ecology & Evolution*, 1, 1129–1135.

Kelly, L.T., Giljohann, K.M., Duane, A., Aquilué, N., Archibald, S., Batllori, E., Bennett, A.F., Buckland, S.T., Canelles, Q., Clarke, M.F., Fortin, M.-J., Hermoso, V., Herrando, S., Keane, R.E., Lake, F.K., McCarthy, M.A., Morán-Ordóñez, A., Parr, C.L., Pausas, J.G., Penman, T.D., Regos, A., Rumpff, L., Santos, J.L., Smith, A.L., Syphard, A.D., Tingley, M.W. & Brotons, L. (2020). Fire and biodiversity in the Anthropocene. *Science*, 370.

La Sorte, F.A., Johnston, A. & Ault, T.R. (2021). Global trends in the frequency and duration of temperature extremes. *Climatic Change*, 166, 1.

Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B. & Ackerly, D.D. (2009). The velocity of climate change. *Nature*, 462, 1052–1055.

Margules, C.R. & Pressey, R.L. (2000). Systematic conservation planning. *Nature*, 405, 243–53.

Martin, A., McGuire, S. & Sullivan, S. (2013). Global environmental justice and biodiversity conservation. *The Geographical Journal*, 179, 122–131.

- Mascia, M.B. & Pailler, S. (2011). Protected area downgrading, downsizing, and degazettement (PADDD) and its conservation implications. *Conservation Letters*, 4, 9–20.
- Maxwell, S.L., Butt, N., Maron, M., McAlpine, C.A., Chapman, S., Ullmann, A., Segan, D.B. & Watson, J.E.M. (2019). Conservation implications of ecological responses to extreme weather and climate events. *Diversity and Distributions*, 25, 613–625.
- McBride, M.F., Wilson, K.A., Bode, M. & Possingham, H.P. (2007). Incorporating the effects of socioeconomic uncertainty into priority setting for conservation investment. *Conservation Biology*, 21, 1463–1474.
- Miller, R.L., Marsh, H., Benham, C. & Hamann, M. (2019). A framework for improving the cross-jurisdictional governance of a marine migratory species. *Conservation Science and Practice*, 1, e58.
- Moilanen, A., Wilson, K. & Possingham, H. (2009). *Spatial conservation prioritization: quantitative methods and computational tools*. Oxford University Press.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.
- Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M.J., Feldman, A., Garon, M., Harrison, M.L.K., Alhusseini, T., Ingram, D.J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D.L.P., Martin, C.D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H.R.P., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White, H.J., Ewers, R.M., Mace, G.M., Scharlemann, J.P.W. & Purvis, A. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*, 520, 45–50.
- Pimm, S.L. & Raven, P. (2000). Extinction by numbers. *Nature*, 403, 843–845.

Pouzols, F.M., Toivonen, T., Di Minin, E., Kukkala, A.S., Kullberg, P., Kuusterä, J., Lehtomäki, J., Tenkanen, H., Verburg, P.H. & Moilanen, A. (2014). Global protected area expansion is compromised by projected land-use and parochialism. *Nature*, 516, 383–386.

Santini, L., Butchart, S.H.M., Rondinini, C., Benítez-López, A., Hilbers, J.P., Schipper, A.M., Cengic, M., Tobias, J.A. & Huijbregts, M.A.J. (2019). Applying habitat and population-density models to land-cover time series to inform IUCN Red List assessments. *Conservation Biology*, 33, 1084–1093.

Schulze, K., Knights, K., Coad, L., Geldmann, J., Leverington, F., Eassom, A., Marr, M., Butchart, S.H., Hockings, M. & Burgess, N.D. (2018). An assessment of threats to terrestrial protected areas. *Conservation Letters*, 11, e12435.

Tesfaw, A.T., Pfaff, A., Kroner, R.E.G., Qin, S., Medeiros, R. & Mascia, M.B. (2018). Land-use and land-cover change shape the sustainability and impacts of protected areas. *Proceedings of the National Academy of Sciences*, 115, 2084–2089.

Venter, O., Fuller, R.A., Segan, D.B., Carwardine, J., Brooks, T., Butchart, S.H.M., Di Marco, M., Iwamura, T., Joseph, L., O’Grady, D., Possingham, H.P., Rondinini, C., Smith, R.J., Venter, M. & Watson, J.E.M. (2014). Targeting Global Protected Area Expansion for Imperiled Biodiversity. *PLOS Biology*, 12, e1001891.


Watson, J.E., Dudley, N., Segan, D.B. & Hockings, M. (2014). The performance and potential of protected areas. *Nature*, 515, 67–73.

Acknowledgments: We thank Alison Johnston, Peter Arcese, Steven Cooke and Lenore Fahrig for helpful discussions. RS was funded by the Liber Ero Fellowship Program. JRB was funded by the Natural Science and Engineering Research Council of Canada (NSERC) Discovery Grant 2016-06147 and Environment and Climate Change Canada Grant GCXE19S058.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data, scripts and full results are available on Open Science Framework (OSF) and will be assigned a DOI once the manuscript is in print:

https://osf.io/e2fuw/?view_only=46eb2e525daf42d29df318a92762d885



433

434

435

436

Figure 1: Spatial representation of priority areas for protection to account for governance, land use and climate risk. Accounting for these risks to protected area effectiveness to produce more resilient conservation networks would require 23.5% of land surface to reach suitable habitat protection goals (26) for vertebrate species from the IUCN Red List of Threatened Species (20).





Figure 2: “No regrets” areas comprising 8.5 million km² of land that was identified as priority habitat for protection regardless of the risks included in our analysis.




441

442 **Figure 3:** Percent country-level variation between the null scenario and the 15 scenarios including risk. Countries whose results are
 443 consistent across the 15 scenarios (e.g., Mexico) have low variation, while countries whose results are less consistent across the 15
 444 scenarios (e.g., Finland) have high variation.

445

446



448

449 **Figure 4:** Contrast of using individual risk objectives (governance, land use, climate) to the null scenario of uniform objective structure.

450 The top panels represent the individual risk data for the focal regions. In the bottom panels brown shows null, green the specific risk
 451 objective scenario results, and purple where both scenarios agree. The figures show how the spatial configuration of the solutions
 452 changes when risk is considered in a scenario. Governance focus is on Finland and Russia, land use risk on Sierra Leone and Liberia, and
 453 climate risk on Serbia, Hungary and Kosovo.

454