

the *Journal of the American Medical Association* (JAMA) and the *New England Journal of Medicine* (NEJM).

For example, the JAMA article by the authors of the NEJM article

concludes that the results of the study are "consistent with the hypothesis

that the use of a single, low-dose, intravenous bolus of 100 mg of

propofol is associated with a lower risk of hypotension than the use of

multiple boluses of 20 mg each." The NEJM article, however, states

that the results of the study are "consistent with the hypothesis that the

use of a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The NEJM article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The NEJM article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The NEJM article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The NEJM article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The NEJM article, however, states that the

results of the study are "consistent with the hypothesis that the use of

a single, low-dose, intravenous bolus of 100 mg of propofol is

associated with a lower risk of hypotension than the use of multiple

boluses of 20 mg each." The JAMA article, however, states that the

results of the study are "consistent with the hypothesis that the use of