Introduction to Abstract Algebra (In Progress)

Nihal Uppugunduri

October 20, 2023

Contents

Preface **v**
1. Introduction v
2. Prerequisites and Notation v
3. Organization and Order of Topics vii
4. Practice Problems viii
5. Final Preliminaries viii

1. Primer: What is Abstract Algebra? **1**

2. Fields **7**
2.1. Definition and Basic Properties 7
2.2. Polynomials Over Fields 15
2.3. Isomorphisms and Homomorphisms 15
2.4. Subfields 16
2.5. Field Extensions 16
2.6. Construction of a Simple Algebraic Extension 16
2.7. Construction of Any Algebraic Extension 16

3. Vector Spaces **17**
3.1. Definition and Basic Properties 17
3.2. Finite-Dimensional Vector Spaces 17
3.3. Matrices and Linear Transformations 17
3.4. Matrices and Subspaces 17
3.5. General Bases 17
3.6. Field Extensions as Vector Spaces 18

4. Groups **19**
4.1. Definition and Basic Properties 19
4.2. Permutation and Symmetry Groups 19
4.3. Cyclic Groups 19
4.4. Order of an Element 19
4.5. Generators, Presentations, and Free Groups 19
4.6. Quotient Groups 20
4.7. The Isomorphism Theorems 20
4.8. Group Actions 20

5. Rings **21**
5.1. Definition and Basic Properties 21
5.2. Examples of Rings 21
5.3. Quotient Rings and the Isomorphism Theorems for Rings 21
5.4. Integral Domains 21
5.5. Characteristic of a Ring 21
5.6. GCD Domains, PIDs, and Euclidean Domains 22
5.7. UFDs 22

6. Galois Theory **23**

7. Finite Fields **25**

A. Appendix **27**
A.1. Construction of the Complex Numbers 27
A.2. The Euclidean Division Algorithm 27
A.3. Euclid's Lemma 27
A.4. Bezout's Identity 27
A.5. Permutations of a Finite Set 27

Preface

1. Introduction

This text introduces a mathematical subject called *abstract algebra*.

Before we begin, I would like to thank Alex Tsun for his suggestions and feedback on parts of this material. Also, the inclusion of the first chapter is heavily influenced by Charles C. Pinter's *A Book of Abstract Algebra*, which is how I first learned this subject.

2. Prerequisites and Notation

In general, we assume a prerequisite of precalculus at the level of Art of Problem Solving's *Precalculus*. In particular, we assume familiarity with algebra (polynomials, functions, etc.), linear algebra (vectors, matrices, etc.), and complex numbers. We always assume comfortability with proofs and mathematical rigor. It is helpful to know some elementary discrete math (e.g., number theory and combinatorics) as well. We use axiomatic set theory implicitly; if you do not know about this, then it doesn't matter, but if you do know about this, and if you care about equivalence to or dependence on the Axiom of Choice, then we will make skippable notes of that for the relevant theorems. We include an appendix where we cover some of the relevant prerequisite background.

We assume familiarity with standard and common mathematical notation. For example:

1. $\mathbb{R}$ denotes the set of real numbers, $\mathbb{Z}$ the set of integers, and $\mathbb{Q}$ the set of

rational numbers.

2. For either of these three sets, attaching the subscript $+$ produces the subset of positive members, so for example $\mathbb{Z}_+$ is the set of positive integers.

3. Similarly, for either of these three sets, attaching the subscript 0 produces the subset of nonnegative members, so for example $\mathbb{R}_0$ is the set of nonnegative real numbers.

4. For any two sets A and B, $A \times B$ denotes the set of ordered pairs where the first "coordinate" is from A and the second is from B: $A \times B = \{(a, b) : a \in A, b \in B\}$. For example, if $A = \{1, 2\}$ and $B = \{1, 2, 3\}$, then $A \times B = \{(1, 2), (1, 3), (2, 2), (2, 3)\}$.

5. In particular, for any set S, $S^2 = S \times S$ denotes the set of ordered pairs of elements of S: $S^2 = \{(a, b) : a, b \in S\}$. For example, if $S = \mathbb{R}$, then $S^2 = \mathbb{R}^2$ is the set of ordered pairs of real numbers (which geometrically is the set of two-dimensional points, aka two-dimensional space.)

6. For any three sets A, B, and C, $A \times B \times C$ denotes the set of ordered triples where the first "coordinate" is from A, the second from B, and the third from C: $A \times B \times C = \{(a, b, c) : a \in A, b \in B, c \in C\}$.

7. In particular, for any set S, $S^3 = S \times S \times S$ denotes the set of ordered triples of elements of S: $S^3 = \{(a, b, c) : a, b, c \in S\}$. For example, if $S = \mathbb{R}$, then $S^3 = \mathbb{R}^3$ is the set of ordered triples of real numbers (which geometrically is the set of three-dimensional points, aka three-dimensional space.)

8. For any n sets $A_1, ..., A_n$, $A_1 \times ... \times A_n$ denotes the set of all ordered n-tuples where the ith "coordinate" is from A_i: $A_1 \times ... \times A_n = \{(a_1, ..., a_n) : a_i \in A_i\}$.

9. In particular, for any set S, S^n, or S $\times$'d with itself n times, is the set of ordered n-tuples of elements of S: $S^n = \{(a_1, ..., a_n) : a_i \in S\}$.

10. For $x \in A_1 \times ... \times A_n$, x_i denotes the ith "coordinate" of x, which would be an element of A_i. For example, if $x = (4, 5, 6) \in \mathbb{R}^3$, then $x_1 = 4$, $x_2 = 5$,

and $x_3 = 6$.

3. Organization and Order of Topics

To provide context for what follows: the main new concept in abstract algebra is an *algebraic structure*, which we will define and elaborate on in the first chapter. In fact, abstract algebra can be *defined* to be the study of algebraic structures. There are many examples of algebraic structures, and there are four main ones covered in a typical intro to abstract algebra course: groups, rings, fields, and vector spaces.

Many (if not most) abstract algebra curricula start with something other than fields; however, here we proceed in order of fields, vector spaces, groups, and rings. We think that the order should not affect the learning experience too strongly, and fields may be a more "familiar" algebraic structure to start with because of experience with rational, real, and complex numbers.

This text is divided naturally into three parts:

1. Chapters 2 and 3 cover the first two major algebraic structures, fields and vector spaces. This already provides sufficient background for some applications of abstract algebra, such as vector space analysis (which has its own major applications such as quantum mechanics.)

2. Chapters 4 and 5 cover the remaining major algebraic structures, groups and rings, that are part of a typical intro to abstract algebra course.

3. Chapters 6 and 7 cover further applications of and results on these four algebraic structures, completing the list of topics that typically make up an intro course.

The "last chapter" is an appendix, which as mentioned previously covers some relevant prerequisite background.

4. Practice Problems

If you want more practice problems in abstract algebra, we recommend that, in addition to standard "plug-and-chug" exercises that may involve just applying formulas in obvious ways, you also look at "brainteaser" problems that involve more creative combinations of theorems. The Putnam Mathematical Competition is a famous college-level extracurricular contest that features these kinds of "brainteaser" problems for typical undergraduate math subjects, including abstract algebra.

5. Final Preliminaries

I would appreciate any comments, feedback, or corrections. You can contact me at `https://riemannzeta5.com/contact`.

Now, without further ado, let's begin!

1. Primer: What is Abstract Algebra?

For this chapter, we only assume familiarity with real numbers, variables, and equations.

Let us do a brief lookback at our previous studies in algebra, to motivate and provide perspective for what is to come.

When we first learned math, we saw how to compute expressions with numbers, such as $2 + 3 = 5$. Then, when we learned algebra, we shifted to representing numbers with letters. This led to the concept of solving an equation for an unknown variable. For example, we would solve $2 + x = 5$ to get $x = 3$. We would then devote our attention to solving all types of equations – linear, quadratic, polynomial, etc. All the quantities involved in these equations were generally real numbers.

But we can in fact do "algebra" with other types of objects. For example, consider the set of all colors, and say we have an operation on colors, denoted by $+$, which is the result of mixing two colors. So yellow+blue = green, white+black = grey, and so on. We can now do "algebra with colors!" We can write down general properties – for instance, since mixing a color with itself produces just that color, we have, for any color x,

$$x + x = x.$$

Also, it does not matter which order we mix the colors (we define the color mixing operation so that proportions are always the same):

$$a + b = b + a.$$

1. *Primer: What is Abstract Algebra?*

We can write down (and solve) equations:

$$x + \text{blue} = \text{green} \to x = \text{yellow}.$$

If we compare the general properties satisfied by real numbers and colors, we see that there are some similarities (like $a + b = b + a$) but also some differences (like $x + x = x$.)

When abstract algebra was founded, mathematicians had been studying many other "algebras" like these – in fact, it is estimated that today over 200 algebras have been studied. Every time they wanted to formulate a new algebra, they would model some operations on objects and use those operations to build the main concepts of algebra: expressions, equations, and so forth. Each of these systems was both similar to and different from the others. As more exotic types of objects and operations were studied, the subject of algebra shifted from focusing on the real numbers to embracing the full landscape of all these algebras. Soon after, mathematicians started asking: instead of working in each of these algebras in isolation, could we prove theorems that could apply to multiple algebras at once?

Enter abstract algebra. We can see that many properties of an algebra can be derived from a few "fundamental" properties. For instance, consider the real numbers, which satisfy properties like:

1. $a + b = b + a$ (The *Commutativity of Addition*)
2. $ab = ba$ (The *Commutativity of Multiplication*)
3. $(a + b) + c = a + (b + c)$ (The *Associativity of Addition*)
4. $(ab)c = a(bc)$ (The *Associativity of Multiplication*)
5. The existence of a *neutral element* or *identity* 0 for addition, where $0+r = r + 0 = r$
6. The existence of a similar identity 1 for multiplication, where $1r = r1 = r$
7. The existence, for each element r, of its *negative* or *additive inverse* $-r$, where $r + (-r) = (-r) + r = 0$

8. The existence, for each nonzero element r, of its *reciprocal* or *multiplicative inverse* r^{-1}, where $rr^{-1} = r^{-1}r = 1$ (The "nonzero" is important!)

9. $a(b+c) = ab + ac$ (The *Distributivity of Multiplication Over Addition*)

10. $0r = r0 = 0$

Now, say we were solving the linear equation $ax+b = 0$. Recall that the solution is $x = -b/a = -(ba^{-1})$ if $a \neq 0$. How did we arrive at that solution? Let us lay it out in terms of very small, basic steps:

$ax+b$	$=$	0	Start
$(ax+b)+(-b)$	$=$	$0+(-b)$	Add $-b$ to Both Sides
$(ax+b)+(-b)$	$=$	$-b$	$0+r = r+0 = r$ for all r
$ax+(b+(-b))$	$=$	$-b$	Associativity of Addition
$ax+0$	$=$	$-b$	$r+(-r) = (-r)+r = 0$ for all r
ax	$=$	$-b$	$0+r = r+0 = r$ for all r
$a^{-1}(ax)$	$=$	$a^{-1}(-b)$	Multiply Both Sides by a^{-1}, $a \neq 0$
$(a^{-1}a)x$	$=$	$a^{-1}(-b)$	Associativity of Multiplication
$1x$	$=$	$a^{-1}(-b)$	$rr^{-1} = r^{-1}r = 1$ for all r
x	$=$	$a^{-1}(-b)$	$1r = r1 = r$ for all r
x	$=$	$(-b)a^{-1}$	Commutativity of Multiplication
$x+ba^{-1}$	$=$	$(-b)a^{-1}+ba^{-1}$	Add ba^{-1} to Both Sides
$x+ba^{-1}$	$=$	$((-b)+b)a^{-1}$	Distributivity
$x+ba^{-1}$	$=$	$0a^{-1}$	$r+(-r) = (-r)+r = 0$ for all r
$x+ba^{-1}$	$=$	0	$0r = r0 = 0$ for all r
$(x+ba^{-1})+(-ba^{-1})$	$=$	$0+(-ba^{-1})$	Add $-ba^{-1}$ to Both Sides
$x+(ba^{-1}+(-ba^{-1}))$	$=$	$0+(-ba^{-1})$	Associativity of Addition
$x+0$	$=$	$0+(-ba^{-1})$	$r+(-r) = (-r)+r = 0$ for all r
x	$=$	$-(ba^{-1})$	$0+r = r+0 = r$ for all r

Now, the fundamental idea that enables abstract algebra is this: we do not actually need a, x, or b to be real numbers for this process to work. All we need

are those properties to be satisfied!

In other words, if we assume that we generally had some set of mathematical objects – they might be real numbers, or they might not be – together with some operations on those objects, that satisfied the properties above, then we can solve linear equations $ax + b = 0$ for those objects, the same way we solve linear equations for real numbers. In fact, anything we do in this general setting can immediately carry over to any algebra that shares these properties with the real numbers. We *abstract* the fact that the objects and operations satisfy the properties away from what the objects or operations actually are, hence the term, *abstract algebra*.

To give an example of this process, consider the generalization of the algebra of real numbers to this concept:

Definition 1.0.1. A *field* is a set F together with operations $+, \times$ on the elements of F. These operations must be *binary*: they both must take in two elements and output one element. They must also satisfy the following properties:

1. *Commutativity*: For all a, b in F, $a + b = b + a$ and $a \times b = b \times a$
2. *Associativity*: For all a, b, c in F, $(a+b)+c = a+(b+c)$ and $(a \times b) \times c = a \times (b \times c)$
3. *Additive and multiplicative identity*: There exist *distinct* elements of F called 0 and 1, such that, for all a in F, $0 + a = a + 0 = a$ and $1 \times a = a \times 1 = a$
4. *Additive inverse*: For all a in F, there exists an element $-a$ such that $a + (-a) = (-a) + a = 0$
5. *Multiplicative inverse*: For all *nonzero* a in F, there exists an element a^{-1} such that $a \times a^{-1} = a^{-1} \times a = 1$
6. *Distributivity*: For all a, b, c in F, $(a + b) \times c = (a \times c) + (b \times c)$ and $a \times (b + c) = (a \times b) + (a \times c)$

From these defining properties of a field, we can actually prove $0x = x0 = 0$ as a consequence. Thus, linear equation solutions for general fields are the same as linear equation solutions for real numbers. We can go on to ask how much real number algebra can carry over – and indeed we can discuss polynomials over fields and theorems for fields that heavily generalize theorems about real numbers.

We can then identify many sets of objects which these theorems can apply to. For example, consider the set of just 0 and 1 with the following "addition" and "multiplication" operations:

$$\begin{aligned}
0+0 &= 0, \\
0+1 &= 1, \\
1+0 &= 1, \\
1+1 &= 0, \\
0\times 0 &= 0, \\
0\times 1 &= 0, \\
1\times 0 &= 0, \\
1\times 1 &= 1.
\end{aligned}$$

This is actually a field! (It is called $\mathbb{Z}_2$.) In fact, this is a special case of a more general kind of construction. Consider say the set of numbers from 0 to 9, with the operations of usual addition and multiplication but "wrapped around" to stay in 0 through 9 (so for example $9 + 9 = 18 \to 18 - 10 = 8$.) In other words, the operations are the remainders when divided by 10 of usual addition and multiplication. We can do this for any number $n > 1$: we define $\mathbb{Z}_n$ to be the set of numbers from 0 to $n - 1$ (including both) with the operations as remainders when divided by n of usual addition and multiplication. Then, it turns out that if n is prime, $\mathbb{Z}_n$ is a field! We will prove this in general later on. In fact, these "finite fields" turn out to have lots of interesting applications, including being foundational to modern computer security.

We can alternatively study a different set of defining properties. For example, another concept in abstract algebra is a *group*:

Definition 1.0.2. A *group* is a set G together with a binary operation $\times$ satisfying:

1. *Associativity*: For all a, b, c in G, $(a \times b) \times c = a \times (b \times c)$
2. *Identity*: There exists an element of G, which we usually call e, such that, for all g in G, $g \times e = e \times g = g$
3. *Inverse*: For all g in G, there exists an element g^{-1} such that $g \times g^{-1} = g^{-1} \times g = e$

This is just a subset of the defining properties of fields. This automatically means that all fields are groups[1], but so are many other algebras that aren't fields – and all of group theory applies to all of these cases. As an example, it turns out that the set of transformations you can apply to a Rubik's cube is a group – in fact, group theory is an important tool for discovering algorithms to solve Rubik's cubes quickly[2].

These abstracted concepts – fields and groups – are called *algebraic structures*. An algebraic structure is essentially any defining list of properties that a set of operations are required to satisfy.

The strength of abstract algebra is many-fold. First, theorems on algebraic structures extend results on familiar algebras like the real numbers to other sets of objects. But in the other direction, studying abstract algebra has actually resulted in new theorems even for familiar algebras like the real numbers! We will see more examples of this phenomenon later on.

[1] Did we mean "all groups are fields"? It may seem wonky, but think about it for a moment and you'll see that it's actually the other way around. A smaller set of defining properties means fewer requirements, which means more possibilities, so groups are more general than fields, not less. Groups include fields, so all fields are groups, but groups include other algebras too.

[2] The application of group theory to Rubik's cubes is not discussed in this text, but if you're interested you should look into it afterwards!

2. Fields

2.1. Definition and Basic Properties

The first algebraic structure we study is the field. Fields abstract the properties of real numbers, as we just saw in the previous chapter:

Definition 2.1.1. A *field* is a set F together with operations $+, \times$ on the elements of F. These operations must be *binary*: they both must take in two elements and output one element. They must also satisfy the following properties:

1. *Commutativity*: For all a, b in F, $a + b = b + a$ and $a \times b = b \times a$
2. *Associativity*: For all a, b, c in F, $(a+b)+c = a+(b+c)$ and $(a \times b) \times c = a \times (b \times c)$
3. *Additive and multiplicative identity*: There exist *distinct* elements of F called 0 and 1, such that, for all a in F, $0 + a = a + 0 = a$ and $1 \times a = a \times 1 = a$
4. *Additive inverse*: For all a in F, there exists an element $-a$ such that $a + (-a) = (-a) + a = 0$
5. *Multiplicative inverse*: For all *nonzero* a in F, there exists an element a^{-1} such that $a \times a^{-1} = a^{-1} \times a = 1$
6. *Distributivity*: For all a, b, c in F, $(a + b) \times c = (a \times c) + (b \times c)$ and $a \times (b + c) = (a \times b) + (a \times c)$

For $a, b \in F$, we write $ab = a \times b$, just like for numbers. We also adopt the standard order of operations conventions when writing expressions, so that for

example $a + bc = a + (b \times c)$.

Note: in abstract algebra we typically use the same symbol to refer to both a set and a set with some operations we have defined on it. The usage is dependent on context.

To remember the terminology of commutativity and associativity, we can visualize them as follows: commutativity ($a+b = b+a$) looks like "a and b commuting to each others' positions," while associativity looks like "b associating first with a and then with c."

Before we dive further into the theory of fields, we mention some examples. Our "prototypal" one, which we based the very definition of a field off of, is $\mathbb{R}$, the set of real numbers, with the usual addition and multiplication. But there are others too: as discussed previously, $\mathbb{Z}_2$ is a field. (As a reminder, we define $\mathbb{Z}_n$, for any positive integer n, to be $\{0, 1, ..., n-1\}$ with addition and multiplication modulo n.) Additionally, $\mathbb{Z}_3$ is also a field. (If you're unsure, check all the properties explicitly to see that these are in fact fields.)

Now, often times, we'd like to consider properties that hold for specific algebras and ask whether they generalize to the abstract setting – that is a large part of the motivation of abstract algebra. But we also want to balance this aspiration with some skepticism: not all properties will carry over to the general case. Here, to gain some intuition for what works, we can take our inspiration from $\mathbb{R}$, which has very rich algebraic properties, but if we are ever in doubt, we can verify whether the property in question is true for something like $\mathbb{Z}_2$ or $\mathbb{Z}_3$. These "look very different" from $\mathbb{R}$ but they are still fields. If the property fails for these cases, then it can't carry over to general fields.

So let's start now by looking at the properties of $\mathbb{R}$. As mentioned earlier:

Theorem 2.1.2. We have $x0 = 0x = 0$.

Think for a moment about how we may prove this from the defining field properties. The only thing we know about 0 is that it is an additive identity, and our desired result involves multiplication. There's actually only one field property that relates addition and multiplication to each other, and that is distributivity.

(Notice that all the other properties talk about either addition or multiplication, not both.) Thus, we expect that we will need to involve $a + 0 = 0 + a = a$ and distributivity at some point.

Let's first just try to show that $x0 = 0$. We have $0 + a = a$; since we want $x0$ to appear at some point, it seems like a good idea to multiply both sides on the left by x. Distributivity further helps our hunch, since $x(0 + a) = x0 + xa$, so we've made our desired term appear. This yields $x0 + xa = xa$. Now, we'd like to cancel xa from both sides like we do with numbers; then, we'd have our desired result. But remember, these are field elements, not numbers. Is cancellation still possible? Yes, it is: that's exactly what the additive inverse allows us to do. Putting this together, we have our proof:

Proof. We have $0 + a = a$. Multiplying both sides by x, $x(0 + a) = xa$. By distributivity, $x(0 + a) = x0 + xa$, so $x0 + xa = xa$. Now, xa has an additive inverse $-(xa)$, so $(x0 + xa) + (-(xa)) = xa + (-(xa))$. The RHS is 0, and we can use associativity of addition on the LHS: $(x0 + xa) + (-(xa)) = x0 + (xa + (-(xa)))$. But this is just $x0 + 0 = x0$. This yields $x0 = 0$, as desired.

See if you can come up with the proof that $0x = 0$. □

An important point that results from this is the fact that, in any field, 0 can't have a multiplicative inverse. How can we prove this? Well, say it did have an inverse, call it 0^{-1}, then we'd have $00^{-1} = 1$, but also what we've just shown implies that $00^{-1} = 0$, and 0 and 1 are required to be different, which is a contradiction. This is why the *nonzero* part in the multiplicative inverse property is necessary, and consequently why there is some asymmetry in the additive and multiplicative inverse properties.

However, just as we were able to cancel added terms by using additive inverses, we can cancel *nonzero* multiplied terms by using multiplicative inverses *of nonzero elements*. (Write this out and prove it if you're unsure.) So while we can't cancel out 0 in multiplication, we can cancel out anything else. This is familiar from real-number algebra, and it holds true for general fields too.

Now, another point that we haven't yet elaborated on is that here, 0 and 1 are identities (for addition and multiplication respectively), but *not necessarily the only identities*. If we don't have uniqueness, then we can't really speak of "the" additive or multiplicative identity. Will this be an issue when trying to generalize more of real number algebra to fields? It turns out that we don't need to worry:

Theorem 2.1.3. The additive and multiplicative identities are unique.

Again, think for a moment about how you would prove this. Consider addition first. Another way of saying this is that if we had two additive identities, say 0_1 and 0_2, then both of them would have to be equal. How do we show this? Well, we want to involve 0_1, 0_2, and addition at some point. It seems like a good idea to consider $0_1 + 0_2$:

Proof. If we had two additive identities 0_1 and 0_2, then consider $0_1 + 0_2$. On the one hand, $0_1 + 0_2 = 0_1$ since 0_2 is an identity; on the other hand, $0_1 + 0_2 = 0_2$ since 0_1 is an identity. Thus, $0_1 = 0_2$, and the desired result follows.

See if you can come up with the proof that multiplicative identities are unique. $\square$

Notice that we didn't actually use any other field properties here: we just used the fact that 0_1 and 0_2 are identities. This shows us that even beyond fields, whenever we have identities in abstract algebra, they will be unique. This will apply later on when we discuss for example group and ring theory.

Let's go back to $\mathbb{R}$. A common way of think of subtraction is as the "inverse" of addition, and similarly of division as the "inverse" of multiplication. Indeed, we can say for numbers that $a - b = a + (-b)$ and $a/b = ab^{-1}$. But we have inverses for any field! Does that mean we can extend subtraction and division to general fields as well?

The main issue in doing that stems again from uniqueness: on a field, we're given that $-b$ is *an* additive inverse for b, not necessarily *the*. So if we wanted to say subtract b from a, via $a - b = a + (-b)$, which value of $-b$ would we pick?

Furthermore, different choices of $-b$ would necessarily yield different values of $a+(-b)$, because of cancellation. (Write this out and explicitly prove it if you're unsure.) However, it turns out that we don't need to worry here either:

Theorem 2.1.4. Additive and multiplicative inverses are unique. (For $x \in F$, there is exactly one element $-x$ that adds with x to 0, and similarly, if $x \neq 0$, exactly one element x^{-1} that multiplies with x to 1.)

As before, think about how we could prove this. Start with addition. A safe bet would be to replicate part of our previous method for uniqueness of identities, by assuming there were two inverses and showing they must be equal. So take $x \in F$, and say that y and z are both additive inverses of x. Then, we'd have $x + y = 0$ and $x + z = 0$. Now, try to complete the proof from here. And similarly, try to establish uniqueness of multiplicative inverses.

Uniqueness of inverses now allows subtraction $a - b = a + (-b)$ and division $a/b = ab^{-1}$, which we will henceforth use throughout regardless of whether we are on $\mathbb{R}$ or a general field.

Let's look at some other properties that can carry over from $\mathbb{R}$; in particular, focus on consequences of the associative property. Take addition first. We know that addition is only required to be a binary operation (takes in two elements), so if we have an expression like $a+b+c$, what does that mean? What order do we evaluate things in? Is it $(a+b)+c$ or $a+(b+c)$? Well, associativity implies in this case that it doesn't matter, since these have to be the same. What about $a+b+c+d$ – is it $((a+b)+c)+d$, or $(a+b)+(c+d)$, or $a+((b+c)+d)$, or what?

Well, we can just repeatedly apply associativity to get that these are all equivalent too! For example, $((a+b)+c)+d = (a+(b+c))+d = a+((b+c)+d) = a+(b+(c+d))$. In a similar vein, with repeated application we can show that associativity implies that we never need parentheses when adding elements: all the possible orders of evaluation will be the same. (Actually proving this is a bit tricky – try establishing it formally! As a hint, induction helps a lot for this sort of argument.) Thus, we don't need to worry about writing out expressions like $a_1 + a_2 + ... + a_n$ when we have associativity. And the same is true for

multiplication, since that is associative as well.

Now, focus on consequences of the commutative property. Again, take addition first. Off the bat, this tells us that it doesn't matter which order we add two elements in: whether we put a or b first, we have $a + b = b + a$. But we just saw that we can add as many elements as we want, without specifying parentheses. So it's natural to ask: if we re-order other numbers of elements and then add them, would the result be the same? For example, do we have that $a + b + c = c + b + a = a + c + b = b + a + c$? (Remember that these are field elements, not necessarily numbers.)

Well, we have that $a+b+c = a+(b+c) = a+(c+b) = (c+b)+a = c+b+a$. Similarly, by repeated application of commutativity and associativity, we can show that in general we can reorder as many elements as we'd like without changing the result. (Again, try to establish this formally; as before, induction can help.)

These two consequences yield that a lot of the algebra we do for real numbers carries over to general fields too. For example, just like in real-number algebra, we can collect like terms: we have $a+b+a+c+b = a+a+b+b+c$, $abacb = aabbc$, and so on. Furthermore, just like for numbers, we can take multiples and powers of general field elements: we can write down something like $10x + 3y$ or $x^{10}y^3$ and know that this is equal to any re-ordering of 10 x's and 3 y's added (or multiplied) together, in any order of evaluation. Thus, for any $x \in F$ and any positive integer n, it makes sense to write nx and x^n.

A natural question then is: what about negative multiples and powers? Let's focus on the multiples first. If an integer is negative, so of the form $-n$ for n a positive integer, then there are multiple "reasonable" ways we could define $(-n)x$: it could be $-(nx)$ (the additive inverse of nx), or $n(-x)$ (the nth multiple of $-x$.) Which of these should we pick? Hopefully, it would be nice if these turned to be equal to each other, then we wouldn't have to pick between two "reasonable-sounding" choices. Also, it seems that we've had some luck so far with such "reasonable" equivalencies holding true for general fields, so maybe the same would be true here as well.

So let's try to prove that they must be equal. Take $n = 2$ as a "guinea pig." We want to show that $(-x)+(-x) = -(x+x)$. Well, rearranging, this is equivalent to $(-x) + (-x) + x + x = 0$, and since we can reorder terms, this is the same as $(-x) + x + (-x) + x = 0$. But we know that $-x + x = 0$, so we then have $(-x) + x + (-x) + x = 0 + 0 = 0$, as desired. Similarly, we can show in general that $-(nx) = n(-x)$. (As an exercise, prove this formally.)

In a similar way, for multiplication, we have for any positive integer n that $(x^{-1})^n = (x^n)^{-1}$ (again, as an exercise, prove this formally), so we can define x^{-n} unambiguously to be either of these two.

If we have positive integer and negative integer multiples/powers, what about the zero multiple/power? Again, look at multiples first. What would $0x$ be? We have to be careful: we already have an element of the field that we're calling 0 as well. To avoid confusion, let's label these $0_{\mathbb{Z}}$ and 0_F. We know that $0_F x = 0_F$, and we are asking what $0_{\mathbb{Z}} x$ should be – i.e., what to define it as. We can get rid of the possible confusion around 0 by just defining these to be equal: we define $0_{\mathbb{Z}} x = 0_F$ for all x. Then, we can unambiguously write $0x = 0$.

What about the zeroth power? Looking at numbers, since $x^0 = 1$ for any nonzero real number x, we can analogously define $x^0 = 1$ for any nonzero x on a field.

So now that we have multiples and powers, do standard results about multiples and powers of numbers carry over to fields? For example, is it true that for integers m, n and field elements x, y, we have $mx + nx = (m + n)x$, or $x^{m+n} = x^m x^n$? As an exercise, prove those two as well as the following also-familiar properties:

$$(mn)x = m(nx),$$

$$x^{mn} = (x^m)^n,$$

$$(nx)y = n(xy).$$

Remember the impact of our theory: though it looks familiar since we've already seen it for numbers, these results apply to *any* field. That includes many fields

that look different from $\mathbb{R}$, like $\mathbb{Z}_2$, $\mathbb{Z}_3$, and many others that we will encounter later on.

What are some other things that we typically do with numbers? Well, one thing is that we often expand products of sums, like

$$(a+b)(c+d) = ac + ad + bc + bd,$$
$$(a+b)(c+d+f) = ac + ad + af + bc + bd + bf,$$

and so on. Are these true for fields as well?

Well, how did we derive them for numbers? It was really just repeatedly applying distributivity! For example, we have

$$\begin{aligned}(a+b)(c+d) &= a(c+d) + b(c+d) \\ &= (ac+ad) + (bc+bd) \\ &= ac + ad + bc + bd.\end{aligned}$$

Well, mainly distributivity, but in conjunction with some other properties like associativity. But these properties hold for fields too (by definition), so these expansions must hold as well for fields. In general, we have that if we multiply two sums A and B, then the result will be the product of each term in A with each term in B, all added together:

$$\sum_{i=1}^{m} a_i \sum_{j=1}^{n} b_j = \sum_{1 \le i \le m, 1 \le j \le n} a_i b_j.$$

(If you are unsure, prove this formally.)

We can obtain similar expansions for multiplying three, or four, or any number of sums together, on a general field.

A particular consequence of all this is the *binomial theorem*, which hopefully is familiar to you from $\mathbb{R}$. As a recap, this is a result that specifies how to expand $(a+b)^2$, $(a+b)^3$, and so on – in general, $(a+b)^n$. It states that

$$(a+b)^n = \sum_{i=0}^{n} \binom{n}{i} a^i b^{n-i}.$$

By what we have just discussed, this must hold generally on all fields.

Remember again that our work here is applicable beyond $\mathbb{R}$. As an example, we know that $\mathbb{Z}_3$ is a field; thus, the binomial theorem must hold on $\mathbb{Z}_3$. Furthermore, each binomial coefficient $\binom{n}{i}$ is divisible by n except for $i = 0$ and $i = n$. Thus, if n is a multiple of 3, then on $\mathbb{Z}_3$, we have that each term corresponding to i other than 0 or n vanishes ($3x = 0$ for all $x \in \mathbb{Z}_3$), yielding just

$$(a+b)^n = a^n + b^n.$$

This is a very "simple" and "nice" expansion that holds on $\mathbb{Z}_3$, but not on say $\mathbb{R}$. (This is sometimes jokingly called the "freshman's dream.")

Constantly keep in mind the generality and power of the work we are doing here, especially as we progress and continue to take intuition from familiar algebras like $\mathbb{R}$.

2.2. Polynomials Over Fields

Section to be added.

2.3. Isomorphisms and Homomorphisms

Section to be added.

2.4. Subfields

Section to be added.

2.5. Field Extensions

Section to be added.

2.6. Construction of a Simple Algebraic Extension

Section to be added.

2.7. Construction of Any Algebraic Extension

Section to be added.

3. Vector Spaces

3.1. Definition and Basic Properties

Section to be added.

3.2. Finite-Dimensional Vector Spaces

Section to be added.

3.3. Matrices and Linear Transformations

Section to be added.

3.4. Matrices and Subspaces

Section to be added.

3.5. General Bases

Section to be added.

3.6. Field Extensions as Vector Spaces

Section to be added.

4. Groups

4.1. Definition and Basic Properties

Section to be added.

4.2. Permutation and Symmetry Groups

Section to be added.

4.3. Cyclic Groups

Section to be added.

4.4. Order of an Element

Section to be added.

4.5. Generators, Presentations, and Free Groups

Section to be added.

4.6. Quotient Groups

Section to be added.

4.7. The Isomorphism Theorems

Section to be added.

4.8. Group Actions

Section to be added.

5. Rings

5.1. Definition and Basic Properties

Section to be added.

5.2. Examples of Rings

Section to be added.

5.3. Quotient Rings and the Isomorphism Theorems for Rings

Section to be added.

5.4. Integral Domains

Section to be added.

5.5. Characteristic of a Ring

Section to be added.

5.6. GCD Domains, PIDs, and Euclidean Domains

Section to be added.

5.7. UFDs

Section to be added.

6. Galois Theory

Chapter to be added.

7. Finite Fields

Chapter to be added.

A. Appendix

A.1. Construction of the Complex Numbers

Section to be added.

A.2. The Euclidean Division Algorithm

Section to be added.

A.3. Euclid's Lemma

Section to be added.

A.4. Bezout's Identity

Section to be added.

A.5. Permutations of a Finite Set

Section to be added.