# Rate

### Riesa Y. Cassano-Coleman

#### 2024-10-31

```
library(tidyverse) # version 2.0.0
library(magrittr) # version 2.0.3
library(nparLD) # version 2.2
select <- dplyr::select # make sure we're using tidyverse's version of select...

Load the data.
data <- read_csv('../data/response_rate_by_sub.csv', show_col_types = FALSE)


Change musician into a factor.
data %<>% mutate(Musician = factor(Musician, levels = c('Yes', 'No')))

Make sure Intact naming is consistent.
data %<>% mutate(scramble = ifelse(scramble == 'intact', 'Intact', scramble))
```

#### Visualize!

(Generate figure 3, minus significance markers)

```
data %>%
  ggplot(aes(x = scramble, y = mean_response_rate, fill = Musician)) +
  geom_bar(position = "dodge", stat = "summary", fun = mean) +
  geom_errorbar(position = position_dodge(width = 0.9), width = 0.2, stat = "summary") +
  geom_point(position = position_jitterdodge(jitter.width = 0.1), alpha = 0.1) +
  theme_gray(base_size = 16) +
  xlab('Scramble Level') +
  ylab('Response Rate (per Minute)') +
  scale_fill_discrete(name="", labels=c('Musicians', 'Non-musicians')) +
  theme(legend.text = element_text(size = 12))
```



#ggsave('rate.png', width = 7, height = 5)

#### Non-parametric ANOVA-type test

Is there an overall effect of scramble condition or musical experience?


```
## FILD FI Model
## -----
## Check that the order of the time and group levels are correct.
```

## Time level: 1B 2B 8B Intact

## Group level: Yes No

## If the order is not correct, specify the correct order in time.order or group.order.

## **Relative Effects**



# condition

```
## expertise 1.406889 1.000000 2.355736e-01
## condition 44.553463 2.311547 8.619521e-23
## expertise:condition 1.544214 2.311547 2.099931e-01
```

- Main effect of scramble condition: F(2.3) = 44.553, p < .001
- No main effect of musical experience: F(1) = 1.407, p = .236
- (Interaction is not significant: F(2.3) = 1.544, p = .210)

#### Post-hoc tests between conditions

What conditions are different from each other?

```
pairwise.wilcox.test(data$mean_response_rate,
                     data$scramble,
                     p.adjust.method = "bonferroni", paired=TRUE)
##
##
   Pairwise comparisons using Wilcoxon signed rank test with continuity correction
##
          data$mean_response_rate and data$scramble
## data:
##
##
          1B
                  2B
                          8B
## 2B
## 8B
          4.7e-11 7.1e-10 -
## Intact 2.3e-08 1.2e-07 1
##
## P value adjustment method: bonferroni
```

Participants responded more frequently in 1B and 2B compared to 8B and Intact (p < .001 all comparisons). There is no difference between 1B and 2B (p = 1.0) or between 8B and Intact (p = 1.00).