

De Giorgi-Nash-Moser Theory

89hao

January 8, 2020

Abstract

This is a learning note about the De Giorgi-Nash-Moser theory, the reference book is Qing Han and Fanghua Lin's *Elliptic Partial Differential Equations*

The main task of this note is to prove the following theorem:

Theorem 1 Suppose $a_{ij} \in L^\infty(B_1)$ and $c \in L^q(B_1)$ for some $q > \frac{n}{2}$ satisfy the following assumptions

$$a_{ij}(x)\xi_i\xi_j \geq \lambda|\xi|^2 \text{ for any } x \in B_1 \text{ and } |a_{ij}|_{L^\infty} + \|c\|_{L^q} \leq \Lambda$$

for some positive constants λ and Λ . Suppose that $u \in H^1(B_1)$ is a subsolution in the following sense

$$\int_{B_1} a_{ij} D_i u D_j \varphi + c u \varphi \leq \int_{B_1} f \varphi \text{ for any } \varphi \in H_0^1(B_1) \text{ and } \varphi \geq 0 \text{ in } B_1.$$

If $f \in L^q(B_1)$, then $u^+ \in L_{\text{loc}}^\infty(B_1)$. Moreover, there holds for any $\theta \in (0, 1)$ and any $p > 0$

$$\sup_{B_\theta} u^+ \leq C \left\{ \frac{1}{(1-\theta)^{\frac{n}{p}}} \|u^+\|_{L^p(B_1)} + \|f\|_{L^q(B_1)} \right\}$$

where $C = C(n, \lambda, \Lambda, p, q)$ is a positive constant.

We present two methods here.

1 Moser's Method

Think about the following undergraduate level question:

Question Let $f \in C[0, 1]$, then what is the value of

$$\lim_{\gamma \rightarrow \infty} \left| \int_0^1 |f(x)|^\gamma dx \right|^{\frac{1}{\gamma}} = ?$$

The answer is $\sup_{0 \leq x \leq 1} |f(x)| = \|f\|_{L^\infty}$. This is exactly the way we transform $\|u^+\|_{L^p}$ to $\sup u^+$ in the proof of the above theorem. To make things more understanding, we assume $f = 0$. In this simplified version, We first establish the inequality

$$\left(\int_{B_r} |u^+|^{\gamma r} \right)^{\frac{1}{r}} \leq C \int_{B_R} |u^+|^\gamma,$$

where $\chi > 1$ and $\gamma \geq 2$, $r < R$. Then we use this inequality to iterate, the iterating step makes $\chi \rightarrow \infty$, then the left side would be more and more likely to the supremum norm of u^+ just as the question. Hence we can get the following inequality by doing this iteration:

$$\sup_{B_{\frac{1}{2}}} u^+ \leq C \|u^+\|_{L^2(B_1)}.$$

Thus the case $f = 0$ and $p = 2$ can be proved. The proof under the condition of $f \neq 0$ needs to be modified slightly. Then the general case that $p = 2$ can be proved easily by using the above special case.

According to the above discussion, we want to establish the inequality like this:

$$\|u^+\|_{L^{\gamma\chi}(B_r)} \leq C \|u^+\|_{L^\gamma(B_R)}$$

for some constant C and $r < R$. This inequality estimate the $L^{\gamma\chi}$ -norm by the weaker L^γ -norm. As a trade-off, we have to make $r < R$, via certain test function.

For some $k > 0$ and $m > 0$, set $\bar{u} = u^+ + k$ and

$$\bar{u}_m = \begin{cases} \bar{u} & \text{if } u \leq m \\ k + m & \text{if } u \geq m \end{cases}$$

The point is that \bar{u}_m is still an element of $H^1(B_1)$, but bounded below by k and from above by $(k + m)$. Then we have $D\bar{u}_m = 0$ whenever $u < 0$ or $u > m$ and $\bar{u}_m \leq \bar{u}$. Set the test function

$$\phi = \eta^2 (\bar{u}_m^\beta \bar{u} - k^{\beta+1}) \in H_0^1(B_1)$$

for some $\beta \geq 0$ and some nonnegative function $\eta \in C_0^1(B_1)$. The function η is a cut-off function to be chosen later on (remember the trade-off $r < R$?). ϕ is an element of $H_0^1(B_1)$ because \bar{u}_m is bounded. Direct calculation yields

$$\begin{aligned} D\phi &= \beta \eta^2 \bar{u}_m^{\beta-1} D\bar{u}_m \bar{u} + D\bar{u} \eta^2 \bar{u}_m^\beta + 2\eta D\eta (\bar{u}_m^\beta \bar{u} - k^{\beta+1}) \\ &= \eta^2 \bar{u}_m^\beta (\beta D\bar{u}_m + D\bar{u}) + 2\eta D\eta (\bar{u}_m^\beta \bar{u} - k^{\beta+1}). \end{aligned}$$

where we used the fact that $\bar{u} = \bar{u}_m$ whenever $D\bar{u}_m \neq 0$. Then we have

$$\begin{aligned} \int a_{ij} D_i u D_j \phi &= \int a_{ij} D_i \bar{u} (\beta D_j \bar{u}_m + D_j \bar{u}) \eta^2 \bar{u}_m^\beta + 2 \int a_{ij} D_i \bar{u} D_j \eta (\bar{u}_m^\beta \bar{u} - k^{\beta+1}) \\ &\geq \lambda \beta \int \eta^2 \bar{u}_m^\beta |D\bar{u}_m|^2 + \lambda \int \eta^2 \bar{u}_m^\beta |D\bar{u}|^2 - \Lambda \int |D\bar{u}| |D\eta| \bar{u}_m^\beta \bar{u} \eta \\ &\geq \lambda \beta \int \eta^2 \bar{u}_m^\beta |D\bar{u}_m|^2 + \frac{\lambda}{2} \int \eta^2 \bar{u}_m^\beta |D\bar{u}|^2 - \frac{2\Lambda^2}{\lambda} \int |D\eta|^2 \bar{u}_m^\beta \bar{u}^2. \end{aligned}$$

Hence we obtain by noting $\bar{u} \geq k$

$$\begin{aligned} &\beta \int \eta^2 \bar{u}_m^\beta |D\bar{u}_m|^2 + \int \eta^2 \bar{u}_m^\beta |D\bar{u}|^2 \\ &\leq C \left\{ \int |D\eta|^2 \bar{u}_m^\beta \bar{u}^2 + \int (|c| \eta^2 \bar{u}_m^\beta \bar{u}^2 + |f| \eta^2 \bar{u}_m^\beta \bar{u}) \right\} \\ &\leq C \left\{ \int |D\eta|^2 \bar{u}_m^\beta \bar{u}^2 + \int c_0 \eta^2 \bar{u}_m^\beta \bar{u}^2 \right\} \end{aligned}$$

where c_0 is defined as

$$c_0 = |c| + \frac{|f|}{k}.$$

Choose $k = \|f\|_{L^q}$ if f is not identically zero. Otherwise choose arbitrary $k > 0$ and eventually let $k \rightarrow 0^+$. By assumption we have

$$\|c_0\|_{L^q} \leq \Lambda + 1.$$

Set $w = \bar{u}_m^{\frac{\beta}{2}} \bar{u}$. Then

$$|Dw| = \bar{u}_m^{\frac{\beta}{2}} \left(\frac{\beta}{2} \cdot D\bar{u}_m + D\bar{u} \right),$$

therefore

$$\begin{aligned} |Dw|^2 &= \bar{u}_m^\beta \left| \frac{\beta}{2} D\bar{u}_m + D\bar{u} \right|^2 \\ &= \bar{u}_m^\beta \left(\frac{\beta^2}{4} |D\bar{u}_m|^2 + \beta D\bar{u}_m D\bar{u} + |D\bar{u}|^2 \right) \\ &= \bar{u}_m^\beta \left(\beta \left(\frac{\beta}{4} + 1 \right) |D\bar{u}_m|^2 + |D\bar{u}|^2 \right) \\ &\leq \bar{u}_m^\beta (\beta + 1) \left(\beta |D\bar{u}_m|^2 + |D\bar{u}|^2 \right). \end{aligned}$$

Therefore we have

$$\int |Dw|^2 \eta^2 \leq C \left((1 + \beta) \int w^2 |D\eta|^2 + (1 + \beta) \int c_0 w^2 \eta^2 \right)$$

and so

$$\begin{aligned} \int |D(w\eta)|^2 &\leq 2 \int \left(|D\eta|^2 w^2 + |Dw|^2 \eta^2 \right) \\ &\leq C \left((1 + \beta) \int w^2 |D\eta|^2 + (1 + \beta) \int c_0 w^2 \eta^2 \right). \end{aligned}$$

C in different lines may not be identical. Hölder inequality implies

$$\int c_0 w^2 \eta^2 \leq \left(\int c_0^q \right)^{\frac{1}{q}} \left(\int (\eta w)^{\frac{2q}{q-1}} \right)^{1-\frac{1}{q}} \leq (\Lambda + 1) \left(\int (\eta w)^{\frac{2q}{q-1}} \right)^{1-\frac{1}{q}}.$$

By interpolation inequality and Sobolev inequality and Sobolev's inequality with $2^* = \frac{2n}{n-2} > \frac{2q}{q-1} > 2$ if $q > \frac{n}{2}$, we have

$$\begin{aligned} \|\eta w\|_{L^{\frac{2q}{q-1}}} &\leq \varepsilon \|\eta w\|_{L^{2^*}} + C(n, q) \varepsilon^{-\frac{n}{2q-n}} \|\eta w\|_{L^2} \\ &\leq \varepsilon \|D(\eta w)\|_{L^2} + C(n, q) \varepsilon^{-\frac{n}{2q-n}} \|\eta w\|_{L^2} \end{aligned}$$

for any small $\varepsilon > 0$. (What we do in this inequality is to split c_0 and $w^2 \eta^2$. If $c = f = 0$, then the operation here would not be needed and the proof can be simpler.) Therefore we obtain

$$\int |D(w\eta)|^2 \leq C \left((1 + \beta) \int w^2 |D\eta|^2 + (1 + \beta)^{\frac{2q}{2q-n}} \int w^2 \eta^2 \right)$$

and in particular

$$\int |D(w\eta)|^2 \leq C(1+\beta)^\alpha \int (|D\eta|^2 + \eta^2) w^2$$

where α is a positive number depending only on n and q . From the Sobolev inequality, with $\chi = n/(n-2) > 1$ for $n > 2$ and any fixed $\chi > 2$ for $n = 2$, we get

$$\left(\int |\eta w|^{2\chi} \right)^{\frac{1}{\chi}} \leq C(1+\beta)^\alpha \int (|D\eta|^2 + \eta^2) w^2.$$

Choose the cut-off function η as follows. For any $0 < r < R \leq 1$ set $\eta \in C_0^1(B_R)$ with the property

$$\eta \equiv 1 \text{ in } B_r \text{ and } |D\eta| \leq \frac{2}{R-r}.$$

Then we obtain

$$\left(\int_{B_r} w^{2\chi} \right)^{\frac{1}{\chi}} \leq \frac{(1+\beta)^\alpha}{(R-r)^2} \int_{B_R} w^2.$$

Since by definition of $w = \bar{u}_m^\beta \bar{u}$, we have

$$\left(\int_{B_r} \bar{u}^{2\chi} \bar{u}_m^{2\beta\chi} \right)^{\frac{1}{\chi}} \leq C \frac{(1+\beta)^\alpha}{(R-r)^2} \int_{B_R} \bar{u}^2 \bar{u}_m^{2\beta}.$$

Set $\gamma = \beta + 2 \geq 2$. Then we obtain

$$\left(\int_{B_r} \bar{u}_m^{2\gamma} \right)^{\frac{1}{\gamma}} \leq C \frac{(\gamma-1)^\alpha}{(R-r)^2} \int_{B_R} \bar{u}^\gamma$$

provided the integral in the right-hand side is bounded. By letting $m \rightarrow \infty$ we obtain

$$\|\bar{u}\|_{L^{\gamma\chi}(B_r)} \leq \left(C \frac{(\gamma-1)^\alpha}{(R-r)^2} \right)^{\frac{1}{\gamma}} \|\bar{u}\|_{L^\gamma(B_R)}$$

provided $\|\bar{u}\|_{L^\gamma(B_R)} < \infty$, where $C = C(n, q, \lambda, \Lambda)$ is a positive constant independent of γ .

Then we do the iteration, taking successively the values $\gamma = 2, 2\chi, 2\chi^2, \dots$. Define, for all $i = 1, 2, \dots$,

$$\gamma_i = 2\chi^i \text{ and } r_i = 2 + \frac{1}{2^{i-1}}.$$

For any $i \geq 0$, $\gamma_{i+1} = \chi\gamma_i$, $r_i - r_{i+1} = \frac{1}{2^{i+2}}$, we have

$$\|\bar{u}\|_{L^{\gamma_{i+1}}(B_{r_{i+1}})} \leq C(n, q, \lambda, \Lambda)^{\frac{1}{\gamma_i}} \|\bar{u}\|_{L^{\gamma_i}(B_{r_i})},$$

that is,

$$\|\bar{u}\|_{L^{\gamma_{i+1}}(B_{r_{i+1}})} \leq C^{\frac{i}{\chi^i}} \|\bar{u}\|_{L^{\gamma_i}(B_{r_i})}.$$

Hence by iteration we obtain

$$\|\bar{u}\|_{L^{\gamma_{i+1}}(B_{r_{i+1}})} \leq C^{\sum_{j=1}^i \frac{j}{\chi^j}} \|\bar{u}\|_{L^2(B_1)}$$

in particular

$$\|\bar{u}\|_{L^{\gamma_{i+1}}\left(B_{\frac{1}{2}}\right)} \leq C^{\sum_{j=1}^i \frac{j}{\gamma_j}} \|\bar{u}\|_{L^2(B_1)}$$

Letting $i \rightarrow \infty$ and using Fatou's Lemma we get

$$\sup_{B_{\frac{1}{2}}} \bar{u} \leq C \|\bar{u}\|_{L^2(B_1)},$$

hence

$$\sup_{B_{\frac{1}{2}}} u^+ \leq C \left(\|u^+\|_{L^2(B_1)} + k \right).$$

Since $k = \|f\|_{L^q}$, we finish the proof for $p = 2$.

2 De Giorgi's Method