

CS 325 Section 401, Project Group 1

David Rigert, Isaiah Perrotte-Fentress, Adam McDaniel

Project 3: Linear Programming

Problem 1: Transshipment Model

Part A: Determine the number of refrigerators to be shipped plants to warehouses and then warehouses to retailers to minimize the cost.

i) Formulate the problem as a linear program with an objective function and all constraints.

//all possible tuples of (plant, warehouse) and (warehouse, retailer) with associated costs

Minimize: $10(P1, W1) + 15(P1, W2) + 11(P2, W1) + 8(P2, W2) + 13(P3, W1) + 8(P3, W2) + 9(P3, W3) + 14(P4, W2) + 8(P4, W3) + 5(W1, R1) + 6(W1, R2) + 7(W1, R3) + 10(W1, R4) + 12(W2, R3) + 8(W2, R4) + 10(W2, R5) + 14(W2, R6) + 14(W3, R4) + 12(W3, R5) + 12(W3, R6) + 6(W3, R7)$

Constraints:

//shipping capacity of each plant

$(P1, W1) + (P1, W2) \leq 150$ //plant 1 supply

$(P2, W1) + (P2, W2) \leq 450$ //plant 2 supply

$(P3, W1) + (P3, W2) + (P3, W3) \leq 250$ //plant 3 supply

$(P4, W2) + (P4, W3) \leq 150$ //plant 4 supply

//warehouses are not endpoints, and must ship all units to retailers

$(P1, W1) + (P2, W1) + (P3, W1) - (W1, R1) - (W1, R2) - (W1, R3) - (W1, R4) = 0$ //warehouse 1

$(P1, W2) + (P2, W2) + (P3, W2) + (P4, W2) - (W2, R3) - (W2, R4) - (W2, R5) - (W2, R6) = 0$ //warehouse 2

$(P3, W3) + (P4, W3) - (W3, R4) - (W3, R5) - (W3, R6) - (W3, R7) = 0$ //warehouse 3

//demand of retailers

$(W1, R1) \geq 100$ //retailer 1 demand

$(W1, R2) \geq 150$ //retailer 2 demand

$(W1, R3) + (W2, R3) \geq 100$ //retailer 3 demand

$(W1, R4) + (W2, R4) + (W3, R4) \geq 200$ //retailer 4 demand

$(W2, R5) + (W3, R5) \geq 200$ //retailer 5 demand

$(W2, R6) + (W3, R6) \geq 150$ //retailer 6 demand

$(W3, R7) \geq 100$ //retailer 7 demand

//nonnegativity

All tuples ≥ 0

ii) Determine the optimal solution for the linear program using any software you want. Include a copy of the code/file in the report.

Lindo code and results:

MIN 10X1 + 15X2 + 11X3 + 8X4 + 13X5 + 8X6 + 9X7 + 14X8 + 8X9 + 5X10 + 6X11 + 7X12 + 10X13 + 12X14 + 8X15 + 10X16 + 14X17 + 14X18 + 12X19 + 12X20 + 6X21

ST

X1 + X2 < 150

X3 + X4 < 450

X5 + X6 + X7 < 250

X8 + X9 < 150

X1 + X3 + X5 - X10 - X11 - X12 - X13 = 0

X2 + X4 + X6 + X8 - X14 - X15 - X16 - X17 = 0

X7 + X9 - X18 - X19 - X20 - X21 = 0

X10 > 100

X11 > 150

X12 + X14 > 100

X13 + X15 + X18 > 200

X16 + X19 > 200

X17 + X20 > 150

X21 > 100

X1 > 0

X2 > 0

X3 > 0

X4 > 0

X5 > 0

X6 > 0

X7 > 0

X8 > 0

X9 > 0

X10 > 0

X11 > 0

X12 > 0

X13 > 0

X14 > 0

X15 > 0

X16 > 0

X17 > 0

X18 > 0

X19 > 0

X20 > 0

X21 > 0

END

LP OPTIMUM FOUND AT STEP 13

OBJECTIVE FUNCTION VALUE

1) 17100.00

VARIABLE	VALUE	REDUCED COST
X1	150.000000	0.000000
X2	0.000000	8.000000
X3	200.000000	0.000000
X4	250.000000	0.000000
X5	0.000000	2.000000
X6	150.000000	0.000000
X7	100.000000	0.000000
X8	0.000000	7.000000
X9	150.000000	0.000000
X10	100.000000	0.000000
X11	150.000000	0.000000
X12	100.000000	0.000000
X13	0.000000	5.000000
X14	0.000000	2.000000
X15	200.000000	0.000000
X16	200.000000	0.000000
X17	0.000000	1.000000
X18	0.000000	7.000000
X19	0.000000	3.000000
X20	150.000000	0.000000
X21	100.000000	0.000000

iii) What are the optimal shipping routes and minimum cost.

Minimum cost: \$17, 100

Optimal shipping routes:

Plant 1 ships 150 units to Warehouse 1.

Plant 2 ships 200 units to Warehouse 1 and 250 units to Warehouse 2.

Plant 3 ships 150 units to Warehouse 2 and 100 units to Warehouse 3.

Plant 4 ships 150 units to Warehouse 3.

Warehouse 1 ships 100 units to Retailer 1, 150 units to Retailer 2, and 100 units to Retailer 3.

Warehouse 2 ships 200 units to Retailer 4 and 200 units to Retailer 5.

Warehouse 3 ships 150 units to Retailer 6 and 100 units to Retailer 7.

Part B: Due to old infrastructure Warehouse 2 is going to close eliminating all of the associated routes. What is the optimal solution for this modified model? Is it feasible to ship all the refrigerators to either warehouse 1 or 3 and then to the retailers without using warehouse 2? Why or why not?

Removing warehouse 2 from the equation results in the modified program below:

//all possible tuples of (plant, warehouse) and (warehouse, retailer) with associated costs

Minimize: $10(P1, W1) + 11(P2, W1) + 13(P3, W1) + 9(P3, W3) + 8(P4, W3) + 5(W1, R1) + 6(W1, R2) + 7(W1, R3) + 10(W1, R4) + 14(W3, R4) + 12(W3, R5) + 12(W3, R6) + 6(W3, R7)$

Constraints:

//shipping capacity of each plant

$(P1, W1) \leq 150$ //plant 1 supply

$(P2, W1) \leq 450$ //plant 2 supply

$(P3, W1) + (P3, W3) \leq 250$ //plant 3 supply

$(P4, W3) \leq 150$ //plant 4 supply

//warehouses are not endpoints, and must ship all units to retailers

$(P1, W1) + (P2, W1) + (P3, W1) - (W1, R1) - (W1, R2) - (W1, R3) - (W1, R4) = 0$ //warehouse 1

$(P3, W3) + (P4, W3) - (W3, R4) - (W3, R5) - (W3, R6) - (W3, R7) = 0$ //warehouse 3

//demand of retailers

$(W1, R1) \geq 100$ //retailer 1 demand

$(W1, R2) \geq 150$ //retailer 2 demand

$(W1, R3) \geq 100$ //retailer 3 demand

$(W1, R4) + (W3, R4) \geq 200$ //retailer 4 demand

$(W3, R5) \geq 200$ //retailer 5 demand

$(W3, R6) \geq 150$ //retailer 6 demand

$(W3, R7) \geq 100$ //retailer 7 demand

//nonnegativity

All tuples ≥ 0

It is not feasible to eliminate Warehouse 2 from the model. While all plants still have at least 1 warehouse available to ship to and all retailers are still serviced by at least 1 warehouse, Retailers 5, 6, and 7 are serviced exclusively by Warehouse 3. Even if Plant 3 and Plant 4 ship all supply to Warehouse 3, Warehouse 3 will have at most 400 units available. The combined demand from Retailers 5, 6, and 7, is 450, and so some demand (50 units) will be unmet (IE, a constraint is unsatisfiable). Therefore, there is no optimal solution.

Lindo code and error message:

```

MIN 10X1 + 11X2 + 13X3 + 9X4 + 8X5 + 5X6 + 6X7 + 7X8 + 10X9 + 14X10 + 12X11 + 12X12 + 6X13
ST
  X1 < 150
  X2 < 450
  X3 + X4 < 250
  X5 < 150
  X1 + X2 + X3 - X6 - X7 - X8 - X9 = 0
  X4 + X5 - X10 - X11 - X12 - X13 = 0
  X6 > 100
  X7 > 150
  X8 > 100
  X9 + X10 > 200
  X11 > 200
  X12 > 150
  X13 > 100
  X1 > 0
  X2 > 0
  X3 > 0
  X4 > 0
  X5 > 0
  X6 > 0
  X7 > 0
  X8 > 0
  X9 > 0
  X10 > 0
  X11 > 0
  X12 > 0
  X13 > 0
END

```

```

Error code: 54
Error text: NO FEASIBLE SOLUTION AT STEP 10.
SUM OF INFEASIBILITIES = 50.0000000000
VIOLATED ROWS HAVE NEGATIVE SLACK, OR
(EQUALITY ROWS) NONZERO SLACKS. ROWS
CONTRIBUTING TO INFEASIBILITY HAVE A
NONZERO DUAL PRICE. USE THE "DEBUG"
COMMAND FOR MORE INFORMATION.

```

```

LINDO Solver Status
Optimizer Status
Status: Infeasible
Iterations: 10
Infeasibility: 50
Objective: 17650
Best IP: N/A
IP Bound: N/A
Branches: N/A
Elapsed Time: 00:01:28

```

Part C: Instead of closing Warehouse 2 management has decide to keep a portion of it open but limit shipments to 100 refrigerators per week. Is this feasible? If so what is the optimal solution when warehouse 2 is limited to 100 refrigerators?

//all possible tuples of (plant, warehouse) and (warehouse, retailer) with associated costs

Minimize: $10(P1, W1) + 15(P1, W2) + 11(P2, W1) + 8(P2, W2) + 13(P3, W1) + 8(P3, W2) + 9(P3, W3) + 14(P4, W2) + 8(P4, W3) + 5(W1, R1) + 6(W1, R2) + 7(W1, R3) + 10(W1, R4) + 12(W2, R3) + 8(W2, R4) + 10(W2, R5) + 14(W2, R6) + 14(W3, R4) + 12(W3, R5) + 12(W3, R6) + 6(W3, R7)$

Constraints:

//shipping capacity of each plant

$(P1, W1) + (P1, W2) \leq 150$ //plant 1 supply

$(P2, W1) + (P2, W2) \leq 450$ //plant 2 supply

$(P3, W1) + (P3, W2) + (P3, W3) \leq 250$ //plant 3 supply

$(P4, W2) + (P4, W3) \leq 150$ //plant 4 supply

//warehouses are not endpoints, and must ship all units to retailers

$(P1, W1) + (P2, W1) + (P3, W1) - (W1, R1) - (W1, R2) - (W1, R3) - (W1, R4) = 0$ //warehouse 1

$(P1, W2) + (P2, W2) + (P3, W2) + (P4, W2) - (W2, R3) - (W2, R4) - (W2, R5) - (W2, R6) = 0$ //warehouse 2

$(P3, W3) + (P4, W3) - (W3, R4) - (W3, R5) - (W3, R6) - (W3, R7) = 0$ //warehouse 3

//NEW constraint – Warehouse 2 cannot receive more than 100 units

$(P1, W2) + (P2, W2) + (P3, W2) + (P4, W2) \leq 100$

//demand of retailers

$(W1, R1) \geq 100$ //retailer 1 demand

$(W1, R2) \geq 150$ //retailer 2 demand

$(W1, R3) + (W2, R3) \geq 100$ //retailer 3 demand

$(W1, R4) + (W2, R4) + (W3, R4) \geq 200$ //retailer 4 demand

$(W2, R5) + (W3, R5) \geq 200$ //retailer 5 demand

$(W2, R6) + (W3, R6) \geq 150$ //retailer 6 demand

$(W3, R7) \geq 100$ //retailer 7 demand

//nonnegativity

All tuples ≥ 0

Adding 100 units of capacity to Warehouse 2 solves the issue we ran into in part B, by ensuring the demands of the retailers formerly only served by Warehouse 3 can now be met.

Lindo code and report:

```

MIN 10X1 + 15X2 + 11X3 + 8X4 + 13X5 + 8X6 + 9X7 + 14X8 + 8X9 + 5X10 + 6X11 + 7X12 + 10X13 + 12X14 + 8X15 +
10X16 + 14X17 + 14X18 + 12X19 + 12X20 + 6X21

ST
  X1 + X2 < 150
  X3 + X4 < 450
  X5 + X6 + X7 < 250
  X8 + X9 < 150
  X1 + X3 + X5 - X10 - X11 - X12 - X13 = 0
  X2 + X4 + X6 + X8 - X14 - X15 - X16 - X17 = 0
  X7 + X9 - X18 - X19 - X20 - X21 = 0
  X2 + X4 + X6 + X8 < 100
  X10 > 100
  X11 > 150
  X12 + X14 > 100
  X13 + X15 + X18 > 200
  X16 + X19 > 200
  X17 + X20 > 150
  X21 > 100
  X1 > 0
  X2 > 0
  X3 > 0
  X4 > 0
  X5 > 0
  X6 > 0
  X7 > 0
  X8 > 0
  X9 > 0
  X10 > 0
  X11 > 0
  X12 > 0
  X13 > 0
  X14 > 0
  X15 > 0
  X16 > 0
  X17 > 0
  X18 > 0
  X19 > 0
  X20 > 0
  X21 > 0
END

```

LP OPTIMUM FOUND AT STEP 15		
OBJECTIVE FUNCTION VALUE		
1)	18300.00	
VARIABLE	VALUE	REDUCED COST
X1	150.000000	0.000000
X2	0.000000	8.000000
X3	350.000000	0.000000
X4	100.000000	0.000000
X5	0.000000	4.000000
X6	0.000000	2.000000
X7	250.000000	0.000000
X8	0.000000	9.000000
X9	150.000000	0.000000
X10	100.000000	0.000000
X11	150.000000	0.000000
X12	100.000000	0.000000
X13	150.000000	0.000000
X14	0.000000	7.000000
X15	50.000000	0.000000
X16	50.000000	0.000000
X17	0.000000	4.000000
X18	0.000000	4.000000
X19	150.000000	0.000000
X20	150.000000	0.000000
X21	100.000000	0.000000

The optimal solution when Warehouse 2 is limited to 100 units of capacity is:

Minimum cost: \$18,300

Optimal shipping routes:

Plant 1 ships 150 units to Warehouse 1.

Plant 2 ships 350 units to Warehouse 1 and 100 units to Warehouse 2.

Plant 3 ships 250 units to Warehouse 3.

Plant 4 ships 150 units to Warehouse 3.

Warehouse 1 ships 100 units to Retailer 1, 150 units to Retailer 2, 100 units to Retailer 3, and 150 units to Retailer 4.

Warehouse 2 ships 50 units to Retailer 4 and 50 units to Retailer 5.

Warehouse 3 ships 150 units to Retailer 5, 150 units to Retailer 6, and 100 units to Retailer 7.

Part D: Formulate a generalized linear programming model for the transshipment problem. Give the objective function and constraints as mathematical formulas.

Minimize $\text{cost}(a, b) + \text{cost}(a, b+1) \dots$ for all valid values of b + $\text{cost}(a+1, b) \dots$ for all valid values of a + $\text{cost}(b, c) + \text{cost}(b, c+1) \dots$ for all valid values of c + $\text{cost}(b+1, c) \dots$ for all valid values of b , where a = Plant #, b = Warehouse #, and c = Retailer #. A valid value is one where plant a is able to ship to warehouse b , or where warehouse b is able to ship to retailer c .

Constraints:

$(a, b) + (a, b+1) \dots$ for all values of $b \leq \text{capacity of } a$

Repeat with an additional constraint for each value of a

$(a, b) + (a+1, b) \dots$ for all values of a , - $(b, c) - (b, c+1)$ for all values of $c = 0$

Repeat with an additional constraint for each value of b

$(b, c) + (b+1, c) \dots$ for all values of $b \geq \text{demand of } c$

Repeat with an additional constraint for each value of c

All values of a , b , and $c \geq 0$

Problem 2: A Mixture Problem

Each salad must contain:

- At least 15 grams of protein
- At least 2 and at most 8 grams of fat
- At least 4 grams of carbohydrates
- At most 200 milligrams of sodium
- At least 40% leafy greens by mass

The nutritional contents of these ingredients (per 100 grams) and cost are:

Ingredient Label	Ingredient	Energy	Protein	Fat	Carbs	Sodium	Cost
I1	Tomato	21.00	0.85	0.33	4.64	9.00	\$1.00
I2	Lettuce	16.00	1.62	0.20	2.37	28.00	\$0.75
I3	Spinach	40.00	2.86	0.39	3.63	65.00	\$0.50
I4	Carrot	41.00	0.93	0.24	9.58	69.00	\$0.50
I5	Sunflower Seeds	585.00	23.40	48.70	15.00	3.80	\$0.45
I6	Smoked Tofu	120.00	16.00	5.00	3.00	120.00	\$2.15
I7	Chickpeas	164.00	9.00	2.60	27.00	78.00	\$0.95
I8	Oil	884.00	0.00	100.00	0.00	0.00	\$2.00

Part A: Determine the combination of ingredients that minimizes calories but meets all nutritional requirements:

i) Formulate the problem as a linear program with an objective function and all constraints.

Decision Variables: I_y = 100 grams of each ingredient “y” to include in the salad. Each ingredient is labeled in order with the letter I and an incrementing number.

Objective Function: $\text{Min } K = I1*21 + I2*16 + I3*40 + I4*41 + I5*585 + I6*120 + I7*164 + I8*884$

Where K = kcal

Resource Constraints:

Protein: $I1*.85 + I2*1.62 + I3*2.86 + I4*.93 + I5*23.40 + I6*16 + I7*9 + I8*0 \geq 15$ g of protein

Fat Min: $I1*.33 + I2*.20 + I3*.39 + I4*.24 + I5*48.70 + I6*5 + I7*2.6 + I8*100 \geq 2$ g of fat

Fat Max: $I1*.33 + I2*.20 + I3*.39 + I4*.24 + I5*48.70 + I6*5 + I7*2.6 + I8*100 \leq 8$ g of fat

Carbs: $I1*4.64 + I2*2.37 + I3*3.63 + I4*9.58 + I5*15 + I6*3 + I7*27 + I8*0 \geq 4$ g of carbs

Sodium: $I1*9 + I2*28 + I3*65 + I4*69 + I5*3.80 + I6*120 + I7*78 + I8*0 \leq 200$ mg of sodium

Leafy Green: $(I1 + I2 + I3 + I4 + I5 + I6 + I7 + I8)*.4 \leq I2 + I3$

Non-Negative: $I_y \geq 0$

ii) Screenshots of code used to determine the optimal solution

```

MAX
Z:\minK.lbx
! Minimize calorie content of the salad
MIN 21 I1 + 16 I2 + 40 I3 + 41 I4 + 585 I5 + 120 I6 + 164 I7 + 884 I8
ST
! Constraints for protein, fatx2, carbs, sodium, and leafy greens
.85 I1 + 1.62 I2 + 2.86 I3 + .93 I4 + 23.40 I5 + 16 I6 + 9 I7 + 0 I8 > 15
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 > 2
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 < 8
4.64 I1 + 2.37 I2 + 3.63 I3 + 9.58 I4 + 15 I5 + 3 I6 + 27 I7 + 0 I8 > 4
9 I1 + 28 I2 + 65 I3 + 69 I4 + 3.80 I5 + 120 I6 + 78 I7 + 0 I8 < 200
.4 I1 + .4 I2 + .4 I3 + .4 I4 + .4 I5 + .4 I6 + .4 I7 + .4 I8 - I1 - I2 < 0
! Ensure no negative values for ingredients
I1 > 0
I2 > 0
I3 > 0
I4 > 0
I5 > 0
I6 > 0
I7 > 0
I8 > 0
END

```

```

MAX
LP OPTIMUM FOUND AT STEP      3
      OBJECTIVE FUNCTION VALUE
    1)      114.7541

```

VARIABLE	VALUE	REDUCED COST
I1	0.000000	10.890710
I2	0.585480	0.000000
I3	0.000000	20.524591
I4	0.000000	36.289616
I5	0.000000	408.387970
I6	0.878220	0.000000
I7	0.000000	97.551910
I8	0.000000	886.404358

ROW	SLACK OR SURPLUS	DUAL PRICES
2)	0.000000	-7.650273
3)	2.508197	0.000000
4)	3.491803	0.000000
5)	0.022248	0.000000
6)	78.220139	0.000000
7)	0.000000	6.010929
8)	0.000000	0.000000
9)	0.585480	0.000000
10)	0.000000	0.000000
11)	0.000000	0.000000
12)	0.000000	0.000000
13)	0.878220	0.000000
14)	0.000000	0.000000
15)	0.000000	0.000000

```

NO. ITERATIONS=      3

```

iii) What is the cost of the low calorie salad?

The solution is 58.55 grams of Lettuce @ \$0.75/100g and 87.82 grams of Smoked Tofu @ \$2.15/100g. This results in calories of 114.75 kcal for a **total cost of \$2.33**.

Part B: Determine the combination of ingredients that minimizes calories but meets all nutritional requirements:

i) Formulate the problem as a linear program with an objective function and all constraints.

Decision Variables: I_y = 100 grams of each ingredient “y” to include in the salad. Each ingredient is labeled in order with the letter I and an incrementing number.

Objective Function: $\text{Min } D = I_1 \cdot 1.00 + I_2 \cdot .75 + I_3 \cdot .50 + I_4 \cdot .50 + I_5 \cdot .45 + I_6 \cdot 2.15 + I_7 \cdot .95 + I_8 \cdot 2.00$

Where D = dollars spent

Resource Constraints:

Protein: $I_1 \cdot .85 + I_2 \cdot 1.62 + I_3 \cdot 2.86 + I_4 \cdot .93 + I_5 \cdot 23.40 + I_6 \cdot 16 + I_7 \cdot 9 + I_8 \cdot 0 \geq 15$ g of protein

Fat Min: $I_1 \cdot .33 + I_2 \cdot .20 + I_3 \cdot .39 + I_4 \cdot .24 + I_5 \cdot 48.70 + I_6 \cdot 5 + I_7 \cdot 2.6 + I_8 \cdot 100 \geq 2$ g of fat

Fat Max: $I_1 \cdot .33 + I_2 \cdot .20 + I_3 \cdot .39 + I_4 \cdot .24 + I_5 \cdot 48.70 + I_6 \cdot 5 + I_7 \cdot 2.6 + I_8 \cdot 100 \leq 8$ g of fat

Carbs: $I_1 \cdot 4.64 + I_2 \cdot 2.37 + I_3 \cdot 3.63 + I_4 \cdot 9.58 + I_5 \cdot 15 + I_6 \cdot 3 + I_7 \cdot 27 + I_8 \cdot 0 \geq 4$ g of carbs

Sodium: $I_1 \cdot 9 + I_2 \cdot 28 + I_3 \cdot 65 + I_4 \cdot 69 + I_5 \cdot 3.80 + I_6 \cdot 120 + I_7 \cdot 78 + I_8 \cdot 0 \leq 200$ mg of sodium

Leafy Green: $(I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8) \cdot .4 \leq I_2 + I_3$

Non-Negative: $I_y \geq 0$

ii) Screenshots of code used to determine the optimal solution

```

Z:\minD.txt
! Minimize cost of the salad
MIN 1 I1 + .75 I2 + .5 I3 + .5 I4 + .45 I5 + 2.15 I6 + .95 I7 + 2 I8
ST
! Constraints for protein, fatx2, carbs, sodium, and leafy greens
.85 I1 + 1.62 I2 + 2.86 I3 + .93 I4 + 23.40 I5 + 16 I6 + 9 I7 + 0 I8 > 15
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 > 2
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 < 8
4.64 I1 + 2.37 I2 + 3.63 I3 + 9.58 I4 + 15 I5 + 3 I6 + 27 I7 + 0 I8 > 4
9 I1 + 28 I2 + 65 I3 + 69 I4 + 3.80 I5 + 120 I6 + 78 I7 + 0 I8 < 200
.4 I1 + .4 I2 + .4 I3 + .4 I4 + .4 I5 + .4 I6 + .4 I7 + .4 I8 - I1 - I2 < 0
! Ensure no negative values for ingredients
I1 > 0
I2 > 0
I3 > 0
I4 > 0
I5 > 0
I6 > 0
I7 > 0
I8 > 0
END

```

```

LP OPTIMUM FOUND AT STEP      0

      OBJECTIVE FUNCTION VALUE

    1)      1.918084

      VARIABLE            VALUE            REDUCED COST
      I1                  0.000000            0.381411
      I2                  0.903303            0.000000
      I3                  0.000000            0.400018
      I4                  0.000000            0.700399
      I5                  0.093199            0.000000
      I6                  0.000000            0.223639
      I7                  1.261756            0.000000
      I8                  0.000000            8.429042

      ROW    SLACK OR SURPLUS    DUAL PRICES
      2)      0.000000           -0.160374
      3)      6.000000            0.000000
      4)      0.000000            0.060941
      5)      33.606209            0.000000
      6)      75.936424            0.000000
      7)      0.000000            0.837303
      8)      0.000000            0.000000
      9)      0.903303            0.000000
     10)      0.000000            0.000000
     11)      0.000000            0.000000
     12)      0.093199            0.000000
     13)      0.000000            0.000000
     14)      1.261756            0.000000
     15)      0.000000            0.000000

NO. ITERATIONS=         0

```

iii) How many calories are in the low cost salad?

The solution is 90.33 grams of Lettuce @ 16 kcal/100g, 9.32 grams of Sunflower Seeds @ 585 kcal/100g, and 126.18 grams of Chickpeas @ 164 kcal/100g. This results in a total cost of \$1.92 and **275.91 kcal for the salad.**

Part C: Compare the results from part A and B. Veronica's goal is to create a Very Veggie Salad that is both low calorie and low cost. She would like to sell the salad for \$5.00 and still have a profit of at least \$3.00. However if she can advertise that the salad has under 250 calories then she may be able to sell more.

	Low Calorie	Low Cost
Kcal	114.75	275.91
Total Cost	\$2.33	\$1.92

i) Suggest some possible ways that she select a combination of ingredients that is "near optimal" for both objectives. This is a type of multi-objective optimization.

To create a Linear Programming problem that can help the user solve for both of these items, the objective from one problem should become a constraint in the other problem. Typically there might be something specific that is driving the user to decide which one should be the constraints. In this case, the 2 goals are a salad that costs < \$2.00, and a salad that has under 250 kcal. In this case, the Low Calorie option is well below the < 250 calorie goal and exceeds the \$2.00 cost benchmark. Meanwhile, the Low Cost option is close to the kcal goal at 275.91 and costs \$1.92, just barely under the cost goal. Since the low cost solution is near optimal, I would recommend the user adds the low calorie constraint to the low cost problem. The user can then manually modify the low calorie constraint to fine tune the desired results. The user can continue to tighten (improve) the low calorie constraint until the increase in total cost is undesirable (the user will have to be able to decide which is more important after a certain point).

ii) What combination of ingredient would you suggest and what is the associated cost and calorie.

The problem setup:

Z:\minDmulti.txt	
Minimize cost of the salad	
MIN 1 I1 + .75 I2 + .5 I3 + .5 I4 + .45 I5 + 2.15 I6 + .95 I7 + 2 I8	
ST	
! Constraints for protein, fatx2, carbs, sodium, and leafy greens	
.85 I1 + 1.62 I2 + 2.86 I3 + .93 I4 + 23.40 I5 + 16 I6 + 9 I7 + 0 I8 > 15	
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 > 2	
.33 I1 + .20 I2 + .39 I3 + .24 I4 + 48.70 I5 + 5 I6 + 2.6 I7 + 100 I8 < 8	
4.64 I1 + 2.37 I2 + 3.63 I3 + 9.58 I4 + 15 I5 + 3 I6 + 27 I7 + 0 I8 > 4	
9 I1 + 28 I2 + 65 I3 + 69 I4 + 3.80 I5 + 120 I6 + 78 I7 + 0 I8 < 200	
.4 I1 + .4 I2 + .4 I3 + .4 I4 + .4 I5 + .4 I6 + .4 I7 + .4 I8 - I1 - I2 < 0	
! Add a kcal minimizing constraint	
21 I1 + 16 I2 + 40 I3 + 41 I4 + 585 I5 + 120 I6 + 164 I7 + 884 I8 < 250	
! Ensure no negative values for ingredients	
I1 > 0	
I2 > 0	
I3 > 0	
I4 > 0	
I5 > 0	
I6 > 0	
I7 > 0	
I8 > 0	
END	

Examples of Solutions:

LP OPTIMUM FOUND AT STEP 9			LP OPTIMUM FOUND AT STEP 0			LP OPTIMUM FOUND AT STEP 0			LP OPTIMUM FOUND AT STEP 0		
OBJECTIVE FUNCTION VALUE			OBJECTIVE FUNCTION VALUE			OBJECTIVE FUNCTION VALUE			OBJECTIVE FUNCTION VALUE		
1) 1.952702			1) 1.966066			1) 1.979431			1) 1.992796		
VARIABLE	VALUE	REDUCED COST	VARIABLE	VALUE	REDUCED COST	VARIABLE	VALUE	REDUCED COST	VARIABLE	VALUE	REDUCED COST
I1	0.000000	0.404085	I1	0.000000	0.404085	I1	0.000000	0.404085	I1	0.000000	0.404085
I2	0.833394	0.000000	I2	0.806404	0.000000	I2	0.779414	0.000000	I2	0.752425	0.000000
I3	0.000000	0.379449	I3	0.000000	0.379449	I3	0.000000	0.379449	I3	0.000000	0.379449
I4	0.000000	0.723834	I4	0.000000	0.723834	I4	0.000000	0.723834	I4	0.000000	0.723834
I5	0.091358	0.000000	I5	0.090647	0.000000	I5	0.089936	0.000000	I5	0.089225	0.000000
I6	0.154791	0.000000	I6	0.214551	0.000000	I6	0.274311	0.000000	I6	0.334071	0.000000
I7	1.003943	0.000000	I7	0.904409	0.000000	I7	0.804875	0.000000	I7	0.705341	0.000000
I8	0.000000	9.057467	I8	0.000000	9.057467	I8	0.000000	9.057467	I8	0.000000	9.057467
ROW	SLACK OR SURPLUS	DUAL PRICES	ROW	SLACK OR SURPLUS	DUAL PRICES	ROW	SLACK OR SURPLUS	DUAL PRICES	ROW	SLACK OR SURPLUS	DUAL PRICES
2)	0.000000	-0.182060	2)	0.000000	-0.182060	2)	0.000000	-0.182060	2)	0.000000	-0.182060
3)	6.000000	0.000000	3)	6.000000	0.000000	3)	6.000000	0.000000	3)	6.000000	0.000000
4)	0.000000	0.055510	4)	0.000000	0.055510	4)	0.000000	0.055510	4)	0.000000	0.055510
5)	26.916334	0.000000	5)	24.33569	0.000000	5)	21.750803	0.000000	5)	19.168037	0.000000
6)	79.435417	0.000000	6)	80.786270	0.000000	6)	82.137131	0.000000	6)	83.487991	0.000000
7)	0.000000	0.812581	7)	0.000000	0.812581	7)	0.000000	0.812581	7)	0.000000	0.812581
8)	0.000000	0.001336	8)	0.000000	0.001336	8)	0.000000	0.001336	8)	0.000000	0.001336
9)	0.000000	0.000000	9)	0.000000	0.000000	9)	0.000000	0.000000	9)	0.000000	0.000000
10)	0.833394	0.000000	10)	0.806404	0.000000	10)	0.779414	0.000000	10)	0.752425	0.000000
11)	0.000000	0.000000	11)	0.000000	0.000000	11)	0.000000	0.000000	11)	0.000000	0.000000
12)	0.000000	0.000000	12)	0.000000	0.000000	12)	0.000000	0.000000	12)	0.000000	0.000000
13)	0.091358	0.000000	13)	0.090647	0.000000	13)	0.089936	0.000000	13)	0.089225	0.000000
14)	0.154791	0.000000	14)	0.214551	0.000000	14)	0.274311	0.000000	14)	0.334071	0.000000
15)	1.003943	0.000000	15)	0.904409	0.000000	15)	0.804875	0.000000	15)	0.705341	0.000000
16)	0.000000	0.000000	16)	0.000000	0.000000	16)	0.000000	0.000000	16)	0.000000	0.000000
NO. ITERATIONS= 9			NO. ITERATIONS= 0			NO. ITERATIONS= 0			NO. ITERATIONS= 0		

Results Table:

Calorie Constraint	Total kcal	Total Cost
<250	250.00	\$1.95
<240	240.00	\$1.97
<230	230.00	\$1.98
<220	220.00	\$1.99

From a business perspective, assuming no incremental gain from lowering calories below 249 (one below 250 so the business is not caught lying and slandered in the media), the optimal solution would be 249 kcal at a cost of \$1.95. This is achieved by using 83.07 grams of Lettuce, 9.13 grams of Sunflower Seeds, 16.08 grams of Smoked Tofu, and 99.40 grams of Chickpeas.

LP OPTIMUM FOUND AT STEP 0		
OBJECTIVE FUNCTION VALUE		
1)	1.954038	
VARIABLE	VALUE	REDUCED COST
I1	0.000000	0.404085
I2	0.830695	0.000000
I3	0.000000	0.379449
I4	0.000000	0.723834
I5	0.091287	0.000000
I6	0.160767	0.000000
I7	0.993989	0.000000
I8	0.000000	9.057467

iii) Note: There is not one “right” answer. Discuss how you derived your solution.

As noted above, the solution was derived through a series of guess and check activities, starting with the minimum accepted answer (kcal below 250 to increase sales). From there, it was apparent that lowering kcal would result in increased costs. Since there is no incremental gain listed between 250 kcal and 220 kcal for this problem, than it is not worth incurring the extra cost. In a real world scenario, it might be worth using the lower kcal values at higher cost because the added marketing leverage could potentially increase sales.

Problem 3: Solving Shortest Path Problems Using Linear Programming

- a) What are the lengths of the shortest paths from vertex a to all other vertices.

VARIABLE	VALUE	REDUCED COST
DA	0.000000	0.000000
DB	2.000000	0.000000
DC	3.000000	0.000000
DD	8.000000	0.000000
DE	9.000000	0.000000
DF	6.000000	0.000000
DG	8.000000	0.000000
DH	9.000000	0.000000
DI	8.000000	0.000000
DJ	10.000000	0.000000
DK	14.000000	0.000000
DL	15.000000	0.000000
DM	17.000000	0.000000

```

MAX da + db + dc + dd + de + df + dg + dh +
di + dj + dk + dl + dm
ST
da = 0
db - da < 2
dc - da < 3
dd - da < 8
dh - da < 9
da - db < 4
dc - db < 5
de - db < 7
df - db < 4
dd - dc < 10
db - dc < 5
dg - dc < 9
di - dc < 11
df - dc < 4
da - dd < 8
dg - dd < 2
dj - dd < 5
df - dd < 1
dh - de < 5
dc - de < 4
di - de < 10
di - df < 2
dg - df < 2
dd - dg < 2
dj - dg < 8
dk - dg < 12
di - dh < 5
dk - dh < 10
da - di < 20
dk - di < 6
dj - di < 2
dm - di < 12
di - dj < 2
dk - dj < 4
dl - dj < 5
dh - dk < 10
dm - dk < 10
dm - dl < 2
da > 0
db > 0
dc > 0
dd > 0
de > 0
df > 0
dg > 0
dh > 0
di > 0
dj > 0
dk > 0
dl > 0
dm > 0

```

END

- b) If a vertex z is added to the graph for which there is no path from vertex a to vertex z, what will be the result when you attempt to find the lengths of shortest paths as in part a).

The distance value d_z for that vertex is unbounded because there are no constraints imposed on its value by edge weights, so you can increase the maximum forever by just increasing d_z .

c) What are the lengths of the shortest paths from each vertex to vertex m. How can you solve this problem with just one linear program?

VARIABLE	VALUE	REDUCED COST
DA	17.000000	0.000000
DB	15.000000	0.000000
DC	15.000000	0.000000
DD	12.000000	0.000000
DE	19.000000	0.000000
DF	11.000000	0.000000
DG	14.000000	0.000000
DH	14.000000	0.000000
DI	9.000000	0.000000
DJ	7.000000	0.000000
DK	10.000000	0.000000
DL	2.000000	0.000000
DM	0.000000	0.000000

For this version, instead of starting with a source vertex s and finding the shortest paths to all other vertices in the graph, we start from target vertex t and find all vertices that point to t, and work our way outward from there. The LINDO program is almost identical, except the target is set with the “= 0” constraint and we swap the operands of the subtraction operator in all of the edge constraints to reverse the direction.

```

MAX da + db + dc + dd + de + df + dg + dh +
di + dj + dk + dl + dm
ST
    dm = 0
    da - db < 2
    da - dc < 3
    da - dd < 8
    da - dh < 9
    db - da < 4
    db - dc < 5
    db - de < 7
    db - df < 4
    dc - dd < 10
    dc - db < 5
    dc - dg < 9
    dc - di < 11
    dc - df < 4
    dd - da < 8
    dd - dg < 2
    dd - dj < 5
    dd - df < 1
    de - dh < 5
    de - dc < 4
    de - di < 10
    df - di < 2
    df - dg < 2
    dg - dd < 2
    dg - dj < 8
    dg - dk < 12
    dh - di < 5
    dh - dk < 10
    di - da < 20
    di - dk < 6
    di - dj < 2
    di - dm < 12
    dj - di < 2
    dj - dk < 4
    dj - dl < 5
    dk - dh < 10
    dk - dm < 10
    dl - dm < 2
    da > 0
    db > 0
    dc > 0
    dd > 0
    de > 0
    df > 0
    dg > 0
    dh > 0
    di > 0
    dj > 0
    dk > 0
    dl > 0
END

```

- d) Suppose that all paths must pass through vertex i . How can you calculate the length of the shortest path from any vertex x to vertex y that pass through vertex i (for all $x, y \in V$)? Calculate the lengths of these paths for the given graph. (Note for some vertices x and y it may be impossible to pass through vertex i).

We can use the same linear program as part c) and set the target to vertex i . This gives us the shortest path from every reachable vertex u to vertex i . Any unreachable vertex is unbounded. Then we run the same linear program as part a) and set the source to vertex i . This gives us the shortest path from vertex i to every reachable vertex v . Once again, any unreachable vertex is unbounded. Finally, we simply add the values together for every permutation (u, i, v) for every $u, v \in V$ to get the shortest path from u to v via i .

VARIABLE	SP TO DI	SP FROM DI
DA	8.000000	20.000000
DB	6.000000	22.000000
DC	6.000000	23.000000
DD	3.000000	28.000000
DE	10.000000	29.000000
DF	2.000000	26.000000
DG	5.000000	28.000000
DH	5.000000	16.000000
DI	0.000000	0.000000
DJ	2.000000	2.000000
DK	15.000000	6.000000
DL	UNBOUNDED	7.000000
DM	UNBOUNDED	9.000000

Shortest Path Via i ($\delta(u,i,v)$)

$v \backslash u$	a	b	c	d	e	f	g	h	i	j	k	l	m
a	28	30	31	36	37	34	36	24	8	10	14	15	17
b	26	28	29	34	35	32	34	22	6	8	12	13	15
c	26	28	29	34	35	32	34	22	6	8	12	13	15
d	23	25	26	31	32	29	31	19	3	5	9	10	12
e	30	32	33	38	39	36	38	26	10	12	16	17	19
f	22	24	25	30	31	28	30	18	2	4	8	9	11
g	25	27	28	33	34	31	33	21	5	7	11	12	14
h	25	27	28	33	34	31	33	21	5	7	11	12	14
i	20	22	23	28	29	26	28	16	0	2	6	7	9
j	22	24	25	30	31	28	30	18	2	4	8	9	11
k	35	37	38	43	44	41	43	31	15	17	21	22	24
l	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
m	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A