

# Poincaré Schemes

Ben Antieau, Viktor Burghardt, Noah Riggenbach, Lucy Yang

## Abstract

We do stuff

N: Change  
this

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Azumaya algebras and their involutions . . . . .	1
1.2	Main results . . . . .	2
1.3	Outline . . . . .	3
1.4	Acknowledgements . . . . .	3
1.5	Conventions . . . . .	3
<b>2</b>	<b>Poincaré Structures on Compact Modules</b>	<b>3</b>
<b>3</b>	<b>Poincaré Ring Spectra</b>	<b>3</b>
3.1	Algebras with genuine involution . . . . .	5
<b>4</b>	<b>The Poincaré Picard space</b>	<b>6</b>
4.1	prime factorization and the picard group of hearts . . . . .	9
4.2	Hermitian line bundles . . . . .	10
<b>5</b>	<b>Poincaré structures on schemes with involution</b>	<b>11</b>
5.1	Poincaré structures associated to rigid symmetric monoidal $\infty$ -categories with involution . . .	11
5.2	From schemes with involution to Poincaré structures on module categories . . . . .	14
<b>6</b>	<b>The Poincaré Brauer Group</b>	<b>17</b>
6.1	Azumaya algebras with genuine involution . . . . .	22
6.2	Stacks associated to Poincaré- $\infty$ -categories . . . . .	24
<b>7</b>	<b>Poincaré schemes</b>	<b>25</b>

## 1 Introduction

### 1.1 Azumaya algebras and their involutions

Let  $X$  be a scheme. Gabber showed that if  $X$  is quasi-compact and separated and admits an ample line bundle, the collection of sheaves of Azumaya  $\mathcal{O}_X$ -algebras up to Morita equivalence is in bijection with the torsion subgroup of  $H_{\text{ét}}^2(X; \mathbb{G}_m)$  [Gab81]. On the other hand, to  $\mathcal{A}$  we may associate the presentable  $\mathcal{O}_X$ -linear  $\infty$ -category  $\mathcal{D}(\mathcal{A})$ . If  $\mathcal{A}$  and  $\mathcal{A}'$  are Morita equivalent, then  $\mathcal{D}(\mathcal{A}')$  and  $\mathcal{D}(\mathcal{A})$  are equivalent in  $\text{Pr}_{\mathcal{O}_X}^L$ , and that  $\mathcal{A}$  is Azumaya implies that  $\mathcal{D}(\mathcal{A})$  is invertible in  $\text{Pr}_{\mathcal{O}_X}^L$  with respect to the  $\mathcal{O}_X$ -linear tensor product.

**Definition 1.1.** The *derived Brauer space* of  $X$  is  $\text{dBr}(X) := \text{Pic}(\text{Pr}_{\mathcal{O}_X}^L)$ .

L: AG nota-  
tion or homo-  
topy notation?

L: who is this  
observation  
due to?

L: cite  
Lieblich/de  
Jong? con-  
nect to twisted  
sheaves!

Toën shows that the assignment  $X \mapsto \mathrm{dBr}(X)$  is an étale sheaf and there is an isomorphism  $\pi_0 \mathrm{dBr}(X) \simeq H_{\mathrm{ét}}^1(X; \mathbb{Z}) \times H_{\mathrm{ét}}^2(X; \mathbb{G}_m)$  [Toë12, Corollary 2.12]. Furthermore, Toën shows that for any qcqs scheme  $X$ , any invertible  $\mathcal{O}_X$ -linear  $\infty$ -category  $\mathcal{C}$  has a compact generator  $G$ ; thus we may write  $\mathcal{C} \simeq \mathrm{Mod}_{\mathrm{End}_{\mathcal{C}}(G)}(\mathrm{Mod}_{\mathcal{O}_X})$  and  $\mathrm{End}_{\mathcal{C}}(G)$  is said to be a *derived/generalized (sheaf of) Azumaya algebras* over  $\mathcal{O}_X$  (see Theorem 3.7 and Corollary 3.8 of [Toë12]). In particular, Toën’s result gives concrete interpretations/realizations of all/not necessarily torsion classes in  $H_{\mathrm{ét}}^2(X; \mathbb{G}_m)$ , at the cost of using derived methods/considering derived objects.

Antieau and Gepner extended Toën’s work to connective ring spectra/affine spectral schemes with connective rings of functions in [AG14].

On the other hand, the theory of anti-involutions on Azumaya algebras is an essential tool for studying the behavior of Brauer classes under multiplication by 2 and/or norm maps/corestriction. An anti-involution on an Azumaya algebra over a ring  $R$  is an equivalence  $\sigma: A \xrightarrow{\sim} A^{\mathrm{op}}$  so that  $\sigma^{\mathrm{op}} \circ \sigma = \mathrm{id}$ . The anti-involution is said to be of type 1 if it acts by the identity on  $R =$  the center of  $A$ . If instead  $R$  is endowed with an involution  $\lambda$  so that the inclusion of the subring of fixed elements  $R^\lambda \rightarrow R$  exhibits  $R$  as a quadratic étale  $R^\lambda$ -algebra and  $\sigma$  agrees with  $\lambda$  on the center of  $A$ , then the involution  $\sigma$  is said to be of *type 2*.

If an Azumaya algebra  $A$  has an anti-involution, its Brauer class  $[A] \in H_{\mathrm{ét}}^2(-; \mathbb{G}_m)$  is manifestly 2-torsion. More surprisingly, if an Azumaya algebra  $A$  is such that  $[A]$  lies in the 2-torsion subgroup of  $H_{\mathrm{ét}}^2(-; \mathbb{G}_m)$ , then there exists an Azumaya algebra  $A'$  in the Brauer class of  $A$  admitting an anti-involution; the result was proved for  $R$  a field by Albert (in fact Albert proves that one can take  $A' = A$ ), for an arbitrary ring by Saltman, and for schemes  $X$  with  $\frac{1}{2} \in \mathcal{O}_X$  by Parimala–Srinivas [Alb61, §9 Theorem 19; Sal78, Theorem 3.1(a); PS92, Theorem 1].

On the other hand, consider an étale cover  $X \rightarrow Y$  of degree 2 where  $\frac{1}{2} \in \mathcal{O}_Y$ , and let  $\lambda$  denote the nontrivial  $C_2$ -Galois action on  $X$ . There is an *involutive Brauer group*  $\mathrm{Br}(X, \lambda)$  consisting of equivalence classes of sheaves of Azumaya  $\mathcal{O}_X$ -algebras with involutions of the second kind [PS92, p.216]. Parimala–Srinivas showed that the involutive Brauer group sits in an exact sequence  $\mathrm{Pic} X \xrightarrow{N} \mathrm{Pic} Y \rightarrow \mathrm{Br}(X, \lambda) \rightarrow \mathrm{Br}(X) \xrightarrow{N} \mathrm{Br}(Y)$  [PS92, Theorem 2].

First and Williams observed that the aforementioned two cases comprise two extreme ends/special cases of a spectrum: multiplication by 2 on  $H_{\mathrm{ét}}^2(-; \mathbb{G}_m)$  may be regarded as a cohomological  $C_2$ -transfer map along the ‘quotient’ map  $X = X$ , where  $X$  is regarded as a scheme with trivial  $C_2$ -action [FW20, §1.2]. In other words, the former comprises trivial  $C_2$ -actions with ‘everywhere ramified’ quotient map, whereas the latter comprises free  $C_2$ -actions with nowhere ramified quotient maps. Moreover, First and William observe that a quotient involves a choice/is extra data; from the perspective of the stacky quotient  $X \rightarrow X//C_2$ , the  $C_2$ -action on  $X$  is free. Thus in order to study involutions systematically, it is necessary to specify a scheme with involution and the choice of a quotient.

The study of involutions on Azumaya algebras goes hand-in-hand with/is inextricably linked to the study of symmetric bilinear/hermitian forms on vector bundles: étale-locally, (classical) Azumaya algebras can be described as endomorphism algebras of vector bundles  $\mathcal{A} \simeq \mathcal{E}nd_X(V)$ . Taking transposes and conjugating by an isomorphism  $q: V \rightarrow V^\vee$  adjoint to a nondegenerate pairing  $q: V \otimes V \rightarrow \mathcal{O}_X$  comprise a ‘prototypical’ example of an involution on  $\mathcal{E}nd_X(V)$ . In [Cal+20a; Cal+20b; Cal+21], Calmès–Dotto–Harpaz–Hebestreit–Land–Moi–Nardin–Nikolaus–Steimle introduce a framework for talking about objects of stable  $\infty$ -categories equipped with the data of nondegenerate hermitian forms. Motivated by this connection between involutions and duality (cf. [Cal+20a, §3]), we use Poincaré  $\infty$ -categories to define a derived enhancement of the involutive Brauer group.

## 1.2 Main results

**Theorem 1.2.** *Let  $\underline{A}$  be an affine Poincaré scheme with underlying  $\mathbb{E}_\infty$ -ring spectrum with involution  $A$ . Then the natural maps*

$$\pi_i(\mathrm{PnPic}(\underline{A})) \rightarrow \pi_i(\mathrm{Pic}(A))$$

*are surjective on 2-torsion.*

**Theorem 1.3.** *Let  $A$  be an  $\mathbb{E}_\infty$  ring with involution, and let  $\underline{NA}$  be the associated Tate affine Poincaré scheme. Let  $\mathrm{Br}_\nu(A)$  be the Brauer group of Azumaya algebras over  $A$  with involution. Then the natural map*

$$\mathrm{PnBr}(\underline{NA}) \rightarrow \mathrm{Br}_\nu(A)$$

L: cite this too?

L: connect to ‘many choices of different Poincaré structures’ on a given  $\infty$ -category eventually

L: this is so close to Spec of a Poincaré ring but also not quite.

L: insert adjointives

L: also cite Jacob

N: I think there is some interaction with the homotopy fixed points, or maybe even the genuine

is an equivalence

N: probably  
of  $\mathbb{E}_\infty$  do-  
dads

**Theorem 1.4.** *The functors  $\mathrm{PnPic}, \mathrm{PnBr} : \mathrm{APS} \rightarrow \mathrm{Sp}$  are fppf sheaves.*

**Theorem 1.5.** *There is a Poincaré group scheme  $\mathbb{G}_m^\circ$  such that*

$$B\mathbb{G}_m^\circ \simeq \mathrm{PnPic}$$

*as fppf stacks.*

### 1.3 Outline

### 1.4 Acknowledgements

The authors wish to thank the Institute for Advanced Study and the organizers of the 2024 Park City Mathematics institute on motivic homotopy theory.

### 1.5 Conventions

$\mathrm{Br}^{\mathrm{P}}$	Poincaré Brauer space
$\mathrm{CAlg}$	$\infty$ -category of $\mathbf{E}_\infty$ -ring spectra
$\mathrm{CAlg}(\mathcal{S})$	$\infty$ -category of $\mathbf{E}_\infty$ -spaces
$\mathrm{CAlg}^{\mathrm{gp}}(\mathcal{S})$	$\infty$ -category of grouplike $\mathbf{E}_\infty$ -spaces
$\mathrm{CAlg}^{\mathrm{P}}$	$\infty$ -category of Poincaré ring spectra
$\mathrm{Cat}_\infty^{\mathrm{ex}}$	$\infty$ -category of small stable $\infty$ -categories and exact functors
$\mathrm{Cat}_\infty^{\mathrm{P}}$	$\infty$ -category of Poincaré $\infty$ -categories
$\mathrm{Cat}_{\infty, \mathrm{idem}}^{\mathrm{P}}$	$\infty$ -category of idempotent complete Poincaré $\infty$ -categories
$\mathrm{Pic}^{\mathrm{P}}$	Poincaré Picard space
$\mathcal{S}$	$\infty$ -category of spaces
$\mathrm{Sp}$	$\infty$ -category of spectra

## 2 Poincaré Structures on Compact Modules

We will use this section to recall notions and results about Poincaré  $\infty$ -categories which we require in the sections to follow. This section can safely be skipped by anyone with extensive knowledge of Poincaré  $\infty$ -categories, as found in [Cal+20a].

**Notation 2.1.** Let  $R$  be an  $\mathbf{E}_\infty$ -ring spectrum. We will drop  $\mathbf{E}_\infty$  from our notation and simply call  $R$  a *ring spectrum*. Moreover, we will denote the  $\infty$ -category  $\mathrm{CAlg}(\mathrm{Sp})$  of commutative algebra objects in the  $\infty$ -category of spectra  $\mathrm{Sp}$  by  $\mathrm{CAlg}$ .

Let  $R$  be a ring spectrum and let  $\mathrm{Mod}_R$  be the  $\infty$ -category of modules over  $R$ . We will study Poincaré structures on the  $\infty$ -category  $\mathrm{Mod}_R^\omega$  of compact modules over  $R$ .

## 3 Poincaré Ring Spectra

In this section we will define the ring theoretic building blocks of Poincaré schemes and the corresponding category they live in. Affine Poincaré Schemes will then be the dual objects, similar to how affine schemes are dual to commutative rings.

**Definition 3.1.** Let  $R$  be a ring spectrum. A *Poincaré structure* on  $R$  is the following data:

- A  $C_2$ -action on  $R$  via maps of ring spectra, i.e. a functor  $\lambda : BC_2 \rightarrow \mathrm{CAlg}$ .
- An  $\mathbb{E}_\infty$ - $R$ -algebra  $R \rightarrow C$ .

V: -  
characterizati  
in terms  
of modules  
with genuine  
involution, -  
characterizati  
of symmetric  
monoidal  
structures,  
-Pn

- An  $\mathbb{E}_\infty$ - $R$ -algebra map  $C \rightarrow R^{tC_2}$ .

Here  $R^{tC_2}$  is the Tate construction with respect to the given action. Since the Tate construction is lax symmetric monoidal,  $R^{tC_2}$  is naturally an  $R$ -algebra via the Tate-valued norm. A ring spectrum equipped with a Poincaré structure will be called a *Poincaré ring spectrum*.

**Remark 3.2.** In [Cal+20a, discussion immediately preceding Examples 5.4.10], Poincaré ring spectra are referred to as  *$\mathbb{E}_\infty$ -ring spectra with genuine involution*.

**Remark 3.3.** Let  $R$  be a ring spectrum. By [Cal+20a, Corollary 5.4.8], a Poincaré structure on  $R$  gives rise to a symmetric monoidal lift of  $\text{Mod}_R^\omega$  to the symmetric monoidal  $\infty$ -category of Poincaré  $\infty$ -categories  $\mathfrak{P}_R : (\text{Mod}_R^\omega)^{\text{op}} \rightarrow \text{Sp}$ . Furthermore, the structure map  $R \rightarrow C$  gives a canonical lift of  $R \in \text{Mod}_R^\omega$  to a Poincaré object  $(R, q) \in \text{Pn}(\text{Mod}_R^\omega, \mathfrak{P}_R)$ .

**Remark 3.4.** A Poincaré structure on a ring spectrum  $R$  with a  $C_2$ -action via maps of ring spectra is a factorization  $R \rightarrow C \rightarrow R^{tC_2}$  in  $\text{CAlg}$  of the Tate Frobenius  $R \rightarrow R^{tC_2}$ .

**Observation 3.5.** There is a forgetful functor from  $C_2$ - $\mathbb{E}_\infty$ -algebras to Poincaré rings which forgets the  $C_2$ -equivariance of the map  $R \rightarrow R^{tC_2}$ .

**Remark 3.6.** By [Cal+20a, §5.1], the assignment  $(R, R \rightarrow C \rightarrow R^{tC_2}) \mapsto (\text{Mod}_R^\omega, \mathfrak{P}_R)$  promotes to a symmetric monoidal functor  $\text{CAlg}^{\text{p}} \rightarrow \text{CAlg}(\text{Cat}_\infty^{\text{p}})$ .

**Notation 3.7.** Let  $R$  be an  $\mathbb{E}_\infty$ -ring spectrum. We will denote by  $\underline{R}$  the spectrum  $R$  with trivial  $C_2$ -action. More precisely,  $\underline{R} : BC_2 \rightarrow \text{Sp}$  is the constant functor.

**Example 3.8.** Let  $R$  be a ring spectrum with a  $C_2$ -action. If  $2 \in \pi_0(R)$  is invertible, we have  $\underline{R}^{tC_2} \simeq 0$  by [NS18, Lemma I.2.8]. A Poincaré structure on  $R$  is equivalent to the data of an  $\mathbb{E}_\infty$ - $R$ -algebra  $R \rightarrow C$ .

**Example 3.9.** Let  $R$  be a ring spectrum equipped with a  $C_2$ -action via maps of ring spectra. The Tate-valued norm endows  $R^{tC_2}$  with a natural  $R$ -algebra structure, which induces a Poincaré structure on  $R$  given by the factorization  $R \xrightarrow{\text{id}} R \rightarrow R^{tC_2}$ . We will call this Poincaré structure the *Tate Poincaré structure on  $R$*  and will denote it by  $(R, \mathfrak{P}_R^t)$ .

**Example 3.10.** The sphere spectrum  $\mathbb{S}$  together with the Tate Poincaré structure will be called the *universal Poincaré ring spectrum* (see [Cal+20a, §4.1]). We will denote it by  $(\mathbb{S}, \mathfrak{P}_u)$ .

**Remark 3.11.** Let  $(R, \mathfrak{P})$  be a ring spectrum associated to a factorization  $R \rightarrow C \rightarrow R^{tC_2}$ . A factorization of the map  $C \rightarrow R^{tC_2}$  through  $R^{hC_2}$  induces a section of the canonical map  $\mathfrak{P}(R) \rightarrow \text{hom}_R(R, C) \simeq C$ . In that case, we have a splitting  $\mathfrak{P}(R) \simeq R_{hC_2} \oplus C$ .

**Example 3.12.** The Tate Frobenius for the sphere spectrum factors through  $\mathbb{S}^{hC_2}$ . Therefore, Remark 3.11 implies  $\mathfrak{P}_u(\mathbb{S}) \simeq \mathbb{S}_{hC_2} \oplus \mathbb{S} \simeq \Sigma^\infty(\mathbb{P}_\mathbb{R}^\infty) \oplus \mathbb{S}$ .

**Example 3.13.** Let  $R$  be a ring spectrum equipped with a  $C_2$ -action via maps of ring spectra. The identity map  $\text{id} : R^{tC_2} \rightarrow R^{tC_2}$  induces a Poincaré structure on  $R$  given by the factorization  $R \rightarrow R^{tC_2} \xrightarrow{\text{id}} R^{tC_2}$ . We will call this Poincaré structure the *symmetric Poincaré structure on  $R$* .

**Example 3.14.** Let  $R$  be a connective ring spectrum equipped with a  $C_2$ -action via maps of ring spectra. The connective cover  $\tau_{\geq 0}(R^{tC_2}) \rightarrow R^{tC_2}$  of  $R^{tC_2}$  induces a Poincaré structure on  $R$  given by the factorization  $R \rightarrow \tau_{\geq 0}(R^{tC_2}) \rightarrow R^{tC_2}$ . We will call this Poincaré structure the *genuine symmetric Poincaré structure on  $R$* .

**Example 3.15.** Let  $R$  be a commutative ring endowed with an involution  $\sigma : R \xrightarrow{\sim} R$ . Write  $\underline{R}^\sigma$  for the  $C_2$ -Green functor with  $C_2$ -fixed points  $R^{C_2}$ , where  $R^{C_2}$  denotes the strict fixed points of the  $C_2$ -action on  $R$ , and underlying object  $R$ . The Mackey functor  $\underline{R}^\sigma$  is a  $C_2$ - $\mathbb{E}_\infty$  ring, therefore in particular we may regard it as a Poincaré ring by Observation 3.5. This is a special case of Example 3.14.

L: This is commonly used for constant Mackey functors—could be ambiguous

V: reference pullback that characterizes all quadratic functors

V: copy more examples from notes

### 3.1 Algebras with genuine involution

**Recollection 3.16.** Assume  $\mathcal{C}$  is a presentable monoidal  $\infty$ -category such that the monoidal product  $- \otimes -: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  preserves small colimits separately in each variable. Then there is an  $\infty$ -category  $\mathrm{LMod}(\mathcal{C})$  [Lur17, Example 4.2.1.18] whose objects are pairs  $(A, M)$  where  $A$  is an associative algebra object of  $\mathcal{C}$  and  $M$  is a left  $A$ -module. Write  $a, m$  respectively for the canonical forgetful functors  $\mathrm{LMod}(\mathcal{C}) \rightarrow \mathrm{Alg}(\mathcal{C})$ ,  $\mathrm{LMod}(\mathcal{C}) \rightarrow \mathcal{C}$  which send  $(A, M)$  to  $A$  and  $M$ , resp. Then  $a$  is a cocartesian fibration [Lur17, Corollary 4.2.3.7], hence it is classified by a functor  $\mathrm{mod}: \mathrm{Alg}(\mathcal{C}) \rightarrow \mathrm{Cat}_\infty$ .

The functor  $s$  of [Lur17, Example 4.2.1.17] determines a natural transformation  $\eta: * \rightarrow \mathrm{mod}$ , where  $*: \mathrm{Alg}(\mathcal{C}) \rightarrow \{*\} \hookrightarrow \mathrm{Cat}_\infty$  is the constant functor at the trivial category, or equivalently

$$\begin{array}{ccc} & & \mathcal{U} \\ & \nearrow \eta & \downarrow \\ \mathrm{Alg}(\mathcal{C}) & \xrightarrow{\mathrm{mod}} & \mathrm{Cat}_\infty \end{array} \quad (3.17)$$

where  $\mathcal{U}$  is the universal cocartesian fibration. Now consider the functor  $o: \mathcal{U} \rightarrow \mathrm{Cat}_\infty$  which sends  $(\mathcal{D}, d \in \mathcal{D})$  to the undercategory  $\mathcal{D}_{d/-}$ . Define  $\mathrm{LMod}(\mathcal{C})_{*/-}$  to be the cocartesian fibration over  $\mathrm{Alg}(\mathcal{C})$  classified by  $o \circ \eta \circ \mathrm{mod}$ .

**Variant 3.18.** Let  $\mathcal{C}$  be as in Recollection 3.16. There is a similar construction where left modules is replaced by *bimodules* [Lur17, Definition 4.3.1.12].

**Construction 3.19.** Regard  $\mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp})$  as a category with  $C_2$ -action given by taking the opposite/reverse algebra. There are functors  $b: \mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp})^{hC_2} \rightarrow \mathrm{LMod}(\mathrm{Sp})$  and  $b: \mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp})^{hC_2} \rightarrow \mathrm{BiMod}(\mathrm{Sp})_{*/-}$  so that  $(a, m) \circ b$  and  $(a, m) \circ b_*$  are (canonically) equivalent to  $(-^e) \otimes (-^e)^{\mathrm{op}}, (-)^e$ . Informally, an  $\mathbb{E}_1$ -algebra with involution  $B$  can be regarded as a  $B \otimes B^{\mathrm{op}}$ -module in a canonical way, and there is a canonical  $B \otimes B^{\mathrm{op}}$ -module map  $B \otimes B^{\mathrm{op}} \rightarrow B$ .

**Definition 3.20.** The category of  $\mathbb{E}_1$ -algebras with genuine involution is defined to be the limit of the  $\mathrm{Cat}_\infty$ -valued diagram

$$\begin{array}{ccccc} & & \mathrm{LMod}(\mathrm{Sp}) & & \\ & & \downarrow a, m & \swarrow (-)^e & \\ & & \mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp}) \times \mathrm{Sp} & & \mathrm{LMod}(\mathrm{Sp}^{C_2}) \\ & \nearrow b \circ (-^e) & & \searrow (-)^e \times (-)^e & \downarrow a, m \\ \mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp})^{hC_2} \times_{\mathrm{Sp}} \mathrm{Sp}^{\Delta^1} & \xrightarrow{N^{C_2}(-^e) \times U} & \mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp}^{C_2}) \times \mathrm{Sp}^{C_2} & & \end{array} \quad (3.21)$$

where

- $b$  is the functor/section of Construction 3.19
- $U$  is the ‘underlying’  $C_2$ -spectrum functor  $\mathbb{E}_1 \mathrm{Alg}(\mathrm{Sp})^{BC_2} \times_{\mathrm{Sp}} \mathrm{Sp}^{\Delta^1} \rightarrow \mathrm{Sp}^{BC_2} \times_{\mathrm{Sp}} \mathrm{Sp}^{\Delta^1} \simeq \mathrm{Sp}^{C_2}$
- The upper right trapezoid commutes canonically by definition of  $\mathrm{LMod}$  (and the fact that the functors  $a, m$  are given by restriction to subcategories of  $LM^\otimes$ ).

Write  $\mathbb{E}_1 \mathrm{Alg}^{\mathrm{gi}}(\mathrm{Sp}^{C_2})$  for the  $\infty$ -category of  $\mathbb{E}_1$ -algebras with genuine involution.

**Definition 3.22.** The category of  $\mathbb{E}_\sigma$ -algebras is defined to be the limit of the  $\mathcal{C}at_\infty$ -valued diagram

$$\begin{array}{ccc}
 & \text{BiMod}(\text{Sp})_{*/-} & \\
 & \downarrow a,m & \swarrow (-)^e \\
 & \mathbb{E}_1\text{Alg}(\text{Sp}) \times \text{Sp} & \text{BiMod}(\text{Sp}^{C_2})_{*/-} \\
 & \downarrow a,m & \\
 & \mathbb{E}_1\text{Alg}(\text{Sp}^{C_2}) \times \text{Sp}^{C_2} & \\
 \uparrow (-)^e \otimes (-)^{op}, (-)^e & & \uparrow (-)^e \times (-)^e \\
 \mathbb{E}_1\text{Alg}(\text{Sp})^{hC_2} \times_{\text{Sp}} \text{Sp}^{\Delta^1} & \xrightarrow{N^{C_2}(-)^e \times U} & \mathbb{E}_1\text{Alg}(\text{Sp}^{C_2}) \times \text{Sp}^{C_2}
 \end{array} \quad (3.23)$$

Write  $\mathbb{E}_\sigma\text{Alg}(\text{Sp}^{C_2})$  for the  $\infty$ -category of  $\mathbb{E}_\sigma$ -algebras.

**Variant 3.24.** Let the base be  $R$  an  $\mathbb{E}_\infty$ -algebra or Poincaré ring instead of  $\mathbb{S}^0$ .

**Remarks 3.25.** 1. Compare [AGH21, Corollary 3.10].

2. There are canonical forgetful functors  $\mathbb{E}_\sigma\text{Alg} \rightarrow \mathbb{E}_1\text{Alg}^{\text{gi}} \rightarrow \mathbb{E}_1\text{Alg}^{hC_2} \rightarrow \mathbb{E}_1\text{Alg}(\text{Sp})$ .

**Construction 3.26.** Let  $R, R^{\varphi C_2} \rightarrow R^{tC_2}$  be a Poincaré ring. There is a functor  $(\text{Mod}_{(-)}^\omega, \Omega_{(-)}) : \mathbb{E}_\sigma\text{Alg}_R \rightarrow (\text{Cat}_R^h)_{(\text{Mod}_R^\omega, \Omega_R)/-}$ .

**Lemma 3.27.** Let  $R, R^{\varphi C_2} \rightarrow R^{tC_2}$  be a Poincaré ring. The functor of Construction 3.26 is fully faithful.

*Proof.* □

Now we observe that given a hermitian object  $(x, q)$  of  $(\mathcal{C}, \Omega_{\mathcal{C}})$ , its endomorphism algebra admits a canonical lift to a  $\mathbb{E}_\sigma$ -algebra.

**Construction 3.28.** There is a functor  $\text{End}(-) : (\text{Cat}_R^h)_{(\text{Mod}_R^\omega, \Omega_R)} \rightarrow \mathbb{E}_\sigma\text{Alg}$  lifting the functor  $(\text{Cat}_R^h)_{(\text{Mod}_R^\omega, \Omega_R)} \rightarrow \mathbb{E}_1\text{Alg}^{hC_2}$  of [Cal+20a, Proposition 3.1.16].

**Theorem 3.29.** The functors of Construction 3.26 and 3.28 form an adjoint pair.

## 4 The Poincaré Picard space

Recall that the Poincaré space functor  $\text{Pn} : \text{Cat}_\infty^{\text{P}} \rightarrow \text{CAlg}(\mathcal{S})$  is lax symmetric monoidal with respect to tensor product of Poincaré  $\infty$ -categories and smash product of  $\mathbf{E}_\infty$ -spaces [Cal+20a, Corollary 5.2.8]. In particular, we can consider invertible objects in  $\text{Pn}(A)$  for a Poincaré ring spectrum  $A$ .

**Definition 4.1.** Let  $A$  be a Poincaré ring spectrum. We define the *Picard space* of  $A$  to be

$$\text{Pic}^{\text{P}}(A) := \text{Pic}(\text{Pn}(A)).$$

**Remark 4.2.** Let  $(\text{Mod}_R^\omega, \Omega_R)$  be a Poincaré ring spectrum, where  $(M_R = R, N_R = R^{\varphi C_2}, R^{\varphi C_2} \rightarrow R^{tC_2})$  is the module with genuine involution associated to  $\Omega_R$ . Then a point in the Poincaré Picard space is the data of a pair  $(\mathcal{L}, q)$ , where  $\mathcal{L}$  is an invertible module in  $\text{Mod}_R^\omega$  and  $q$  is a point in  $\Omega^\infty \Omega_R(\mathcal{L})$ . By [Cal+20a, Proposition 1.3.11], the data of  $q$  is equivalent to the data of points in the lower left and upper right corner of the square

$$\begin{array}{ccc}
 \Omega(\mathcal{L}) & \longrightarrow & \text{hom}_R(\mathcal{L}, R^{\varphi C_2}) \ni \ell(q) \\
 \downarrow & & \downarrow \\
 b(q) \in \text{hom}_{R \otimes R}(\mathcal{L} \otimes \mathcal{L}, R)^{hC_2} & \longrightarrow & \text{hom}_R(\mathcal{L}, R^{tC_2})
 \end{array} \quad (4.3)$$

and a path between their images in the lower right corner. In particular, the adjoint of  $b(q)$  must define a nondegenerate hermitian form on  $\mathcal{L}$ , that is, an equivalence  $\mathcal{L} \simeq \text{hom}_R(\mathcal{L}, R^*)$  where  $R^*$  is considered as an  $R$ -module via the action of the generator of  $C_2$ .

Write  $(\mathcal{L}^\vee, q^\vee)$  is for the inverse of  $(\mathcal{L}, q)$ . By definition of invertibility, there exists an  $R$ -linear map  $\ell(q^\vee): \mathcal{L}^\vee \rightarrow R^{\varphi C_2}$  so that the following diagram commutes

$$\begin{array}{ccc} \mathcal{L} \otimes_R \mathcal{L}^\vee & \xrightarrow{\ell(q) \otimes \ell(q^\vee)} & R^{\varphi C_2} \otimes_R R^{\varphi C_2} \\ \downarrow \sim \text{ev} & & \downarrow \text{multiplication} \\ R & \xrightarrow{\text{given}} & N_R \end{array} \quad (4.4)$$

**Lemma 4.5.** *Let  $(R, \mathfrak{Q})$  be a connective Poincaré ring spectrum. Then, for any integer  $n$ , the spectrum  $\mathfrak{Q}(\Sigma^n R)$  is  $(-2n)$ -connective.*

*Proof.* This follows from the fiber sequence

$$(\Sigma^{-2n} R)_{hC_2} \rightarrow \mathfrak{Q}(\Sigma^n R) \rightarrow \text{hom}_R(\Sigma^n R, C) \simeq \Sigma^{-n} C.$$

□

**Remark 4.6.** The functor  $\text{Pic}^P : \text{CAlg}^P \rightarrow \text{CAlg}^{\text{gp}}(\mathcal{S})$  preserves certain structures. Let  $A$  be a Poincaré ring spectrum. Since  $A$  is a module over  $(\mathbb{S}, \mathfrak{Q}_u)$ , the space  $\text{Pic}^P(A)$  is something over  $\text{Pic}^P(\mathbb{S}, \mathfrak{Q}_u)$ .

Since the forgetful functor  $\text{Pn}(A) \rightarrow \text{Mod}_A^\omega$  is symmetric monoidal we get an induced map

$$U : \text{Pic}^P(A) \rightarrow \text{Pic}(A)$$

of spectra. For a point  $(\mathcal{L}, q) \in \pi_0(\text{Pic}^P(A))$  we will refer to  $\mathcal{L} := U(\mathcal{L}, q)$  as the *underlying invertible module*. Note that the  $A$ -module  $A^*$  is (nonequivariantly) isomorphic to  $A$  via the involution, and so the fact that  $\mathcal{L} \simeq \text{hom}_A(\mathcal{L}, A^*)$  forces  $\mathcal{L}$  to be 2-torsion. In particular we get a refined map

$$U : \text{Pic}^P(A) \rightarrow \text{Pic}(A)[2]$$

which factors the underlying invertible module map.

**Example 4.7.** Let  $(\mathbb{S}, \mathfrak{Q}_u)$  be the universal Poincaré ring spectrum from Example 3.10. The only 2-torsion element of  $\text{Pic}(\mathbb{S}) \simeq \mathbf{Z}$  is  $\mathbb{S}$ . Therefore, any element in  $\text{Pic}^P(\mathbb{S}, \mathfrak{Q}_u)$  lies above  $\mathbb{S}$  under  $U$ . With Remark 3.12, we conclude  $\pi_0(\text{Pic}^P(\mathbb{S}, \mathfrak{Q}_u)) \simeq \pi_0(\mathbb{S}_{hC_2} \oplus \mathbb{S}^\times)^\times \simeq (\mathbf{Z} \times \mathbf{Z}/2)^\times \simeq \mathbf{Z}/2 \times \mathbf{Z}/2$ .

**Remark 4.8.** One might hope that the map  $\text{Pic}^P(A) \rightarrow \text{Pic}(A)[2]$  is close to an equivalence. This however is quite far from being true. Let  $k$  be a finite field of characteristic 2, and let  $\mathbb{S}_{W(k)}$  be the spherical Witt vectors on  $k$  in the sense of [Lur18, Example 5.2.7]. Then by [Nik23, Example 3.4] we know that  $\mathbb{S}_{W(k)}$  must satisfy that the map  $\varphi_2 : \mathbb{S}_{W(k)} \rightarrow \mathbb{S}_{W(k)}^{tC_2}$  is an equivalence where the action is trivial.

Consider now the Poincaré ring  $(\text{Mod}_{\mathbb{S}_{W(k)}}^\omega, \mathfrak{Q}_{\mathbb{S}_{W(k)}}^u)$  where  $\mathfrak{Q}_{\mathbb{S}_{W(k)}}^u$  is the Tate Poincaré structure. We have that  $\pi_0(\text{Pic}(\mathbb{S}_{W(k)})) \cong \mathbf{Z}$  and is generated by  $\Sigma \mathbb{S}_{W(k)}$ . To see this note that for  $\mathcal{L}$  an invertible module over  $\mathbb{S}_{W(k)}$ ,  $\mathcal{L}$  must be bounded below since otherwise it would not be perfect. Then for  $\pi_n(\mathcal{L})$  its bottom homotopy group, we have that  $\pi_n(\mathcal{L}/2) \cong k$  since it must be an invertible  $k$ -module and  $k$  is a field. Thus we get a map  $\mathbb{S}^n \rightarrow \mathcal{L}$  lifting a generator of  $k$ , and by adjunction an  $\mathbb{S}_{W(k)}$ -module map  $\Sigma^n \mathbb{S}_{W(k)} \rightarrow \mathcal{L}$  which on  $\pi_n((-)/2)$  gives an isomorphism  $k \cong k$ . Therefore

$$\mathbb{S}_{W(k)}[n] \otimes k \simeq k[n] \rightarrow k[n] \simeq \mathcal{L} \otimes k$$

is an equivalence, where the equivalence  $k[n] \simeq \mathcal{L} \otimes k$  follows from the fact that base change preserves invertible objects. The map  $\mathbb{S}_{W(k)}[n] \rightarrow \mathcal{L}$  is then a  $k$ -local, and therefore an  $\mathbb{F}_p$ -local, equivalence. Both sides are connective and  $p$ -complete so it follows that the map  $\mathbb{S}_{W(k)}[n] \rightarrow \mathcal{L}$  is an equivalence.

Thus  $\pi_0(\text{Pic}(\mathbb{S}_{W(k)})) = 0$ . On the other hand, we have that the unit map  $\mathbb{S}_{W(k)} \rightarrow \mathfrak{Q}_{\mathbb{S}_{W(k)}}^u(\mathbb{S}_{W(k)})$  is split by the map  $\mathfrak{Q}_{\mathbb{S}_{W(k)}}^u(\mathbb{S}_{W(k)}) \rightarrow \mathbb{S}_{W(k)}^{\varphi C_2} = \mathbb{S}_{W(k)}$ . Consequently  $\pi_0(\mathfrak{Q}_{\mathbb{S}_{W(k)}}^u(\mathbb{S}_{W(k)})) \cong \pi_0(\mathbb{S}_{W(k)} \oplus (\mathbb{S}_{W(k)})_{hC_2}) \cong$

L: add equivariance/symmetry data

V: define connectivity and make conditions here precise. As stated this works for R and C connective. More precisely,  $\text{conn}(\mathfrak{Q}(\Sigma^n R)) \min(\text{conn}(\Sigma^{-$

V: write out details

V: there is no truth in here yet. Work in progress. Noah had an example using Witt vectors which showed that  $\pi_0$  does not need to be 2-torsion

V: did we mod out by isomorphisms here?

N: There is probably a reference for this fact, I'll look around for one.



$W(k) \times W(k)$ . As a ring this is  $W_2(W(k))$ , and in order for  $q \in W_2(W(k))$  to induce a Poincaré structure we must have that  $q \in W_2(W(k))^\times \cong W(k)^\times \times W(k)^\times$ .

We then have that  $\pi_0(\text{Pic}^P(\mathbb{S}_{W(k)})) \cong W(k)^\times \times W(k)^\times / H$  where  $H$  is the subgroup of Poincaré structures  $q$  on  $\mathbb{S}_{W(k)}$  which are identified by some automorphism  $f : \mathbb{S}_{W(k)} \rightarrow \mathbb{S}_{W(k)}$ . By the defining property of spherical Witt vectors there is an equivalence  $\text{Maps}_{\text{CAlg}}(\mathbb{S}_{W(k)}, \mathbb{S}_{W(k)}) \simeq \text{Maps}_{\text{Perf}}(k, k) = \text{Gal}(k/\mathbb{F}_2)$  and the action on  $W(k)^\times \times W(k)^\times$  is given by  $g \in \text{Gal}(k/\mathbb{F}_2)$  acts via  $W(g) \times W(g)$ . Consequently

$$\pi_0(\text{Pic}^P(\mathbb{S}_{W(k)})) \cong (W(k)^\times \times W(k)^\times) / \text{Gal}(k/\mathbb{F}_2)$$

which even for  $k = \mathbb{F}_2$  is not zero and in fact not even  $2^\infty$ -torsion.

In the usual Picard spectrum one has the relationship  $\text{Pic} = B\mathbb{G}_m$ , where  $\mathbb{G}_m$  is the spectral algebraic group scheme sending a ring spectrum  $E$  to the spectrum of  $E$ -linear equivalences of  $E \text{ gl}_1 E := \text{Aut}_E(E)$ .<sup>1</sup> Equivalently  $\mathbb{G}_m$  is the affine groupscheme given by  $\mathbb{G}_m = \text{Spét}(\mathbb{S}\{x^{\pm 1}\})$ , where  $\mathbb{S}\{x^{\pm 1}\}$  is the free  $\mathbb{E}_\infty$  ring on the  $\mathbb{E}_\infty$  space  $\mathbb{Z}$ . This relationship between  $\text{Pic}$  and  $\mathbb{G}_m$  has many important applications, for example relating the higher homotopy groups of  $\text{Pic}(A)$  with those of  $A$ . We will spend the rest of this section on establishing such an equivalence in the Poincaré setting.

**Construction 4.9.** The underlying  $\mathbb{E}_\infty$  ring of  $\mathbb{G}_m^\Omega$  will again be  $\mathbb{S}\{x^{\pm 1}\}$ , but in order to promote this ring to a Poincaré ring it will be helpful to write it as

$$\mathbb{S}\{x^{\pm 1}, y^{\pm 1}\} \otimes_{\mathbb{S}\{z^{\pm 1}\}} \mathbb{S}$$

where the map  $\mathbb{S}\{z^{\pm 1}\} \rightarrow \mathbb{S}\{x^{\pm 1}, y^{\pm 1}\}$  is induced by  $z \mapsto xy$ . This ring naturally lifts to a Borel  $C_2$ -ring given by  $C_2$  swaps  $x$  and  $y$  and does nothing to  $z$ . Now take  $\mathbb{G}_m^\Omega$  to be the Poincaré ring with underlying Borel  $C_2$  structure as described above and geometric fixed points  $(\mathbb{G}_m^\Omega)^{\varphi_{C_2}} = \mathbb{S}$  and the map  $(\mathbb{G}_m^\Omega)^{\varphi_{C_2}} \rightarrow (\mathbb{G}_m^\Omega)^{t_{C_2}}$  given by the unit map. Endowing  $(\mathbb{G}_m^\Omega)^{\varphi_{C_2}}$  with the  $\mathbb{G}_m^\Omega$ -module structure given by  $x, y \mapsto 1$ , it remains to show that the unit map  $(\mathbb{G}_m^\Omega)^{\varphi_{C_2}} \rightarrow (\mathbb{G}_m^\Omega)^{t_{C_2}}$  factors the Tate valued Frobenius  $\mathbb{G}_m^\Omega \rightarrow (\mathbb{G}_m^\Omega)^{t_{C_2}}$  in order to promote  $\mathbb{G}_m^\Omega$  to a Poincaré ring.

By construction of  $\mathbb{G}_m^\Omega$  this amounts to showing that on  $\pi_0$  the Tate valued Frobenius sends  $x, y \mapsto 1$  in  $\pi_0((\mathbb{G}_m^\Omega)^{t_{C_2}})$ . This map sends both  $x$  and  $y$  to  $xy \in \pi_0((\mathbb{G}_m^\Omega)^{t_{C_2}})$ . These are equal to 1 in  $\pi_0((\mathbb{G}_m^\Omega)^{t_{C_2}})$  since the functor  $(-)^{t_{C_2}}$  is lax-monoidal so  $(\mathbb{G}_m^\Omega)^{t_{C_2}}$  is a module over  $\mathbb{S}\{x^{\pm 1}, y^{\pm 1}\}^{t_{C_2}} \otimes_{\mathbb{S}\{z^{\pm 1}\}^{t_{C_2}}} \mathbb{S}^{t_{C_2}}$  which has the image of  $xy$  equal to 1.

**Theorem 4.10.** *There is a natural equivalence of*

$$\Omega \text{Pic}^P(-) \simeq \mathbb{G}_m^\Omega$$

*of functors on Poincaré rings.*

*Proof.* This amounts to identifying the space  $\text{Aut}_{\text{Pn}(\text{Mod}_A)}(A, u)$  functorially, where  $(A, u)$  is the Poincaré object  $A$  with bilinear form given by the unit map  $\mathbb{S} \rightarrow \mathcal{Q}_A(A)$ . Note that any automorphism of Hermetian objects will automatically be Poincaré and so we may instead describe the automorphisms as a Hermetian object. We then have that  $\text{He}(\text{Mod}_A) \rightarrow \text{Mod}_A$  is a cocartesian fibration by definition, and classified by the functor which takes a module  $M$  to the groupoid  $\Omega^\infty \mathcal{Q}_A(M)$ . Thus we get that  $\text{Aut}_{\text{He}(\text{Mod}_A)}((A, u))$  is exactly the fiber of the map

$$\text{Aut}_{\text{Mod}_A}(A) \rightarrow \mathcal{Q}_A(A)$$

or in other words an automorphism  $(A, u) \rightarrow (A, u)$  is the data of an automorphism  $a \in \text{Aut}(A)$  together with a path  $q : u \mapsto a^*u$  in  $\Omega^{\infty+1} \mathcal{Q}_A(A)$ .

There is a natural transformation  $\mathbb{G}_m^\Omega(-) \rightarrow \Omega \text{Pic}^P(-)$  given as follows: we get a map  $\mathbb{G}_m^\Omega((\text{Mod}_A, \mathcal{Q}_A)) \rightarrow \text{Aut}_A(A)$  given by forgetting the Poincaré structure everywhere, and so it is enough to see that on  $\pi_0$  the automorphisms of  $A$  coming from  $\mathbb{G}_m^\Omega$  preserve  $u$ . By using the linear and quadratic decomposition of  $\mathcal{Q}_A$ , for an element  $a \in \pi_0(A)^\times$  send  $u$  to  $u$  must be sent to  $1 \in \pi_0(A^{\varphi_{C_2}})$  and must act by 1 on  $A^{h_{C_2}}$ . By the following Lemma this second condition is equivalent to  $a\sigma(a) \in \pi_0(A)^\times$  being equal to 1, but then these two conditions are exactly describing a map out of  $\mathbb{G}_m^\Omega$  as desired.

<sup>1</sup>Normally the automorphism space of an object is only  $\mathbb{A}_\infty$ , but as the unit in a symmetric monoidal category, the automorphisms of  $E$  inherit a canonical and in fact functorial  $\mathbb{E}_\infty$  structure and this construction makes sense.



Consequently we have a comparison map  $\mathbb{G}_m^{\mathfrak{Q}}(\mathrm{Mod}_A, \mathfrak{Q}_A) \rightarrow \Omega \mathrm{Pic}^{\mathrm{P}}(\mathrm{Mod}_A, \mathfrak{Q}_A)$ , and the above argument in fact shows that this map is an equivalence on  $\pi_0$ . To finish the argument, note that the pushout description of  $\mathbb{G}_m^{\mathfrak{Q}}$  induces a pullback of mapping spaces

$$\begin{array}{ccc} \mathbb{G}_m^{\mathfrak{Q}}(\mathrm{Mod}_A, \mathfrak{Q}_A) & \longrightarrow & \mathrm{Maps}_{\mathrm{CAlg}(\mathrm{Sp}^{C_2})}(\mathbb{S}\{x^{\pm 1}, y^{\pm 1}\}, A) \simeq \mathrm{gl}_1(A) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \mathrm{Maps}_{\mathrm{CAlg}(\mathrm{Sp}^{C_2})}(\mathbb{S}\{z\}, A) \simeq \Omega^{\infty} \mathfrak{Q}_A(A) \end{array}$$

which finishes the proof.  $\square$

**Lemma 4.11.** *Let  $A \in \mathrm{CAlg}(\mathrm{Sp}^{BC_2})$  and  $s \in \pi_0(A)^{\times}$ . Then  $a\sigma(a) = 1$  in  $\pi_0(A)$  if and only if  $(a \otimes a)^*$  acts by 1 on  $\pi_0(A^{hC_2}) = \pi_0(\mathrm{Hom}_{A \otimes A}(A \otimes A, A)^{hC_2})$ .*

*Proof.* The only if direction follows from the fact that the evaluation map  $\mathrm{Hom}_{A \otimes A}(A \otimes A, A) \rightarrow A$  is an  $A \otimes A$ -module map. Now suppose that  $a\sigma(a) = 1$  in  $A$ . Then before taking homotopy fixed points the induced map  $a^* = id$  because  $A$  is  $\mathbb{E}_{\infty}$ .<sup>2</sup>  $\square$

## 4.1 prime factorization and the picard group of hearts

We establish a Poincaré analogue of Fausk's result which describes the picard group of the derived category of a scheme  $X$  in terms of connected components of  $X$  and the classical picard group of  $X$ .

Let  $X$  be a simplicial set. The set  $\pi_0(X)$  is the set of connected components of  $X$ , i.e. simplicial subsets which are connected and form  $X$  via a coproduct. In other words, the functor  $\pi_0$  records a unique and maximal decomposition of  $X$  into coproducts. To establish the result mentioned above, we study the dual analogue of connected components in the sense of a maximal decomposition of  $X$  into products.

**Definition 4.12.** Let  $X$  be a simplicial set. We call a map of simplicial sets  $X \rightarrow Y$  a factor of  $X$ , if there is an isomorphism  $X \simeq Y_{\times} Z$ , for some simplicial set  $Z$ , such that  $Y \times Z \simeq X \rightarrow Y$  is a structure map of the given product.

**Definition 4.13.** Let  $X$  be a simplicial set. We say that  $X$  is prime, or indecomposable, if it is nonempty and every factor of  $X$  is isomorphic to either  $\Delta^0$  or  $X$ . We let  $\mathrm{prim}(X)$  denote the set of prime factors of  $X$ .

**Proposition 4.14.** *Let  $X_{\bullet}$  be a simplicial set, then  $X_{\bullet}$  is the product of its prime factors.*

*Proof.*

**Proposition 4.15.** *Let  $X$  be a simplicial set. Then  $\mathrm{prim}(X) \simeq \mathrm{prim}(X^{\simeq})$ .*

*Proof.* Let  $f : X \rightarrow Y$  be a weak equivalence of simplicial sets. Then  $f^{\simeq} : X^{\simeq} \rightarrow Y^{\simeq}$  is a weak equivalence of spaces.

**Proposition 4.16.** *Let  $X$  be a simplicial set. Then  $\mathrm{prim}(X) \simeq \pi_0(\mathrm{Spec}(X))$ , where  $\mathrm{Spec}(X^{\simeq})$  is the Balmer spectrum of  $X$  with respect to the symmetric monoidal structure given by cartesian product.*

*Proof.*

**Remark 4.17.** Let  $R$  be a commutative ring. Then the scheme  $\mathrm{Spec}(R)$  is isomorphic to the Balmer spectrum of  $\mathrm{Mod}_R^{\heartsuit}$ . When we view  $R$  as a discrete simplicial set, we thus have

$$\mathrm{prim}(\mathrm{Mod}_R^{\heartsuit}) \simeq \pi_0(\mathrm{Spec}(R)).$$

**Definition 4.18.** Let  $X$  be a prime simplicial set. A  $c$ -structure on  $X$  is a map of simplicial sets  $X \rightarrow \mathbf{Z}$  satisfying (todo). Let  $Y$  be a simplicial set, then a  $c$ -structure on  $Y$  is a product of  $c$ -structures on each of its prime components. We write  $X_{\geq n}$  for the homotopy pullback of  $\mathbf{Z}_{\geq n}$  along  $c$ ,  $X_{\leq n}$  for the homotopy pullback of  $\mathbf{Z}_{\leq n}$  along  $c$ , and  $X^{\heartsuit}$  for the pullback of  $X_{\leq n}$  along  $X_{\geq n} \rightarrow X$ .

V: still in development

V:

V:

V:

V: does this need a proof?

V: when  $X$  is a stable infinity category and prime, then a  $c$ -structure should be a

**Theorem 4.19.** *Let  $X$  be a prime simplicial set and  $c : X \rightarrow \mathbf{Z}$  a  $c$ -structure. Then we have a fiber sequence of monoids  $>$*

$$X^\heartsuit \rightarrow X^\simeq \rightarrow \mathbf{Z}.$$

*Proof.* □

**Corollary 4.20** (Fausk). *Let  $R$  be a discrete ring. Then we have a short exact sequence:  $>$*

$$0 \rightarrow \mathrm{Pic}(\mathrm{Mod}(R)^\heartsuit) \rightarrow \pi_0(\mathrm{Pic}(\mathrm{Mod}(R))) \rightarrow H^0(\mathrm{Spec}(R); \mathbf{Z}) \rightarrow 0.$$

*Proof.* □

## 4.2 Hermitian line bundles

**Definition 4.21.** Let  $R$  be a commutative discrete ring with a  $C_2$ -action  $\sigma : R \rightarrow R$ . Write  $\sigma_* R$  for the  $R$ -module with underlying abelian group  $R$  and action  $r \cdot m = \sigma(r) \cdot m$ . Let  $M$  be an  $R$ -module. Define the *adjoint* of  $M$  to be the  $R$ -module  $M^\dagger := \mathrm{hom}_R(M, \sigma_* R)$ . Also recall that there is a canonical  $R$ -linear isomorphism  $(M^\dagger)^\dagger \simeq M$ . Note that given two  $R$ -modules  $M, N$ , the adjoint satisfies  $M^\dagger \otimes N^\dagger \simeq (M \otimes N)^\dagger$ . Let  $I$  be a projective  $R$ -module (in particular, there is a canonical identification  $(I^\dagger)^\dagger \simeq I$ ). A  $\sigma$ -hermitian form on  $I$  is an  $R$ -linear isomorphism  $\varphi : I \xrightarrow{\sim} I^\dagger$  so that  $\varphi^\dagger = \varphi$ .

**Observation 4.22.** Let  $R$  be a commutative discrete ring with a  $C_2$ -action  $\sigma : R \rightarrow R$ . Given two discrete  $R$ -modules  $M, N$  equipped with  $\sigma$ -hermitian forms  $\varphi, \psi$ , respectively,  $\varphi \otimes \psi$  defines a  $\sigma$ -hermitian form on  $M \otimes_R N$ . Using the canonical isomorphism mentioned above, if  $\varphi$  is a  $\sigma$ -hermitian form on  $M$ , then  $\varphi^\dagger$  induces a  $\sigma$ -hermitian form on  $M^\dagger$ . Finally, observe that  $R$  has a canonical  $\sigma$ -hermitian form which is the adjoint of the map  $R \otimes R \rightarrow R, r \otimes s \mapsto r\sigma(s)$ .

**Definition 4.23.** Let  $R$  be a commutative discrete ring with a  $C_2$ -action  $\sigma : R \rightarrow R$ . Define the *hermitian Picard group of  $R$*  to have underlying set consisting of pairs  $(I, \varphi)$  where  $I$  is an invertible  $R$ -module and  $\varphi$  is a  $\sigma$ -hermitian form on  $I$ .

By Observation 4.22, this set inherits a group structure. We write  $\mathrm{hPic}(R)$  for the group of  $\sigma$ -hermitian line bundles on  $\mathrm{Spec} R$ .

**Theorem 4.24.** *Let  $R$  be a discrete commutative ring with a  $C_2$ -action  $\sigma : R \xrightarrow{\sim} R$  via ring maps. Regard  $R$  as a Poincaré ring via Example 3.15. Then there is a split short exact sequence of abelian groups*

$$0 \rightarrow \mathrm{hPic}(R) \rightarrow \pi_0 \mathrm{PnPic}(\underline{R}^\sigma) \rightarrow C_{C_2}(\mathrm{Spec} R, \mathbb{Z}^-) \rightarrow 0$$

where  $R$  is endowed with the genuine symmetric Poincaré structure and  $\mathbb{Z}^-$  is endowed with the  $C_2$ -action given by multiplication by  $-1$  and  $C_{C_2}$  denotes continuous functions which are moreover  $C_2$ -equivariant. Moreover, forgetting the hermitian form (resp. forgetting the  $C_2$ -action) induces a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{hPic}(R) & \longrightarrow & \pi_0 \mathrm{PnPic}(R) & \longrightarrow & C_{C_2}(\mathrm{Spec} R, \mathbb{Z}^-) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{Pic}^{\mathrm{cl}}(R) & \longrightarrow & \pi_0 \mathrm{Pic}(\mathrm{Perf}_R) & \longrightarrow & C(\mathrm{Spec} R, \mathbb{Z}) \longrightarrow 0 \end{array}$$

where the bottom row is that of [Fau03, Theorem 3.5].

*Proof.* An object of  $\pi_0 \mathrm{PnPic}(R)$  is a pair  $(I, q)$  where  $I$  is an invertible  $R$ -module and  $q$  is a point in  $\pi_0 \Omega^\infty \mathcal{Q}_{R^{\mathrm{gs}}}(I)$ . By the proof of [Fau03, Theorem 3.5],  $I$  induces a continuous map  $\Psi(I) : \mathrm{Spec} R \rightarrow \mathbb{Z}$ . Write  $\sigma$  for the involution on  $R$ . Now  $q$  in particular induces an equivalence  $q : I \xrightarrow{\sim} I^\dagger \simeq (\sigma_* I)^\vee$ . For each point  $\mathfrak{p} \in \mathrm{Spec} R$ , localizing  $q$  gives an equivalence

$$q_{\mathfrak{p}} : I_{\mathfrak{p}} \xrightarrow{\sim} (\sigma_* I)_{\mathfrak{p}}^\vee \simeq (\sigma_*(I_{\sigma(\mathfrak{p})}))^\vee.$$

<sup>2</sup>Or just  $\mathbb{E}_2$ .

V:

V: what

V: apply pic to the previous sequence and take  $\pi_0$

L: see 3.8-3.11 in this paper

L: workshop the name later

Since  $I_{\mathfrak{p}}$  is an invertible module over a local ring, [Fau03, Proposition 3.2] implies that  $q_{\mathfrak{p}}$  induces an equivalence

$$I_{\mathfrak{p}} \simeq R_{\mathfrak{p}}[\varphi(\mathfrak{p})] \xrightarrow{\sim} (\sigma_*(R_{\sigma(\mathfrak{p})}[\varphi(\sigma(\mathfrak{p}))]))^{\vee} \simeq (\sigma_*(R_{\sigma(\mathfrak{p})}))^{\vee}[-\varphi(\sigma(\mathfrak{p}))].$$

Since  $R$  is discrete, this implies in particular that  $\Psi(I)(\sigma(\mathfrak{p})) = -\Psi(I)(\mathfrak{p})$ , i.e. that  $\Psi(I)$  is  $C_2$ -equivariant. It follows immediately from [Fau03, Theorem 3.5] that  $\Psi$  is a homomorphism and that an element of the kernel of  $\Psi$  lifts to  $\mathrm{hPic}(R)$ .

Now consider a  $C_2$ -equivariant map  $g: \mathrm{Spec} R \rightarrow \mathbb{Z}$ . As in *loc. cit.*, the image of  $g$  is finite and  $C_2$ -invariant, say  $\{n_1, -n_1, \dots, n_m, -n_m\}$  or  $\{0, n_1, -n_1, \dots, n_m, -n_m\}$  for some  $n_i \neq 0$ . As in *loc. cit.*, the disjoint subsets  $U_{\pm n_i} := g^{-1}(\pm n_i)$  correspond to an orthogonal basis of idempotents  $e_{U_{\pm n_i}}$  in  $R$ . Since  $g$  is  $C_2$ -equivariant with respect to the sign action on  $\mathbb{Z}$ , we have  $\sigma(U_{n_i}) = U_{-n_i}$ . Moreover, it follows from Lemma 3.4 *ibid.* that  $\sigma(e_{U_{n_i}}) = e_{U_{-n_i}}$ . Consider the  $R$ -module  $\Phi(g) := \bigoplus_{n \in \mathrm{Im}(g)} e_{g^{-1}(\{n\})} R[n]$ . In other words,  $\Phi(g) := \bigoplus_{i=1}^m (e_{U_{n_i}} R[n_i] \oplus e_{U_{-n_i}} R[-n_i])$  if 0 is not in the image of  $g$  and  $\Phi(g) := e_{U_0} \oplus \bigoplus_{i=1}^m (e_{U_{n_i}} R[n_i] \oplus e_{U_{-n_i}} R[-n_i])$  otherwise. Observe that  $(e_{U_{-n_i}} R[-n_i])^{\dagger} = \mathrm{hom}_R(e_{U_{-n_i}} R[-n_i], \sigma_* R) = \mathrm{hom}_R(\sigma_*(e_{U_{-n_i}} R), R)[n_i] = \mathrm{hom}_R(e_{U_{n_i}} R, R)[n_i]$ . Finally, we claim that there is a canonical  $\sigma$ -hermitian form  $q_g \in \Omega^{\infty} \mathfrak{Q}_{R^{\mathrm{ss}}}(\Phi(g))$  whose adjoint  $q_g^{\dagger}: \Phi(g) \xrightarrow{\sim} \Phi(g)^{\dagger}$  corresponds to the identity. That  $q_g$  defines a point of  $\mathrm{hom}_{R^{\otimes 2}}(\Phi(g)^{\otimes 2}, R)^{hC_2}$  is evident. Observe that to give a lift of  $q_g$  to  $\mathfrak{Q}_{R^{\mathrm{ss}}}(\Phi(g)) = \mathrm{hom}_{N^{C_2}R}(N^{C_2}\Phi(g), R)$  is equivalent to giving a commutative diagram

$$\begin{array}{ccc} \Phi(g) \otimes_R R^{\varphi C_2} & \overset{\exists?}{\dashrightarrow} & R^{\varphi C_2} \\ \downarrow & & \downarrow \\ (\Phi(g)^{\otimes 2})^{tC_2} & \xrightarrow{q_g^{tC_2}} & R^{tC_2} \end{array} \quad (4.25)$$

Let us write  $\eta: R \rightarrow \pi_0 R^{\varphi C_2}$  for the ring map induced by the structure map. Since  $R$  is a  $C_2$ - $\mathbb{E}_{\infty}$ -ring,  $\eta$  is invariant with respect to the given action on  $R$  and the trivial action on  $\pi_0 R^{\varphi C_2}$ . Consider  $e_{U_{n_i}}$  an idempotent corresponding to an element of the image of  $g$  so that  $n_i \neq 0$ . Then

$$\begin{aligned} \eta(e_{U_{n_i}}) &= \eta(e_{U_{n_i}})^2 && \text{ring maps preserve idempotents} \\ &= \eta(e_{U_{n_i}}) \cdot \eta(e_{U_{-n_i}}) && C_2\text{-invariance of } \eta \\ &= \eta(e_{U_{n_i}} e_{U_{-n_i}}) && \eta \text{ is a ring map} \\ &= 0 && \text{orthogonality and } n_i \neq 0. \end{aligned}$$

In particular, if 0 is not in the image of  $g$ ,  $\Phi(g) \otimes_R R^{\varphi C_2} \simeq 0$  and (4.25) commutes vacuously. If 0 is in the image of  $g$ , then  $e_{U_0} R$  is a discrete/projective  $e_{U_0} R$ -module and  $q_g$  evidently defines a genuine hermitian form on  $e_{U_0} R$  (compare [Cal+20a, Remark 4.2.21]).

Thus,  $g \mapsto (\Phi(g), q_g)$  defines a splitting of  $\Psi$  which agrees with the splitting constructed in [Fau03, Theorem 3.5] on underlying objects.  $\square$

## 5 Poincaré structures on schemes with involution

Let  $X$  be a scheme with an involution  $\sigma: X \xrightarrow{\sim} X$ . We want to introduce a Poincaré structure  $\mathfrak{Q}$  on  $\mathrm{Perf}(X)$  so that the duality is given by  $E \mapsto E^{\vee} \otimes \sigma_* (\mathcal{O}_X)$  (contrast with §3 of [this paper](#)).

### 5.1 Poincaré structures associated to rigid symmetric monoidal $\infty$ -categories with involution

Let  $\mathcal{C}$  be a stably symmetric monoidal  $\infty$ -category with an involution, i.e. an exact autoequivalence  $\sigma: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$  and a functor  $BC_2 \rightarrow \mathbb{E}_{\infty} \mathrm{Alg} \mathcal{C}^{\mathrm{ex}}$  sending  $*$  to  $\mathcal{C}$  and a generator of  $\mathrm{End}_{BC_2}(*) \simeq C_2$  to  $\sigma$ . Then  $\sigma$  induces a  $C_2$ -action on the  $\infty$ -groupoid of  $\infty$ -invertible objects  $\mathrm{Pic}(\mathcal{C})$ . Let  $L \in \mathrm{Pic}(\mathcal{C})^{hC_2}$  be a homotopy fixed point of this action. In other words,  $L$  is endowed with the choice of an equivalence  $\varphi: L \simeq \sigma(L)$ , a homotopy from  $\sigma(\varphi) \circ \varphi$  to the identity on  $L$ , and higher coherences.

Consider a functor  $f: \mathcal{C}^{\text{op}} \rightarrow \text{Sp}$  which is  $C_2$ -equivariant with respect to the  $\sigma$ -action on  $\mathcal{C}$  and the trivial action on  $\text{Sp}$ . In particular, the “ $C_2$ -equivariance” of  $f$  is additional data: for each  $x \in \mathcal{C}$ , an equivalence of spectra  $c_x: f(x) \simeq f(\sigma(x))$  which is natural in  $x$ , a homotopy from  $c_{\sigma(x)} \circ c_x$  to  $\text{id}_x$ , and higher coherences.

**Lemma 5.1.** *Let  $\mathcal{C}$  be a stably symmetric monoidal  $\infty$ -category with an involution, and suppose given  $L \in \text{Pic}(\mathcal{C})^{hC_2}$  a homotopy fixed point of this action. Then the functor  $\text{hom}_{\mathcal{C}}(-, L)$  promotes canonically to a  $C_2$ -equivariant functor  $\mathcal{C}^{\text{op}} \rightarrow \text{Sp}$  in the sense of the previous paragraph.*

*Proof.* Note that the Yoneda embedding  $y: \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \text{Sp})$  is equivariant with respect to the given action on  $\mathcal{C}$  and the action of  $C_2$  on the functor category via  $F \mapsto \sigma^* F = F \circ \sigma$ . Now since  $\text{Pic}(\mathcal{C}) \subseteq \mathcal{C}$  induces  $\text{Pic}(\mathcal{C})^{hC_2} \rightarrow \mathcal{C}^{hC_2}$ , we may take the image of  $L$  under the Yoneda embedding:  $y(L) \in \text{Fun}(\mathcal{C}^{\text{op}}, \text{Sp})^{hC_2} \simeq \text{Fun}_{C_2}(\mathcal{C}^{\text{op}}, \text{Sp})$ .  $\square$

Now fix a presentably symmetric monoidal  $\infty$ -category  $\mathcal{D}$ , and regard it as having the trivial  $C_2$ -action. Recall the  $\text{Fin}_*$ -cartesian fibration  $(\text{Cat}_{\text{op}/q^*\mathcal{D}}^{BC_2})^{\otimes} \simeq ((\text{Cat}_{\text{op}/\mathcal{D}})^{BC_2})^{\otimes} \rightarrow (\text{Cat}^{BC_2})^{\times}$  from [CHN24, p. 13]. Set

$$\mathcal{W}_{\mathcal{D}}^{\otimes} := \mathbb{E}_{\infty} \text{Alg}(\text{Cat}^{BC_2})^{\times} \times_{(\text{Cat}^{BC_2})^{\times}} (\text{Cat}_{\text{op}/q^*\mathcal{D}}^{BC_2})^{\otimes}$$

This is a  $\text{Fin}_*$ -cartesian fibration classified by the lax symmetric monoidal functor

$$\begin{aligned} \mathbb{E}_{\infty} \text{Alg}(\text{Cat})^{BC_2} &\rightarrow \text{Cat} \\ \mathcal{C}^{\otimes} &\mapsto \text{Fun}_{C_2}(\mathcal{C}^{\text{op}}, q^*\mathcal{D}). \end{aligned} \tag{5.2}$$

In particular, an object of the underlying  $\infty$ -category of  $\mathcal{W}_{\mathcal{D}}^{\otimes}$  is a pair  $(\mathcal{C}, f)$  where  $\mathcal{C}$  is a symmetric monoidal  $\infty$ -category with a  $C_2$ -action (via a symmetric monoidal functor) and  $f: \mathcal{C}^{\text{op}} \rightarrow q^*\mathcal{D}$  is a  $C_2$ -equivariant functor.

**Construction 5.3.** Given a  $C_2$ -equivariant functor  $f: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$ , we may regard the data of the  $C_2$ -equivariance of  $f$  as a commutative diagram

$$\begin{array}{ccc} \widetilde{\mathcal{C}^{\text{op}}} & \xrightarrow{\tilde{f}} & \mathcal{D} \times BC_2 \\ \downarrow & & \downarrow \\ BC_2 & \xlongequal{\quad} & BC_2 \end{array}$$

where the vertical maps are cocartesian fibrations and the restriction of  $\tilde{f}$  to the fiber over the point  $*$  in  $BC_2$  recovers  $f$ . The diagram induces a map on cocartesian sections

$$\overline{f}: \text{Fun}_{BC_2}^{\text{cocart}}(BC_2, \widetilde{\mathcal{C}^{\text{op}}}) \rightarrow \mathcal{D}^{BC_2}.$$

Now if  $\mathcal{C}$  is a symmetric monoidal  $\infty$ -category, we can associate to  $f$  the composite

$$T_f: \mathcal{C}^{\text{op}} \xrightarrow{x \mapsto x \otimes \sigma(x)} \text{Fun}_{BC_2}^{\text{cocart}}(BC_2, \widetilde{\mathcal{C}^{\text{op}}}) \xrightarrow{\overline{f}} \mathcal{D}^{BC_2}.$$

Finally, if  $\mathcal{D}$  admits  $BC_2$ -indexed limits, we can take homotopy fixed points of  $C_2$ -objects in which case we define the functor  $\mathcal{Q}_f^s: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$  as the composite

$$\mathcal{Q}_f^s: \mathcal{C}^{\text{op}} \xrightarrow{T_f} \mathcal{D}^{BC_2} \xrightarrow{(-)^{hC_2}} \mathcal{D}.$$

**Observation 5.4.** Let  $\mathcal{C}$ ,  $L$ , be as before. Then the cross effect  $B_L$  of  $\mathcal{Q}_L^s$  is given by  $B_L(x, y) = \text{hom}_{\mathcal{C}}(x \otimes \sigma(y), L)$ .

**Proposition 5.5.** *Let  $\mathcal{D}$  be a symmetric monoidal  $\infty$ -category which admits  $BC_2$ -indexed limits; endow  $\mathcal{D}$  with the trivial  $C_2$ -action. Then the assignment  $(\mathcal{C}, f) \mapsto (\mathcal{C}, \mathcal{Q}_f^s)$  of Construction 5.3 assembles to form a lax symmetric monoidal functor*

$$\mathcal{W}_{\mathcal{D}}^{\otimes} \rightarrow (\text{Cat}^{BC_2})_{\text{op}/\mathcal{D}}^{\otimes}$$

L: This may be ‘overkill,’ but this is true because  $\mathcal{C}$  can be regarded as a  $\mathcal{O}_{C_2}^{\text{op}}$ -parametrized  $\infty$ -category (with empty fiber over  $C_2/C_2$ ). Then there is a parametrized Yoneda embedding.

L: I’m just using the same notation as Harpaz-Nardin-Shah here, but  $(-)^s$  is maybe a little weird because it should be ‘hermitian,’ not ‘symmetric.’

sitting in a commutative diagram

$$\begin{array}{ccc}
\mathcal{W}_{\mathcal{D}}^{\otimes} & \longrightarrow & (\text{Cat}^{BC_2})_{\text{op}/\mathcal{D}}^{\otimes} \\
\downarrow & & \downarrow \\
\mathbb{E}_{\infty}\text{Alg}(\text{Cat}^{BC_2})^{\times} & \xrightarrow{\mathcal{C}^{\otimes} \mapsto \mathcal{C}} & (\text{Cat}^{BC_2})^{\times}
\end{array} \tag{5.6}$$

in which both vertical arrows are both  $\text{Fin}_*$ -cartesian fibrations and cocartesian fibrations of  $\infty$ -operads.

**Construction 5.7.** Consider the functors  $r: \mathbb{E}_{\infty}\text{Alg}(\text{Cat}^{BC_2}) \xrightarrow{\mathcal{C}^{\otimes} \mapsto \mathcal{C}} \text{Cat}^{BC_2}$  and  $p: \mathbb{E}_{\infty}\text{Alg}(\text{Cat}^{BC_2}) \xrightarrow{\text{forget}} \text{Cat} \xrightarrow{q^*} \text{Cat}^{BC_2}$  where  $q: BC_2 \rightarrow *$ . We construct a symmetric monoidal natural transformation  $\tau: r^{\times} \Rightarrow p^{\times}$  whose component at a given category  $\mathcal{C}$  with  $C_2$ -action  $\sigma: \mathcal{C} \simeq \mathcal{C}$  is the  $C_2$ -equivariant functor  $\mathcal{C} \xrightarrow{x \mapsto x \otimes \sigma(x)} p(\mathcal{C})$ .

Let  $\text{Span}(\text{Fin}_{C_2}^{\text{free}})$  be the span  $\infty$ -category of finite sets with free  $C_2$ -action. For an  $\infty$ -category with finite products  $\mathcal{E}$ , there is a natural equivalence  $\text{Fun}^{\times}(\text{Span}(\text{Fin}_{C_2}^{\text{free}}), \mathcal{E}) \simeq \text{CMon}(\mathcal{E})^{BC_2}$  between product-preserving functors  $\text{Span}(\text{Fin}_{C_2}^{\text{free}}) \rightarrow \mathcal{E}$  and commutative monoids in  $\mathcal{E}$  with  $C_2$ -action. Taking  $\mathcal{E} \simeq \text{Cat}$ , we may identify the functor  $r$  as restriction along the inclusion  $i: BC_2 \rightarrow \text{Span}(\text{Fin}_{C_2}^{\text{free}})$  of the maximal subgroupoid in the full subcategory on a finite  $C_2$ -set with a single orbit. On the other hand, we can identify the functor  $p$  as restriction along the map  $j: BC_2 \rightarrow \{*\} \xrightarrow{* \mapsto C_2} \text{Span}(\text{Fin}_{C_2}^{\text{free}})$ . Now the span  $C_2 \xleftarrow{\pi_1} C_2 \times C_2 \xrightarrow{\pi_2} C_2$  determines a morphism in  $\text{Span}(\text{Fin}_{C_2}^{\text{free}})$  which is equivariant with respect to the given action on the source  $C_2$  and the *trivial action* on the target  $C_2$ . This morphism determines a functor  $\Delta^1 \times BC_2 \rightarrow \text{Span}(\text{Fin}_{C_2}^{\text{free}})$  whose restriction to  $\{0\} \times BC_2$  agrees with  $i$  and whose restriction to  $\{1\} \times BC_2$  agrees with  $j$ . This determines a natural transformation  $i^* \Rightarrow j^*$  of functors  $\text{Fun}(\text{Span}(\text{Fin}_{C_2}^{\text{free}}), \text{Cat}) \rightarrow \text{Fun}(BC_2, \text{Cat})$ , and precomposing with the inclusion  $\text{Fun}^{\times}(\text{Span}(\text{Fin}_{C_2}^{\text{free}}), \text{Cat}) \subset \text{Fun}(\text{Span}(\text{Fin}_{C_2}^{\text{free}}), \text{Cat})$  gives the desired natural transformation  $\tau: r \Rightarrow p$ . Since  $r$  and  $p$  preserve products, we may lift them to symmetric monoidal functors  $r^{\times}, p^{\times}: \mathbb{E}_{\infty}\text{Alg}(\text{Cat}^{BC_2})^{\times} \rightarrow (\text{Cat}^{BC_2})^{\times}$ , and  $\tau$  refines to a symmetric monoidal natural transformation  $\tau^{\times}: r^{\times} \Rightarrow p^{\times}$ .

*Proof of Proposition 5.5.* Horizontally composing the natural transformation of Construction 5.7 with the functor 5.3 and unstraightening induces a commutative diagram

$$\begin{array}{ccc}
\mathcal{W}_{\mathcal{D}}^{\otimes} & \longrightarrow & (\text{Cat}_{\text{op}/\mathcal{D}}^{BC_2})^{\otimes} \\
\downarrow & & \downarrow \\
(\text{Cat}^{BC_2})_{\text{op}/\mathcal{D}}^{\otimes} & \xrightarrow{\mathcal{C}^{\otimes} \mapsto \mathcal{C}} & (\text{Cat}^{BC_2})^{\times}
\end{array}, \tag{5.8}$$

where we have used that  $(q^*\mathcal{D})^{hC_2} \simeq \mathcal{D}^{BC_2}$ . □

**Definition 5.9.** Define

$$\mathcal{W}_{\text{ex}}^{\otimes} \subseteq \mathcal{W}_{\text{Sp}}^{\otimes} \times_{\mathbb{E}_{\infty}\text{Alg}(\text{Cat})} \mathbb{E}_{\infty}\text{Alg}(\text{Cat}_{\infty}^{\text{ex}})$$

to be the full sub-operad on those colors  $(\mathcal{C}, f)$  so that  $f$  is exact.

**Observation 5.10.** The commutative square (5.6) restricts to a commutative square of  $\infty$ -operads

$$\begin{array}{ccc}
\mathcal{W}_{\text{ex}}^{\otimes} & \xrightarrow{(\mathcal{C}, f) \mapsto (\mathcal{C}, \Omega_f^{\circ})} & \text{Cat}_{\infty}^{\text{h}} \otimes \\
\downarrow & & \downarrow \\
(\mathbb{E}_{\infty}\text{Alg}(\text{Cat}_{\infty}^{\text{ex}})^{BC_2})^{\otimes} & \longrightarrow & \text{Cat}_{\infty}^{\text{ex}} \otimes
\end{array} \tag{5.11}$$

L: This is basically Construction 3.1.4 of [CHN24] with minor edits.

L: Pretty similar to proof of [CHN24, Proposition 3.1.3], the main thing is the fact (stated after proof of Lemma 3.1.1 of *op. cit.*) that  $\text{Fin}_*$ -cartesian fibrations are classified by lax symmetric monoidal functors.

where  $\mathcal{Y}_f^s$  was defined in Construction 5.3.

**Lemma 5.12.** *Both vertical maps in 5.11 are cocartesian fibrations of  $\infty$ -operads. In particular,  $\mathcal{W}_{\text{ex}}^{\otimes}$  is a symmetric monoidal  $\infty$ -category.*

*Proof.* That  $\text{Cat}_{\infty}^{\text{h}}{}^{\otimes}$  is symmetric monoidal and the right-hand projection is a symmetric monoidal functor is [Cal+20a, Theorem 5.2.7]. Furthermore, by Proposition 5.5 and base change, we have a cocartesian fibration of  $\infty$ -operads:

$$\pi: (\mathbb{E}_{\infty} \text{Alg}(\text{Cat}_{\infty}^{\text{ex}})^{BC_2})^{\otimes} \times_{(\text{Cat}^{BC_2})^{\times}} (\text{Cat}_{\text{op}/\mathcal{D}}^{BC_2})^{\otimes} \rightarrow (\mathbb{E}_{\infty} \text{Alg}(\text{Cat}_{\infty}^{\text{ex}})^{BC_2})^{\otimes}$$

so that  $\mathcal{W}_{\text{ex}}^{\otimes}$  includes as a full sub-operad on the fiber product on the left on those colors  $(\mathcal{C}, f: \mathcal{C}^{\text{op}} \rightarrow \text{Sp})$  so that  $f$  is exact. It suffices to show that if  $\alpha: (\mathcal{C}_i, f_i: \mathcal{C}_i^{\text{op}} \rightarrow \text{Sp})_{i \in I} \rightarrow (\mathcal{D}, f: \mathcal{D}^{\text{op}} \rightarrow \text{Sp})$  is a  $\pi$ -cocartesian arrow so that  $(\mathcal{C}_i, f_i: \mathcal{C}_i^{\text{op}} \rightarrow \text{Sp})_{i \in I}$  is in  $\mathcal{W}_{\text{ex}}^{\otimes}$ , then  $(\mathcal{D}, f: \mathcal{D}^{\text{op}} \rightarrow \text{Sp})$  is also in  $\mathcal{W}_{\text{ex}}^{\otimes}$ . This holds because the left Kan extension of the multi-exact functor  $\Pi_i f_i: \Pi_i \mathcal{C}_i^{\text{op}} \rightarrow \text{Sp}$  along  $\Pi_i \mathcal{C}_i^{\text{op}} \rightarrow \bigotimes_i \mathcal{C}_i^{\text{op}} \xrightarrow{\alpha} \mathcal{D}^{\text{op}}$  is exact.  $\square$

As in [CHN24, p. 15], we can identify objects of the underlying  $\infty$ -category of  $\mathcal{W}_{\text{ex}}^{\otimes}$  with pairs  $(\mathcal{C}^{\otimes}, \sigma_{\mathcal{C}}, L, \lambda)$  where  $\mathcal{C}$  is a symmetric monoidal stable  $\infty$ -category with involution  $\sigma_{\mathcal{C}}$ , and  $(L, \lambda) \in \text{Ind}(\mathcal{C})^{hC_2}$  is a fixed point with respect to the induced action on  $\text{Ind}(\mathcal{C})$ .

**Recollection 5.13.** A symmetric monoidal  $\infty$ -category  $\mathcal{C}$  is said to be *rigid* if every object in  $\mathcal{C}$  is dualizable.

**Proposition 5.14.** *Suppose  $\mathcal{C}$  is a rigid stably symmetric monoidal  $\infty$ -category with a  $C_2$ -action  $\sigma_{\mathcal{C}}$  via symmetric monoidal functors, and let  $(L, \lambda) \in \text{Ind}(\mathcal{C})^{BC_2}$ . Then the hermitian structure  $\mathcal{Y}_L^s$  is non-degenerate if and only if  $L$  belongs to  $\mathcal{C}$ , and it is furthermore Poincaré if and only if the underlying object  $L$  is tensor-invertible in  $\mathcal{C}$ . In addition, if  $g: \mathcal{C} \rightarrow \mathcal{C}'$  is a symmetric monoidal  $C_2$ -equivariant exact functor,  $(L, \lambda) \in \mathcal{C}^{BC_2}$  and  $(L', \lambda') \in (\mathcal{C}')^{BC_2}$  are tensor-invertible, and  $g(L, \lambda) \simeq (L', \lambda')$  an equivalence in  $(\mathcal{C}')^{BC_2}$ , then the induced hermitian functor  $(\mathcal{C}, \mathcal{Y}_L^s) \rightarrow (\mathcal{C}', \mathcal{Y}_{L'}^s)$  is Poincaré.*

*Proof.* If the bilinear part  $B_L$  is represented by  $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$ , then  $D(1_{\mathcal{C}}) = \sigma_{\mathcal{C}}(L)$ ; in particular,  $\sigma_{\mathcal{C}}(L) \simeq L$  is an object of  $\mathcal{C}$ . On the other hand, if  $L$  belongs to  $\mathcal{C}$ , then for  $x, y \in \mathcal{C}$ , the bilinear part  $B_L$  is  $B_L(x, y) = \text{hom}_{\text{Ind}(\mathcal{C})}(x \otimes \sigma_{\mathcal{C}}(y), L) \simeq \text{hom}_{\mathcal{C}}(x \otimes \sigma_{\mathcal{C}}(y), L) \simeq \text{hom}_{\mathcal{C}}(x, L \otimes \sigma_{\mathcal{C}}(y)^{\vee})$ . The natural transformation  $\text{id}_{\mathcal{C}} \rightarrow D^{\text{op}} \circ D$  is given by  $y \rightarrow L \otimes \sigma_{\mathcal{C}}(L \otimes \sigma_{\mathcal{C}}(y)^{\vee})^{\vee}$  induced by the adjoint of the equivalence  $L \simeq \sigma_{\mathcal{C}}(L)$ , which is an equivalence if and only if  $L$  is invertible.

Finally, the natural transformation  $g \circ D_L \Rightarrow D_{L'} \circ g$  is  $g(L \otimes \sigma_{\mathcal{C}}(y)^{\vee}) \simeq g(L) \otimes g(\sigma_{\mathcal{C}}(y)^{\vee}) \rightarrow L' \otimes \sigma_{\mathcal{C}'}(g(y))^{\vee}$  which is an equivalence if and only if the map  $g(L) \rightarrow L'$  is an equivalence.  $\square$

## 5.2 From schemes with involution to Poincaré structures on module categories

**Recollection 5.15** ([FW20, Remark 4.20]). Let  $X$  be a scheme with an involution  $\lambda: X \rightarrow X$ . A map  $\pi: X \rightarrow Y$  is called a good quotient of  $X$  relative to  $\lambda$  if  $\pi$  is  $C_2$ -invariant and affine and  $\pi_{\#}: \mathcal{O}_Y \rightarrow \pi_* \mathcal{O}_X$  induces an isomorphism  $\mathcal{O}_Y \simeq (\pi_* \mathcal{O}_X)^{C_2}$ . A good quotient of  $X$  exists if and only if every  $C_2$ -orbit is contained in an affine open subscheme, in which case it is unique up to isomorphism.

**Definition 5.16.** Define a category  $\text{qSch}^{C_2}$  so that

- an object of  $\text{qSch}^{C_2}$  consists of the data of qcqs schemes  $X$  and  $Y$ , an involution  $\lambda: X \rightarrow X$ , and a morphism  $p: X \rightarrow Y$  which exhibits  $Y$  as a *good quotient* of the involution on  $X$  in the sense of [FW20, Remark 4.20].
- a morphism from  $(X, \lambda, Y, p)$  to  $(Z, \nu, W, q)$  consists of a  $C_2$ -equivariant morphism  $X \rightarrow Z$  and a morphism  $Y \rightarrow W$  so that the diagram

$$\begin{array}{ccc} X & \longrightarrow & Z \\ \downarrow & & \downarrow \\ Y & \longrightarrow & W \end{array}$$

commutes.

L: reference for when derived categories of connective objects over more general objects are compactly generated

L: how much equivariance on  $g(L) \rightarrow L'$  is necessary?

L: explain why using good quotients/choice of quotient is useful.



**Remark 5.17.** Suppose  $(X, \lambda, Y, p)$  is an object of  $\text{qSch}^{C_2}$  and  $j: U \rightarrow Y$  is a flat map. Then  $(X_U, \lambda|_U, U, p|_U)$  is an object of  $\text{qSch}^{C_2}$ . Affineness and invariance under the  $C_2$ -action are stable under pullback, so it suffices to show that  $j^*(\pi)$  satisfies  $\mathcal{O}_U \simeq (j^*(\pi))_*(\mathcal{O}_{j^*U})^{C_2}$ . This follows from the proof of [FW20, Theorem 4.35(i)].

**Proposition 5.18.** Write  $U: \text{qSch}^{C_2} \rightarrow \text{qSch}$  for the functor so that  $U(X, \lambda, Y, p) = X$ . The category  $\text{qSch}^{C_2}$  has a symmetric monoidal structure  $\boxtimes$  so that  $U$  is symmetric monoidal, where  $\text{qSch}$  is endowed with the product symmetric monoidal structure.

*Proof.* If  $X, Z$  are schemes with involutions  $\lambda_X, \lambda_Z$ , then  $\lambda_X \times \lambda_Z$  endows  $X \times Z$  with an involution. It suffices to show that  $X \times Z$  admits a good quotient, as a good quotient is a categorical quotient and is therefore unique up to isomorphism. By [FW20, Remark 4.20], a good quotient exists if and only if every  $C_2$ -orbit is contained in an affine open subscheme. Consider a  $C_2$ -orbit in  $X \times Z$ . Its image under the projection  $\pi_1: X \times Z \rightarrow X$  (resp.  $\pi_2: X \times Z \rightarrow Z$ ) is contained in an affine open subscheme  $U \subseteq X$  (resp.  $V \subseteq Z$ ). Thus the orbit under consideration is contained in  $U \times Z$ , which is affine.  $\square$

**Construction 5.19.** Assume that  $X$  has a good quotient  $Y$  in the sense of [FW20, Remark 4.20]. We write  $p: X \rightarrow Y$  for the quotient map. Let  $j: \text{Spec } A \simeq U \subseteq Y$  be an affine open subscheme of  $Y$ . Because  $p$  is an affine map, the fiber product  $\text{Spec } B := \text{Spec } A \times_Y X$  is an affine open of  $X$  which is invariant under the  $C_2$ -action. In particular  $\text{Spec } B$  inherits a  $C_2$ -action from  $X$  (hence so does its ring of functions  $B$ ). Now  $A \rightarrow B$  acquires the structure of a  $C_2$ -Green functor  $\underline{\mathcal{Q}}(U)$ . Regarding  $\underline{\mathcal{Q}}(U)$  as a  $C_2$ -spectrum, by the isotropy separation sequence, we have an equivalence of  $A$ -modules  $\underline{\mathcal{Q}}(U)^{\varphi_{C_2}} \simeq \text{cofib}(\text{tr}: B_{hC_2} \rightarrow A)$ .

**Lemma 5.20.** Let  $X$  be a scheme with an involution. Assume that  $X$  has a good quotient  $Y$  in the sense of [FW20, Remark 4.20], and write  $p: X \rightarrow Y$  for the quotient map.

- (i) The assignment of Construction 5.19 lifts to a contravariant functor from (the nerve of) the category of affine opens of  $Y$  to the  $\infty$ -category of Poincaré rings/ $C_2$ - $\mathbb{E}_\infty$ -rings/Tambara functors.
- (ii) The presheaf  $\underline{\mathcal{Q}}$  of (i) defines a Zariski sheaf.
- (iii) Write  $p_*\mathcal{O}_X$  for the sheaf of  $\mathbb{E}_\infty$ - $\mathcal{O}_Y$ -algebras (all functors are derived). Then the pushforward  $p_*$  induces an equivalence  $\mathcal{D}(X) \xrightarrow{\sim} \text{Mod}_{p_*\mathcal{O}_X}$ .

*Proof.* Part (i) follows from a similar argument to [Yan23, Theorem 5.1]; functoriality follows from noting that  $\tau_{\geq 0}$  is a functor. Part (ii) follows from Lemma 5.21. To prove part (iii), consider a Zariski cover  $\{j_i: U_i \rightarrow Y\}$  of  $Y$  by affine opens. By Zariski descent,  $\mathcal{D}(X) \simeq \lim_{p^*(j_i)=p \times_Y j_i: U_i \times_Y X \rightarrow X} \text{Mod}_{\mathcal{O}_X(U_i \times_Y X)}$  and  $\text{Mod}_{p_*\mathcal{O}_X} \simeq \lim_{j_i: U_i \rightarrow Y} \text{Mod}_{p_*\mathcal{O}_X(U_i)}$ , hence the result follows.  $\square$

**Lemma 5.21.** Let  $K$  be a simplicial set, and let  $f: K^\triangleleft \rightarrow C_2\mathbb{E}_\infty\text{Alg}(\text{Sp}^{C_2})$  be a diagram. Then  $f$  is a limit diagram if and only if  $f^e: K^\triangleleft \rightarrow \mathbb{E}_\infty\text{Alg}(\text{Sp})$  and  $f^{C_2}: K^\triangleleft \rightarrow \mathbb{E}_\infty\text{Alg}(\text{Sp})$  are both limit diagrams.

*Proof.* The result follows from the observation that limits in  $\mathbb{E}_\infty\text{Alg}(\text{Sp}^{C_2})$  are computed in  $\text{Sp}^{C_2}$ .  $\square$

**Construction 5.22.** Let  $p: X \rightarrow Y$  as before. Consider the composites

$$\begin{aligned} \text{Mod}_{\underline{\mathcal{Q}}}: \text{Op}(Y)^{\text{op}} &\xrightarrow{\underline{\mathcal{Q}}} C_2\mathbb{E}_\infty\text{Alg}(\text{Sp}^{C_2}) \xrightarrow{\text{Mod}_{(-)}} \text{Cat} \\ \text{Mod}_{\underline{\mathcal{Q}}}^\otimes: \text{Op}(Y)^{\text{op}} &\xrightarrow{\underline{\mathcal{Q}}} C_2\mathbb{E}_\infty\text{Alg}(\text{Sp}^{C_2}) \xrightarrow{\text{Mod}_{(-)}^\otimes} C_2 \otimes \text{Cat}, \end{aligned} \tag{5.23}$$

where  $C_2 \otimes \text{Cat}$  denotes the  $\infty$ -category of (small)  $C_2$ -symmetric monoidal  $C_2$ - $\infty$ -categories. In the notation of Construction 5.19, this functor sends the affine open  $\text{Spec } A \subseteq Y$  to the category of modules in  $C_2$ -spectra over the  $C_2$ - $\mathbb{E}_\infty$ -algebra which has underlying  $C_2$ -Mackey functor  $A \rightarrow B$ . Define  $\text{Mod}_{\underline{\mathcal{Q}}}, \text{Mod}_{\underline{\mathcal{Q}}}^\otimes$  to be the limits in  $\text{Cat}, C_2 \otimes \text{Cat}$ , resp. of the functors in (5.23). In particular, if we write  $s: \int \text{Mod}_{\underline{\mathcal{Q}}} \rightarrow \text{Op}(Y)^{\text{op}}$  for the cocartesian fibration obtained by taking the Grothendieck construction on (5.23), an object of  $\text{Mod}_{\underline{\mathcal{Q}}}$  is a cocartesian section of  $s$ . In other words, it is a choice, for each affine open  $\text{Spec } A$  of  $Y$  (same notation as before), of a module over the  $C_2$ - $\mathbb{E}_\infty$ -algebra which has underlying  $C_2$ -Mackey functor  $A \rightarrow B$  which glue compatibly.

L: bleh...cardinals  
L: invent better notation later



Observe that for each  $A \rightarrow B$ , there is a quadratic norm functor  $N^{C_2}: \text{Mod}_B(\text{Sp}) \rightarrow \text{Mod}_{N^{C_2}B}(\text{Sp}^{C_2})$  and a quadratic relative norm functor  $N^{C_2}: \text{Mod}_B(\text{Sp}) \rightarrow \text{Mod}_{A \rightarrow B}(\text{Sp}^{C_2})$ .

**Construction 5.24.** Let  $X$  be a scheme with an involution, and let  $p: X \rightarrow Y$  exhibit  $Y$  as a good quotient of  $X$ . Assume that  $p$  is affine. The norm functors (resp. relative norm functors)  $N_e^{C_2}$  assemble under Construction 5.19 to a ‘global’ norm functor  $N_Y^{C_2}: \pi_{\#}\mathcal{O}_X \text{Mod} \rightarrow N^{C_2}\pi_{\#}\mathcal{O}_X \text{Mod}$  (resp. relative norm functor  $N_Y^{C_2}: \pi_{\#}\mathcal{O}_X \text{Mod} \rightarrow \mathcal{Q} \text{Mod}$ ). Moreover, these functors are quadratic.

For each affine open  $j: \text{Spec } A \subseteq Y$ , write  $B = \Gamma\mathcal{O}_{\text{Spec } A \times_Y X}$ , consider the composite

$$\pi_{\#}\mathcal{O}_X \text{Mod} \xrightarrow{j^*} \text{Mod}_B(\text{Sp}) \xrightarrow{N^{C_2}} \text{Mod}_{N^{C_2}B}(\text{Sp}^{C_2}) \xrightarrow{-\otimes_{N^{C_2}B}(A \rightarrow B)} \text{Mod}_{A \rightarrow B}(\text{Sp}^{C_2}),$$

where the last map is base change along the map  $N^{C_2}B \rightarrow (A \rightarrow B)$  which is a structure map for the  $C_2$ - $\mathbb{E}_{\infty}$ -algebra structure on  $A \rightarrow B$ . Now since quadratic functors are closed under limits [Lur17, Theorem 6.1.1.10] and  $N_Y^{C_2}$  can be written as a limit of a diagram of quadratic functors,  $N_Y^{C_2}$  is also quadratic.

**Definition 5.25.** Varying  $X \rightarrow Y$ , Constructions 5.22 and 5.24 define a functor

$$\begin{aligned} (\text{qSch}^{C_2})^{\text{op}} &\rightarrow C_2 \otimes \text{Cat} \\ (X, \lambda, Y, p) &\mapsto \underline{\text{Mod}}_{\mathcal{Q}}(\underline{\text{Sp}}^{C_2}) \end{aligned}$$

**Definition 5.26.** Suppose  $\mathcal{C}$  is a  $C_2$ -stable  $C_2$ -symmetric monoidal  $C_2$ - $\infty$ -category. Define a functor

$$\begin{aligned} \text{eInv}: C_2 \otimes \text{Cat}^{\text{ex}} &\rightarrow \mathcal{S} \\ (\mathcal{C}, \otimes) &\mapsto (C^{C_2})^{\simeq} \times_{(C^e)^{\simeq, hC_2}} \text{Pic}(C^e)^{\simeq, hC_2}. \end{aligned}$$

In other words,  $\text{eInv}$  sends a  $C_2$ -symmetric monoidal  $C_2$ - $\infty$ -category to the full subgroupoid of  $\mathcal{C}^{C_2}$  on those objects  $L$  so that  $L^e$  is an invertible object in  $C^e$ .

Write  $\widetilde{\text{eInv}}$  for the Grothendieck construction on  $\text{eInv}$ .

There is a functor

$$\begin{aligned} \widetilde{\text{eInv}} &\rightarrow \mathbb{E}_{\infty} \text{Alg}(\text{Cat}^h) \\ (\mathcal{C}, L) &\mapsto (C^e, C^e \xrightarrow{N_{\mathcal{C}}} C^{C_2} \xrightarrow{\text{hom}_{C^{C_2}}(-, L)} \text{Sp}) \end{aligned} \tag{5.27}$$

**Lemma 5.28.** The functor of (5.27) lifts to a functor  $\widetilde{\text{eInv}} \rightarrow \text{Cat}_{\infty}^{\text{p}}$ .

**Lemma 5.29.** Let  $X$  be a scheme with involution  $\sigma: X \xrightarrow{\sim} X$  equipped with a good quotient  $\pi: X \rightarrow Y$ . Let  $L$  be a line bundle on  $Y$ . Then the canonical map

$$L \rightarrow \pi_{\#}\pi^*L \tag{5.30}$$

promotes (5.30) to a sheaf of  $\mathcal{Q}$ -modules on  $Y$ . We will write  $\underline{L}$  for (5.30).

*Proof.* Follows from naturality of the unit and the canonical identification  $\pi^*\mathcal{O}_Y$  with  $\mathcal{O}_X$ .  $\square$

**Definition 5.31.** Let  $X$  be a scheme with involution  $\sigma: X \xrightarrow{\sim} X$  equipped with a good quotient  $\pi: X \rightarrow Y$ . Let  $L$  be a line bundle on  $Y$ . Define  $\mathfrak{Q}_{\sigma, L}$  to be the functor

$$\text{Perf}_X^{\text{op}} \xrightarrow{\pi_{\#}} \pi_{\#}\mathcal{O}_X \text{Mod}^{\omega, \text{op}} \xrightarrow{N^{C_2}} N^{C_2}\pi_{\#}\mathcal{O}_X \text{Mod}(\text{Sp}^{C_2})^{\text{op}} \xrightarrow{\text{hom}_{N^{C_2}\pi_{\#}\mathcal{O}_X}(-, \underline{L})} \text{Sp},$$

where  $\underline{L}$  is a  $\mathcal{Q}$ -module by Lemma 5.29 and  $\mathcal{Q}$  is a  $N^{C_2}\pi_{\#}\mathcal{O}_X$ -algebra by Lemma 5.20. By Construction 5.24 and the fact that the composite of an exact (1-excise) functor and an  $m$ -exciseive functor is  $m$ -exciseive (see [Bar+22, §2.2]),  $\mathfrak{Q}_{\sigma, L}$  is quadratic.

L: todo: use effective descent/limit definition for  $\mathcal{Q}$ -modules.

L: want: codomain consists of  $C_2$ -stable  $C_2$ -presentable  $C_2$ - $\infty$ -categories

L: Pretty sure distributive norm functors are 2-exciseive (results here should generalize readily)...if not, can define the problem away.

L: Example: Special case where  $X$  has trivial  $C_2$  action and  $X = Y$ .

**Example 5.32.** Suppose  $L = \mathcal{O}_Y$ . Then we drop  $L$  from notation and the quadratic functor  $\mathfrak{Q}_\sigma$  of Definition 5.31 takes the form

$$\mathrm{Perf}_X^{\mathrm{op}} \xrightarrow{\pi_\#} \pi_\# \mathcal{O}_X \mathrm{Mod}^{\omega, \mathrm{op}} \xrightarrow{N^{C_2}} N^{C_2} \pi_\# \mathcal{O}_X \mathrm{Mod} \left( \mathrm{Sp}^{C_2} \right)^{\mathrm{op}} \xrightarrow{\mathrm{hom}_{N^{C_2} \pi_\# \mathcal{O}_X}(-, \mathcal{O})} \mathrm{Sp}.$$

**Lemma 5.33.** *Let  $X$  be a scheme with involution  $\sigma: X \xrightarrow{\sim} X$ , and let  $Y$  be a good quotient of  $X$ . Let  $L$  be a line bundle on  $Y$ , and let  $\mathfrak{Q}_{\sigma, L}$  be the quadratic functor on  $\mathrm{Perf}_X$  of Definition 5.31. Then the bilinear part of  $\mathfrak{Q}_{\sigma, L}$  agrees with that of Observation 5.4. In particular,  $(\mathrm{Perf}_X, \mathfrak{Q}_{\sigma, L})$  is a Poincaré  $\infty$ -category.*

*Proof.* By definition of the bilinear part of a quadratic functor, it suffices to show that there is an equivalence  $\mathrm{hom}_{\pi_\# \mathcal{O}_X \mathrm{Mod}}(\pi_\# E \otimes_{\pi_\# \mathcal{O}_X} \pi_\# E, \pi_\# \mathcal{O}_X) \simeq \mathrm{hom}_{\mathcal{O}_X \mathrm{Mod}}(E \otimes_{\mathcal{O}_X} \sigma^* E, \mathcal{O}_X)$  for any perfect complex  $E$  on  $X$ . This follows from Lemma 5.20(iii).  $\square$

**Remark 5.34.** Compare the description of the space of bilinear forms in Lemma 5.33 with the description of a  $\delta$ -hermitian form  $H$  in [PS92, p. 216].

## 6 The Poincaré Brauer Group

Let  $A$  be a Poincaré ring spectrum. By Remark 3.3,  $\mathrm{Mod}_A^\omega$  promotes to a commutative algebra object in the  $\infty$ -category of Poincaré  $\infty$ -categories  $\mathrm{Cat}_\infty^{\mathrm{p}}$ , and we may thus consider modules over it. In this section, we will use modules over Poincaré ring spectra to define derived analogues of the involutive Brauer group for Poincaré ring spectra.

Recall that a Poincaré  $\infty$ -category is called idempotent complete if the underlying stable  $\infty$ -category is idempotent complete. The full subcategory of  $\mathrm{Cat}_\infty^{\mathrm{p}}$  spanned by idempotent complete Poincaré  $\infty$ -categories is denoted by  $\mathrm{Cat}_{\infty, \mathrm{idem}}^{\mathrm{p}}$  [Cal+20b, Definition 1.3.2].

**Definition 6.1.** Let  $A$  be a Poincaré ring spectrum. We define the *Poincaré Brauer space* of  $A$  as

$$\mathrm{Br}^{\mathrm{p}}(A) := \mathrm{Pic}(\mathrm{Mod}_A(\mathrm{Cat}_{\infty, \mathrm{idem}}^{\mathrm{p}})).$$

The assignment  $A \mapsto \mathrm{Br}^{\mathrm{p}}(A)$  defines a functor

$$\mathrm{Br}^{\mathrm{p}}: \mathrm{CAlg}^{\mathrm{p}} \rightarrow \mathrm{CAlg}^{\mathrm{gp}}(\mathcal{S})$$

valued in grouplike  $\mathbf{E}_\infty$ -spaces.

**Remark 6.2.** The symmetric monoidal forgetful functor  $\mathrm{Mod}_A(\mathrm{Cat}_{\infty, \mathrm{idem}}^{\mathrm{p}}) \rightarrow \mathrm{Mod}_A(\mathrm{Cat}_\infty^{\mathrm{ex}})$  induces a map  $\mathrm{Br}^{\mathrm{p}}(A) \rightarrow \mathrm{Br}(A)$  of grouplike  $\mathbf{E}_\infty$ -spaces, where  $\mathrm{Br}(A)$  is the Brauer space  $\mathrm{br}_{\mathrm{alg}}(A)$  of [AG14, pp. 1154–1155].

**Proposition 6.3.** *Let  $A$  be a Poincaré ring spectrum. Then we have a canonical equivalence*

$$\Omega \mathrm{Br}^{\mathrm{p}}(A) \simeq \mathrm{Pic}^{\mathrm{p}}(A).$$

*Proof.* Since  $\Omega \mathrm{Br}^{\mathrm{p}}(R)$  is given by the space of automorphisms of any object in  $\mathrm{Br}^{\mathrm{p}}(R)$ , it suffices to determine the space of autoequivalences of  $(\mathrm{Mod}_R^\omega, \mathfrak{Q}_R)$ . An autoequivalence is the data of a pair  $(f, \eta)$  where  $f: \mathrm{Mod}_R^\omega \xrightarrow{\sim} \mathrm{Mod}_R^\omega$  is an exact  $R$ -linear autoequivalence and  $\eta: \mathfrak{Q}_R \xrightarrow{\sim} \mathfrak{Q}_R \circ f^{\mathrm{op}}$  is a natural equivalence. Since  $\mathrm{Cat}_{\infty R}^{\mathrm{p}} \rightarrow \mathrm{Cat}_{\infty R}^{\mathrm{ex}}$  is symmetric monoidal,  $f$  is of the form  $- \otimes_R \mathcal{L}$  where  $\mathcal{L}$  is an invertible  $R$ -module. Since taking bilinear and linear parts is functorial/by [Cal+20a, Proposition 1.3.11],  $\eta$  is equivalently the data of a pair of equivalences

$$\begin{aligned} b(\eta): \mathrm{hom}_{R \otimes R}((-\otimes \mathcal{L}) \otimes (-\otimes \mathcal{L}), R)^{hC_2} &\simeq \mathrm{hom}_{R \otimes R}(-\otimes -, R)^{hC_2} \\ \ell(\eta): \mathrm{hom}_R(-\otimes \mathcal{L}, R^{\varphi C_2}) &\simeq \mathrm{hom}_R(-, R^{\varphi C_2}) \end{aligned}$$

L: What else do we need to do to show that we have an equivalence of functors?

plus a path between their images in  $\text{hom}_R(\mathcal{L}, R^{tC_2})$ . The transformation  $b(\eta)$  is equivalent to the data of an  $R$ -bilinear equivalence  $R \simeq \mathcal{L}^\vee \otimes \mathcal{L}^\vee$ , and the transformation  $\ell(\eta)$  is equivalent to the data of an  $R^{\varphi C_2}$ -linear equivalence  $\ell(\eta): R^{\varphi C_2} \otimes_R \mathcal{L}^\vee \xrightarrow{\sim} R^{\varphi C_2}$ .

Now consider the composites

$$\begin{aligned} R \otimes_R \mathcal{L}^\vee &\xrightarrow{\text{unit} \otimes \text{id}} R^{\varphi C_2} \otimes \mathcal{L}^\vee \xrightarrow{\ell(\eta)} R^{\varphi C_2} \\ R \otimes_R \mathcal{L} &\xrightarrow{\text{unit} \otimes \text{id}} R^{\varphi C_2} \otimes \mathcal{L} \xrightarrow{\ell(\eta)^{-1} \otimes \text{id}_{\mathcal{L}}} R^{\varphi C_2}. \end{aligned}$$

These correspond to the  $\ell(q^\vee), \ell(q)$  of Remark 4.2, respectively. In particular, the condition that  $\ell(q^\vee), \ell(q)$  make the diagram (4.4) commute is equivalent to the condition that  $\ell(\eta)$  is an equivalence by an adjunction argument.  $\square$

**Proposition 6.4.** *Let  $(\text{Mod}_R^\omega, \mathfrak{Y}_R)$  be a Poincaré ring spectrum.*

- (1) *The  $\infty$ -category  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_{\infty, \text{idem}}^{\text{P}})$  admits all small limits and colimits, and it inherits a canonical symmetric monoidal structure, and for every morphism  $(R, R^{\varphi C_2} \rightarrow R^{tC_2}) \rightarrow (S, S^{\varphi C_2} \rightarrow S^{tC_2})$ , the functor  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_{\infty, \text{idem}}^{\text{P}}) \rightarrow \text{Mod}_{(\text{Mod}_S^\omega, \mathfrak{Y}_S)}(\text{Cat}_{\infty, \text{idem}}^{\text{P}})$  is a symmetric monoidal left adjoint.*
- (2) *Let  $A$  be an  $\mathbb{E}_1$ - $R$ -algebra in spectra, and regard the category of compact right  $A$ -modules  $\text{Mod}_A^\omega$  as left-tensored over  $\text{Mod}_R^\omega$  in the canonical way. Then the pullback*

$$\begin{array}{ccc} & \text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_\infty^{\text{h}}) & \\ & \downarrow & \\ \{\text{Mod}_A^\omega\} & \longrightarrow & \text{Cat}_{\infty}^{\text{ex}} \end{array} \quad (6.5)$$

*is canonically equivalent to  $\text{Mod}_{N_R A \otimes_{N_R R} R^L}(\text{Sp}^{C_2})$  where  $R^L$  is the  $\mathbb{E}_\infty$ - $N_R R$ -algebra with  $(R^L)^e \simeq R$  and  $(R^L)^{\varphi C_2} \simeq C$ .*

*A  $N_R A \otimes_{N_R R} R^L$ -module classifies a  $(\text{Mod}_R^\omega, \mathfrak{Y}_R)$ -module in Poincaré  $\infty$ -categories if its underlying  $R$ -module is invertible in the sense of [Cal+20a, Definition 3.1.4].*

- (3) *Let  $A, B$  be  $R$ -algebras with associated  $(R$ -linear) modules with genuine involution  $(M_A, N_A, N_A \rightarrow M_A^{tC_2})$  and  $(M_B, N_B, N_B \rightarrow M_B^{tC_2})$ , respectively so that (under item (2))  $(\text{Mod}_A^\omega, \mathfrak{Y}_A)$  and  $(\text{Mod}_B^\omega, \mathfrak{Y}_B)$  are objects of  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_{\infty, \text{idem}}^{\text{P}})$ . Then the symmetric monoidal structure of item (1) is so that the underlying  $R$ -linear  $\infty$ -category with perfect duality  $(\text{Mod}_A^\omega, \mathfrak{Y}_A) \otimes_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)} (\text{Mod}_B^\omega, \mathfrak{Y}_B)$  is  $\text{Mod}_A^\omega \otimes_{\text{Mod}_R^\omega} \text{Mod}_B^\omega \simeq \text{Mod}_{A \otimes_R B}^\omega$ , and the associated module with genuine involution is given by  $M_A \otimes_R M_B, N_A \otimes_{R^{\varphi C_2}} N_B$ , and the structure map is  $N_A \otimes_{R^{\varphi C_2}} N_B \rightarrow M_A^{tC_2} \otimes_{R^{tC_2}} M_B^{tC_2} \rightarrow (M_A \otimes_R M_B)^{tC_2}$  where the latter map arises canonically from lax monoidality of the Tate construction.*
- (4) *Let  $(\mathcal{C}, \mathfrak{Y}_{\mathcal{C}}), (\mathcal{D}, \mathfrak{Y}_{\mathcal{D}})$  be objects of  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_\infty^{\text{h}})$ . Then the forgetful functor induces  $\text{hom}_{\text{Cat}_{\infty}^{\text{h}}}((\mathcal{C}, \mathfrak{Y}_{\mathcal{C}}), (\mathcal{D}, \mathfrak{Y}_{\mathcal{D}})) \rightarrow \text{hom}_{\text{Cat}_{\infty}^{\text{ex}}}(\mathcal{C}, \mathcal{D})$  on mapping spaces so that the fiber over an  $R$ -linear functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is the mapping space  $\text{map}_{\mathfrak{Y}_R}(F_! \mathfrak{Y}_{\mathcal{C}}, \mathfrak{Y}_{\mathcal{D}}) \simeq \text{map}_{\mathfrak{Y}_R}(\mathfrak{Y}_{\mathcal{C}}, \mathfrak{Y}_{\mathcal{D}} \circ F^{\text{op}})$ , where the mapping space is taken in  $\text{Fun}_{\mathfrak{Y}_R}^q(\mathcal{D}^{\text{op}}, \text{Sp})$  and  $\text{Fun}_{\mathfrak{Y}_R}^q(\mathcal{C}^{\text{op}}, \text{Sp})$ , respectively.<sup>3</sup>*
- (5) *The symmetric monoidal forgetful functor  $\theta: \text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Y}_R)}(\text{Cat}_\infty^{\text{h}}) \rightarrow \text{Mod}_{\text{Mod}_R^\omega}(\text{Cat}_\infty^{\text{ex}})$  is a (co)cartesian fibration.*

**Remark 6.6.** A special case of part (2) is [Cal+20a, Example 5.4.13].

*Proof.* (1) The first part of the statement follows from [Cal+20a, §6.1] and [Lur17, §4.2.3].

<sup>3</sup>The proof of (2) in particular shows that  $\text{Fun}^q(\mathcal{C}^{\text{op}}, \text{Sp})$  is left-tensored over  $\text{Fun}^q(\text{Mod}_R^{\omega, \text{op}}, \text{Sp})$  in the sense of [Lur17, Definition 4.2.1.19], so this makes sense.

L: maybe one of these should be conjugate dual here?

L: is the  $R^{\varphi C_2}$ -linearity of this  $\simeq$  correct?

L: under construction— not sure what to say about the  $(-)^{tC_2}$  part yet.

L: what is it classified by?

- (2) Let  $\mathcal{LM}^\otimes$  denote the  $\infty$ -operad of [Lur17, Definition 4.2.1.7]. Our strategy of proof will be similar to that of [Cal+20a, §5.3]: First, we show that an  $\mathcal{LM}^\otimes$ -algebra object in  $\text{Cat}_\infty^h$  is equivalent to an  $\mathcal{LM}^\otimes$ -algebra object in an operad of functor categories. Then, we use a (suitably coherent version of) the classification of hermitian structures on module categories as categories of modules over the Hill–Hopkins–Ravenel norm [Cal+20a, Theorem 3.3.1] to conclude. Recall that the action of  $\text{Mod}_R^\omega$  on  $\text{Mod}_A^\omega$  is given by a functor  $\mathcal{LM}^\otimes \rightarrow \text{Cat}_\infty^\times$ , and define  $\text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes$  via the following pullback square of  $\infty$ -operads:

$$\begin{array}{ccc} \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \xrightarrow{p} & \mathcal{LM}^\otimes \\ \downarrow & & \downarrow \text{Mod}_R^\omega, \text{Mod}_A^\omega \\ (\text{Cat}_\infty)_{\text{op}/-/\text{Sp}}^\otimes & \longrightarrow & \text{Cat}_\infty^\times \end{array} \quad (6.7)$$

Informally, an object  $F \in \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes$  is a functor  $F: \text{Mod}_R^{\omega, \text{op}} \rightarrow \text{Sp}$  and an object  $G$  over the fiber of  $\mathfrak{m}$  is a functor  $G: \text{Mod}_A^{\omega, \text{op}} \rightarrow \text{Sp}$ . The  $p$ -cocartesian edge over the canonical map  $(\mathfrak{a}, \mathfrak{m}) \rightarrow \mathfrak{m}$  in  $\mathcal{LM}^\otimes$  sends  $(F, G)$  to the lower arrow in the diagram

$$\begin{array}{ccc} \text{Mod}_R^{\omega, \text{op}} \times \text{Mod}_A^{\omega, \text{op}} & \xrightarrow{F \times G} & \text{Sp} \times \text{Sp} \\ \downarrow - \otimes_R - & & \downarrow \otimes_{\text{Sp}} \\ \text{Mod}_A^{\omega, \text{op}} & \xrightarrow{F \otimes G := \text{LKE}_{\otimes_R}(\otimes_{\text{Sp}} \circ (F \times G))} & \text{Sp} \end{array}$$

Now define  $\text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^q(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes$  to consist of the full subcategory of  $\text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes$  consisting of those tuples of functors which are all quadratic. The inclusion  $\text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^q(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes \rightarrow \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes$  exhibits the former as an  $\infty$ -operad, and moreover the localization is compatible with the  $\mathcal{LM}^\otimes$ -monoidal structure in the sense of [Lur17, Definition 2.2.1.6]. We can extend the previous diagram to

$$\begin{array}{ccccc} \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^q(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \longrightarrow & \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \xrightarrow{p} & \mathcal{LM}^\otimes \\ \downarrow & & \downarrow & & \downarrow \text{Mod}_R^\omega, \text{Mod}_A^\omega \\ \text{Cat}_\infty^h \otimes & \longrightarrow & (\text{Cat}_\infty)_{\text{op}/-/\text{Sp}}^\otimes & \longrightarrow & \text{Cat}_\infty^\otimes \end{array} \quad (6.8)$$

Modifying [Cal+20a, Construction 5.3.15 & Lemma 5.3.15] slightly (note that Corollary 5.1.4 did not assume the tensor factors to be equivalent), we obtain an analogous commutative diagram of  $\infty$ -operads

$$\begin{array}{ccccccc} \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^p(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \longrightarrow & \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^q(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \longrightarrow & \text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes & \xrightarrow{p} & \mathcal{LM}^\otimes \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \text{Mod}_R^\omega, \text{Mod}_A^\omega \\ \text{Cat}_\infty^p \otimes & \longrightarrow & \text{Cat}_\infty^h \otimes & \longrightarrow & (\text{Cat}_\infty)_{\text{op}/-/\text{Sp}}^\otimes & \longrightarrow & \text{Cat}_\infty^\otimes \end{array} \quad (6.9)$$

in which all squares are pullbacks. Now suppose  $A$  is given a module with genuine involution  $(M_A, N_A, N_A \rightarrow M_A^{tC_2})$  and call the associated Poincaré  $\infty$ -category  $\overline{\text{Mod}}_A$ . Then to lift  $\overline{\text{Mod}}_A$  to a module over  $(\text{Mod}_R^\omega, \mathfrak{I}_R)$  compatibly with the  $\text{Mod}_R^\omega$ -module structure on  $\text{Mod}_A^\omega$  is to give a map of  $\infty$ -operads  $\mathcal{LM}^\otimes \rightarrow \text{Cat}_\infty^h \otimes$  so that the restriction along the canonical inclusion  $\text{Assoc}^\otimes \rightarrow \mathcal{LM}^\otimes$  gives the algebra object  $(\text{Mod}_R^\omega, \mathfrak{I}_R)$  and postcomposing with the canonical projection to  $\text{Cat}_\infty^{\text{ex}}$  recovers the given  $\text{Mod}_R^\omega$ -module structure on  $\text{Mod}_A^\omega$ . By the pullback square (6.8), this is equivalent to giving an object of  $\text{Alg}_{\mathcal{LM}/\mathcal{LM}}(\text{Fun}_{\text{Mod}_R^\omega, \text{Mod}_A^\omega}^q(\text{Mod}_A^{\omega, \text{op}}, \text{Sp})^\otimes)$ . Now let us identify the bilinear functor  $\text{Mod}_R^\omega \times \text{Mod}_A^\omega \xrightarrow{- \otimes_R -} \text{Mod}_A^\omega$  with the exact functor  $\text{Mod}_R^\omega \otimes \text{Mod}_A^\omega \simeq \text{Mod}_{R \otimes A}^\omega \rightarrow \text{Mod}_A^\omega$  which is induction along the action map  $R \otimes A \rightarrow A$ . Using [Cal+20a, Corollary 3.4.1] and unravelling definitions gives the claim for  $R$ -linear hermitian structures. The proof for  $R$ -linear Poincaré structures considers (6.9) instead but otherwise proceeds in an identical fashion.

- (3) By [Lur17, Theorem 4.4.2.8], the relative tensor product  $(\mathrm{Mod}_A^\omega, \mathfrak{P}_A) \otimes_{(\mathrm{Mod}_R^\omega, \mathfrak{P}_R)} (\mathrm{Mod}_B^\omega, \mathfrak{P}_B)$  is computed as the geometric realization of the bar construction

$$p: \Delta^{\mathrm{op}} \rightarrow \mathrm{Cat}_\infty^{\mathrm{h}} \\ [n] \mapsto (\mathrm{Mod}_A^\omega, \mathfrak{P}_A) \otimes (\mathrm{Mod}_R^\omega, \mathfrak{P}_R)^{\otimes n} \otimes (\mathrm{Mod}_B^\omega, \mathfrak{P}_B)$$

Write  $f: \mathrm{Cat}_\infty^{\mathrm{h}} \rightarrow \mathrm{Cat}_\infty^{\mathrm{ex}}$  for the forgetful functor. Then  $f \circ p$  has a colimit with value  $\mathrm{Mod}_A^\omega \otimes_{\mathrm{Mod}_R^\omega} \mathrm{Mod}_B^\omega \simeq \mathrm{Mod}_{A \otimes_R B}^\omega$ . Writing  $g: \mathrm{Cat}_\infty^{\mathrm{ex}} \rightarrow \{*\}$ , by Example 4.3.1.3 of [Lur09] we see that  $f \circ p$  is a  $g$ -colimit. By Proposition 4.3.1.5(2) and Example 4.3.1.3 of [Lur09],  $p$  admits a colimit in  $\mathrm{Cat}_\infty^{\mathrm{h}}$  if and only if it admits an  $f$ -colimit. Now recall that  $f$  is a cocartesian fibration with pushforward given by left Kan extension [Cal+20a, Corollary 1.4.2]. We show that  $f$  satisfies the conditions of [Lur09, Corollary 4.3.1.11].

- Condition (1) follows from Theorem 6.1.1.10 of [Lur17] applied to  $\mathrm{Sp}^{\mathrm{op}}$  (see the end of [Cal+20a, Construction 1.1.26]).
- Condition (2) follows from [Cal+20a, Corollary 1.4.2], the adjoint functor theorem, and presentability of  $\mathrm{Fun}^q(\mathcal{C})$ , which is discussed in the proof of [Cal+20a, Lemma 5.3.3] (also see [Lur17, Remark 6.1.1.11]).

Thus the preceding discussion shows that there exists a map of simplicial sets  $p'$  making the diagram commute

$$\begin{array}{ccc} \Delta^{\mathrm{op}} & \xrightarrow{p} & \mathrm{Cat}_\infty^{\mathrm{h}} \\ \downarrow & \nearrow p' & \downarrow f \\ (\Delta^{\mathrm{op}})^\triangleright & \longrightarrow & \mathrm{Cat}_\infty^{\mathrm{ex}} \end{array} .$$

Since  $\{0\} \rightarrow \Delta^1$  is left anodyne, by [Lur09, Corollary 2.1.2.7] the inclusions

$$\begin{aligned} \{0\} \times \Delta^{\mathrm{op}} &\rightarrow \Delta^1 \times \Delta^{\mathrm{op}} \\ \iota: (\{0\} \times (\Delta^{\mathrm{op}})^\triangleright) \sqcup_{\{0\} \times \Delta^{\mathrm{op}}} (\Delta^1 \times \Delta^{\mathrm{op}}) &\rightarrow \Delta^1 \times (\Delta^{\mathrm{op}})^\triangleright \end{aligned}$$

are left anodyne. The former implies that there exists a map  $p''$  making the diagram

$$\begin{array}{ccc} \{0\} \times \Delta^{\mathrm{op}} & \xrightarrow{p} & \mathrm{Cat}_\infty^{\mathrm{h}} \\ \downarrow & \nearrow p'' & \downarrow f \\ \Delta^1 \times \Delta^{\mathrm{op}} & \longrightarrow & \mathrm{Cat}_\infty^{\mathrm{ex}} \end{array}$$

commute. The maps  $p'$  and  $p''$  assemble to give a map  $p''' := p' \sqcup_p p''$  making the diagram

$$\begin{array}{ccc} \{0\} \times \Delta^{\mathrm{op}} & \xrightarrow{p} & \mathrm{Cat}_\infty^{\mathrm{h}} \\ \downarrow & \nearrow p''' & \downarrow f \\ (\{0\} \times (\Delta^{\mathrm{op}})^\triangleright) \sqcup_{\{0\} \times \Delta^{\mathrm{op}}} (\Delta^1 \times \Delta^{\mathrm{op}}) & \xrightarrow{\iota} \Delta^1 \times (\Delta^{\mathrm{op}})^\triangleright & \xrightarrow{\bar{p}} \mathrm{Cat}_\infty^{\mathrm{ex}} \end{array}$$

commute, and likewise  $\bar{p}$  exists making the diagram commute since  $\iota$  is left anodyne. Now we show that  $\bar{p}$  satisfies the conditions of [Lur09, Proposition 4.3.1.9]. By (the opposite/dual/cocartesian version of) [Lur09, Remark 3.1.1.10] and Proposition 3.1.1.5(2'') *ibid.* and the fact that  $f$  is a cocartesian fibration, we can choose  $\bar{p}$  so that for all  $k \in (\Delta^{\mathrm{op}})^\triangleright$ ,  $\bar{p}|_{\Delta^1 \times \{k\}}$  is  $f$ -cocartesian. Furthermore, since we can choose  $\Delta^{\mathrm{op}}$ ,  $(\Delta^{\mathrm{op}})^\triangleright$  to have the markings  $(-)^\triangleright$  in [Lur09, Remark 3.1.1.10],  $f \circ \bar{p}|_{\Delta^1 \times \{*\}}$  is a degenerate edge in  $\mathrm{Cat}_\infty^{\mathrm{ex}}$ .

Now [Lur09, Proposition 4.3.1.9] implies that  $\bar{p}_0$  is an  $f$ -colimit diagram if and only if  $\bar{p}_1$  is an  $f$ -colimit diagram. Now notice that  $\bar{p}|_{\{1\} \times (\Delta^{\mathrm{op}})^\triangleright}$  has image contained in the fiber of  $f$  over  $\mathrm{Mod}_{A \otimes_R B}^\omega$ .

By [Lur09, Proposition 4.3.1.10], it suffices to show that  $\bar{p}_1$  is a colimit diagram in  $\text{Fun}^q(\text{Mod}_{A \otimes_R B}^\omega)$ . Write  $\bar{M}_A \in \text{Mod}_{N^{C_2}A}$  and  $\bar{M}_B \in \text{Mod}_{N^{C_2}B}$  for the corresponding modules (see introduction to §3.3 of [Cal+20a]). Unraveling definitions and using [Cal+20a, Theorem 3.3.1 & Corollary 3.4.1 & Lemma 5.4.6], it follows that the diagram  $\bar{p}_1|_{\{1\} \times \Delta^{\text{op}}}$  is the bar construction

$$[n] \mapsto \bar{M}_A \otimes_{N^{C_2}R} R^{\otimes_{N^{C_2}R} n} \otimes_{N^{C_2}R} \bar{M}_B.$$

This proves the result.

- (4) Let  $(\mathcal{C}, \mathfrak{Q}_{\mathcal{C}})$  be an object of  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Q}_R)}(\text{Cat}_\infty^h)$  and let  $F: \mathcal{C} = \theta(\mathcal{C}, \mathfrak{Q}_{\mathcal{C}}) \rightarrow \mathcal{D}$  be an  $R$ -linear functor. Now define  $\mathfrak{Q}_{\mathcal{D}}: \mathcal{D}^{\text{op}} \rightarrow \text{Sp}$  to be the left Kan extension of  $\mathfrak{Q}_{\mathcal{C}}$  along  $F^{\text{op}}$ . Now  $(\mathcal{D}, \mathfrak{Q}_{\mathcal{D}}) \in \text{Cat}_\infty^h$  and there is a canonical map  $(f, \eta): (\mathcal{C}, \mathfrak{Q}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathfrak{Q}_{\mathcal{D}})$ . Now  $F$  is classified by a functor  $\Delta^1 \times \mathcal{LM}^\otimes \rightarrow \text{Cat}_\infty^{\text{ex} \otimes}$ , and we may form the pullback

$$\begin{array}{ccc} \mathcal{N} & \longrightarrow & \Delta^1 \times \mathcal{LM}^\otimes \\ \downarrow & & \downarrow \\ \text{Cat}_\infty^h & \xrightarrow{p} & \text{Cat}_\infty^{\text{ex} \otimes} \end{array} \quad (6.10)$$

Since  $p$  is a cocartesian fibration [Cal+20a, Theorem 5.2.7],  $\mathcal{N} \rightarrow \Delta^1 \times \mathcal{LM}^\otimes$  is a cocartesian fibration, and the nontrivial morphism in  $\Delta^1$  classifies a map  $F_!: \text{Fun}_{\text{Mod}_R^\omega}^q(\mathcal{C}^{\text{op}}, \text{Sp})^\otimes \rightarrow \text{Fun}_{\text{Mod}_R^\omega}^q(\mathcal{D}^{\text{op}}, \text{Sp})^\otimes$  of  $\infty$ -operads over  $\mathcal{LM}^\otimes$ . Passing to algebra objects, we obtain the desired result on mapping spaces.

- (5) By [Lur09, Proposition 2.4.2.8], it suffices to show that  $\theta$  is a locally (co)cartesian fibration, and that locally (co)cartesian edges are closed under composition. We give the proof that  $\theta$  is a cocartesian fibration; the proof that  $\theta$  is a cartesian fibration is formally dual and will be left to the reader.

Let  $(\mathcal{C}, \mathfrak{Q}_{\mathcal{C}})$  be an object of  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Q}_R)}(\text{Cat}_\infty^h)$  and let  $F: \mathcal{C} = \theta(\mathcal{C}, \mathfrak{Q}_{\mathcal{C}}) \rightarrow \mathcal{D}$  be an  $R$ -linear functor. Now define  $\mathfrak{Q}_{\mathcal{D}}: \mathcal{D}^{\text{op}} \rightarrow \text{Sp}$  to be the left Kan extension of  $\mathfrak{Q}_{\mathcal{C}}$  along  $F^{\text{op}}$ . By the proof of (4), we see that the image of  $\mathfrak{Q}_{\mathcal{C}}$  under  $F_!$  is a lift of  $(\mathcal{D}, \mathfrak{Q}_{\mathcal{D}})$  to an object of  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Q}_R)}(\text{Cat}_\infty^h)$  and  $(f, \eta)$  to a morphism in  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Q}_R)}(\text{Cat}_\infty^h)$ .

Now by Lemma 2.4.4.1 and the locally cocartesian version of Proposition 2.4.1.10 of [Lur09], we must show that for all choices  $\mathfrak{Q}'_{\mathcal{D}}$  of an  $R$ -linear Hermitian structure on  $\mathcal{D}$ , precomposition with  $F_!$  induces a pullback square

$$\begin{array}{ccc} \text{hom}_{\text{Cat}_{\infty R}^h}((\mathcal{D}, \mathfrak{Q}_{\mathcal{D}}), (\mathcal{D}, \mathfrak{Q}'_{\mathcal{D}})) & \longrightarrow & \text{hom}_{\text{Cat}_{\infty R}^h}((\mathcal{C}, \mathfrak{Q}_{\mathcal{C}}), (\mathcal{D}, \mathfrak{Q}'_{\mathcal{D}})) \\ \downarrow & & \downarrow \\ \text{hom}_{\text{Cat}_{\infty R}^{\text{ex}}}(\mathcal{D}, \mathcal{D}) & \longrightarrow & \text{hom}_{\text{Cat}_{\infty R}^{\text{ex}}}(\mathcal{C}, \mathcal{D}) \end{array} \quad (6.11)$$

By (4),  $F_!$  induces equivalences on the fibers of the vertical maps, hence  $(f, \eta)$  is locally  $\theta$ -cocartesian. The locally  $\theta$ -cocartesian maps are manifestly closed under composition, hence we are done.  $\square$

**Corollary 6.12.** *Let  $R$  be a Poincaré ring, and let  $A, B$  be  $\mathbb{E}_1$ - $R$ -algebras with genuine involution. Then there is an equivalence  $\text{hom}_{\text{Cat}_{\infty, \text{idem}_R}^p}((\text{Mod}_A^\omega, \mathfrak{Q}_A), (\text{Mod}_B^\omega, \mathfrak{Q}_B)) \simeq (\text{BiMod}_{A \otimes_R B^{\text{op}}})_{A \varphi^{C_2} \otimes_R B \varphi^{C_2}} / -$ .*

*Proof.*

$\square$

As in the Picard group case, the symmetric monoidal forgetful functor  $\theta: \text{Cat}_{\infty R}^p \rightarrow \text{Cat}_{\infty R}^{\text{ex}}$  induces a map of spectra  $\theta: \text{Br}^p(A) \rightarrow \text{Br}(A^e)$ . When  $A^e$  is endowed with the trivial action,  $\theta$  will factor through the 2-torsion on  $\pi_0$ . As a consequence of Proposition 6.4(2) we can identify the fiber of this map.

**Corollary 6.13.** *Let  $(\text{Mod}_A^\omega, \mathfrak{Q}_A)$  be a Poincaré ring with underlying genuine  $C_2$  spectrum  $A^L$  as in Proposition 6.4(2). Then the fiber of the map*

$$\text{Br}^p(A) \rightarrow \text{Br}(A)$$

*can be naturally identified with  $\text{Pic}(\text{Mod}_{A^L}(\text{Sp}^{C_2}))$ .*

L: todo—  
probably  
need to fix the  
statement with  
duals when  
the proof is  
written

*Proof.* Since  $\theta : \text{Mod}_{(\text{Mod}_A^\omega, \mathcal{Q}_A)}(\text{Cat}_{\infty, \text{idem}}^{\text{p}}) \rightarrow \text{Mod}_{\text{Mod}_{A^e}}(\text{Cat}_{\infty}^{\text{ex}})$  is symmetric monoidal and conservative, it induces a map  $\theta^\simeq : \text{PnBr}(A) \rightarrow \text{br}(A^e)$  on the groupoid core of invertible objects. Now observe that  $\theta$  is an isofibration; it follows that  $\theta^\simeq$  is a Kan fibration by [Lur24, Proposition 01EZ]. Consequently we need only identify the fiber instead of the homotopy fiber, which follows from the identification of the fiber in Proposition 6.4(2).  $\square$

Write  $\text{Pn}$  for the composite  $\text{Cat}_{\infty, R}^{\text{p}} \xrightarrow{U} \text{Cat}_{\infty}^{\text{p}} \xrightarrow{\text{Pn}} \mathcal{S}$ .

**Proposition 6.14.** *Let  $(R, R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring. Then  $(\text{Mod}_R^\omega, \mathcal{Q}_R)$  corepresents the functor  $\text{Pn} : \text{Cat}_{\infty, R}^{\text{p}} \rightarrow \mathcal{S}$ .*

*Proof.* Recall that Proposition 6.4(1) furnishes an adjoint pair  $\text{Cat}_{\infty, R}^{\text{p}} \rightleftarrows \text{Cat}_{\infty}^{\text{p}}$  of functors. Write  $\bar{\mathcal{C}} = (\mathcal{C}, \mathcal{Q}_{\mathcal{C}}) \in \text{Cat}_{\infty, \text{idem}_R}^{\text{p}}$ . Then

$$\text{Pn}(\mathcal{C}) = \text{hom}_{\text{Cat}_{\infty}^{\text{p}}} \left( (\text{Sp}^f, \Omega^u), U(\bar{\mathcal{C}}) \right) \simeq \text{hom}_{\text{Cat}_{\infty, R}^{\text{p}}} \left( (\text{Mod}_R^\omega, \mathcal{Q}_R) \otimes (\text{Sp}^f, \Omega^u), \bar{\mathcal{C}} \right),$$

where the first equivalence is [Cal+20a, Proposition 4.1.3].  $\square$

## 6.1 Azumaya algebras with genuine involution

Let  $R$  be an  $\mathbb{E}_\infty$ -ring spectrum.

**Recollection 6.15.** Recall [BRS12; AG14] that an  $\mathbb{E}_1$ - $R$ -algebra  $A$  is said to be *Azumaya* if it is a compact generator of  $\text{Mod}_R$  and if the natural  $R$ -algebra map giving the bimodule structure on  $A$

$$A \otimes_R A^{\text{op}} \rightarrow \text{End}_R(A)$$

is an equivalence of  $R$ -algebras.

**Definition 6.16.** Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring spectrum. An *Azumaya algebra with genuine type 1 (anti-)involution* over  $R$  is the data of

- (a) An  $\mathbb{E}_1$ - $R$ -algebra  $A$  equipped with an anti-involution  $\tau : A \rightarrow A^{\text{op}}$  so that  $A$  is an Azumaya  $R$ -algebra in the sense of Recollection 6.15
- (b) an  $(A \otimes_R A^{\text{op}})^{\otimes R^2}$ -linear equivalence  $\text{hom}_{R \otimes R}(A \otimes_R A, R) \simeq A \otimes_R A^{\text{op}}$
- (c) A left  $A \otimes_R R^{\varphi C_2}$ -module  $P$  and an  $A^{\text{op}} \otimes_R R^{\varphi C_2}$ -module  $\bar{P}$
- (d) An  $A$ -linear map  $P \rightarrow A^{tC_2}$  and an  $A^{\text{op}}$ -linear map  $\bar{P} \rightarrow A^{tC_2}$ . Here we regard  $A^{tC_2}$ , which is canonically a  $(A \otimes A)^{tC_2}$ -module, as an  $A$ -module (resp.  $A^{\text{op}}$ -module) via the twisted Tate-valued diagonal  $A \rightarrow (A \otimes A^{\text{op}})^{tC_2}$  (resp.  $A^{\text{op}} \rightarrow (A \otimes A^{\text{op}})^{tC_2}$ ).
- (e) An equivalence of  $A \otimes_R A^{\text{op}}$ -modules

$$\text{hom}_R(A, R^{\varphi C_2}) \simeq P \otimes_{R^{\varphi C_2}} \bar{P}$$

and a homotopy making the diagram

$$\begin{array}{ccc} \text{hom}_R(A, R^{\varphi C_2}) & \xrightarrow{\hspace{10em}} & P \otimes_{R^{\varphi C_2}} \bar{P} \\ \downarrow & & \downarrow \\ \text{hom}_R(A, R^{tC_2}) \simeq \text{hom}_R(A \otimes A^{\text{op}}, R)^{tC_2} & \longrightarrow & \text{hom}_{(A \otimes A^{\text{op}})^{\otimes 2}}((A \otimes A^{\text{op}})^{\otimes 2}, A \otimes A^{\text{op}})^{tC_2} \simeq (A \otimes A^{\text{op}})^{tC_2} \end{array}$$

commute, where the lower horizontal arrow is induced by (b) and the right vertical arrow is induced by (d).

Similarly, an *Azumaya algebra with genuine type 2 (anti-)involution* over  $R$  is obtained by replacing  $\tau$  in item (a) with  $\tau : A \rightarrow \sigma_R^* A^{\text{op}}$ , where  $\sigma_R : R \xrightarrow{\sim} R$  is the given involution on  $R$ .  $\square$

N: Needs a lot more detail I know, but I think the skeleton is here.

L: define the category!



**Remark 6.17.** If  $A$  is an Azumaya algebra with genuine involution over  $R$ , then in particular  $M_A = A$ ,  $N_A = P$  is a module with genuine involution over  $A$  in the sense of [Cal+20a, Definition 3.2.3].

With ordinary Azumaya algebras, the prototypical Azumaya algebra with anti-involution arises from endomorphism rings of perfect modules. Choosing a (nondegenerate symmetric bilinear) form on a perfect module  $P$  endows its endomorphism algebra with additional structure.

**Example 6.18.** Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring, and let  $(P, q) \in \text{Pn}(\text{Mod}_R^\omega, \mathfrak{Q}_R)$ .

Then  $A := \text{End}_R(P)$  admits a canonical lift to an  $\mathbb{E}_1$  with genuine involution over  $R$  with  $A^{\varphi C_2} := \text{hom}_R(P, R^{\varphi C_2})$ . If  $P$  is a generator of  $\text{Mod}_R^\omega$ , then  $A$  is furthermore Azumaya.

By [Cal+20a, Proposition 3.1.16],  $A$  inherits a canonical anti-involution. To exhibit (b), observe that  $q^\dagger$  induces a canonical  $A \otimes A^{\text{op}}$ -linear equivalence  $A = \text{End}(P) \simeq \text{End}(P^\vee) \simeq A^\vee$ . If  $P$  is a generator,  $\text{hom}_R(P, -)$  induces an equivalence  $\text{Mod}_R^\omega \simeq \text{Mod}_A^\omega$ , thus we can regard  $\text{Mod}_A^\omega$  as equipped with a Poincaré structure. By the classification of  $R$ -linear Poincaré structures of Proposition 6.4(2), the Poincaré structure on  $\text{Mod}_A^\omega$  is associated to an  $A$ -module with genuine involution  $(M_A, N_A, N_A \rightarrow M_A^{tC_2})$ . We claim that  $M_A \simeq A$  with the canonical  $A$ - $A$ -bimodule structure: By [Cal+20a, Proposition 3.1.6], as an  $A^{\text{op}}$ -module  $M_A$  is the image of  $A$  under the composite

$$\text{Mod}_A^\omega \xrightarrow{\text{hom}_R(P, -)^{-1}} \text{Mod}_R^\omega \xrightarrow{D_R = \text{hom}_R(-, R)} \text{Mod}_R^{\omega, \text{op}} \xrightarrow{\text{hom}_R(P, -)} \text{Mod}_A^{\omega, \text{op}}.$$

Observe that the image of  $A$  in  $\text{Mod}_R^{\omega, \text{op}}$  is  $D_R(P)$  and  $q^\dagger$  induces an equivalence  $D_R(P) \simeq P$ , hence  $M_A \simeq A$  as  $A^{\text{op}}$ -modules.

A similar argument with the linear part of  $\mathfrak{Q}$  shows that we have an equivalence  $N_A \simeq \text{hom}_R(P, R^{\varphi C_2})$ .

**Proposition 6.19.** Let  $R$  be a discrete ring with a given  $C_2$ -action  $\lambda$ , and suppose that the ramification locus in  $\text{Spec}(R)/C_2 = \text{Spec}(R^{C_2})$  is empty. Let  $A$  be a classical Azumaya algebra over  $R$  with an involution of type 2. Regard  $R$  as a Poincaré ring spectrum  $\underline{R}^\lambda$  via Example 3.15.

Then there is a canonical Azumaya algebra with genuine involution over  $\underline{R}^\lambda$  whose underlying Azumaya algebra is  $A$ .

**Remark 6.20.** If  $\frac{1}{2} \in R$ , then  $\text{Br}(\text{Spec } R, \lambda)$  is defined [PS92, p. 216]. In view of Propositions 6.19 and 6.21, there is a homomorphism  $\text{Br}(\text{Spec } R, \lambda) \rightarrow \pi_0 \text{PnBr}(\underline{R}^\lambda)$ .

*Proof of Proposition 6.19.* Since  $R^{\varphi C_2} = 0$ , conditions (c)-(e) of Definition 6.16 are vacuous.  $\square$

**Proposition 6.21.** Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring, and let  $(A, A^{\varphi C_2} \rightarrow A^{tC_2})$  be an Azumaya algebra with genuine involution over  $R$ . Then

1.  $(\text{Mod}_A^\omega, \mathfrak{Q}_A)$  defines an  $R$ -linear Poincaré  $\infty$ -category.
2.  $(\text{Mod}_A^\omega, \mathfrak{Q}_A)$  is an invertible object in  $\text{Mod}_{(\text{Mod}_R^\omega, \mathfrak{Q}_R)}(\text{Cat}_{\infty, \text{idem}}^{\text{P}})$ .

*Proof of Proposition 6.21.* The first statement follows from unravelling definitions; we prove the second statement. First, by [Cal+20a, Example 3.2.9], we see that  $(\text{Mod}_A^\omega, \mathfrak{Q}_A)$  is indeed an  $R$ -linear Poincaré  $\infty$ -category (and not merely hermitian). To show that the associated Poincaré  $\infty$ -category is invertible, we must identify a dual  $(\text{Mod}_A^\omega, \mathfrak{Q}_A)^\vee$  and exhibit an equivalence  $(\text{Mod}_A^\omega, \mathfrak{Q}_A) \otimes (\text{Mod}_A^\omega, \mathfrak{Q}_A)^\vee \simeq (\text{Mod}_R^\omega, \mathfrak{Q}_R)$ . Since  $\text{Cat}_{\infty, \text{idem}}^{\text{P}} \rightarrow \text{Cat}_{\infty}^{\text{ex}}$  is symmetric monoidal, we see that the underlying  $R$ -linear  $\infty$ -category associated to the dual must be  $\text{Mod}_{A^{\text{op}}}^\omega$ . Moreover, the canonical evaluation map  $\text{ev}: \text{Mod}_A^\omega \otimes \text{Mod}_{A^{\text{op}}}^\omega \xrightarrow{\sim} \text{Mod}_R^\omega$  sends  $A \otimes A^{\text{op}}$  to  $A$ . Endow  $\text{Mod}_{A^{\text{op}}}^\omega$  with a Poincaré structure corresponding to the module with genuine involution  $M_{A^{\text{op}}} := A^{\text{op}}$ ,  $N_{A^{\text{op}}} := \overline{P}$ . It remains to exhibit a natural equivalence

$$\eta: (\mathfrak{Q}_A \otimes \mathfrak{Q}_{A^{\text{op}}}) \xrightarrow{\sim} \text{ev}^* \mathfrak{Q}_R \quad (6.22)$$

of [quadratic] functors  $\text{Mod}_A^\omega \otimes \text{Mod}_{A^{\text{op}}}^\omega \rightarrow \text{Sp}$ . By [Cal+20a, Theorem 3.3.1], it suffices to exhibit equivalences on the bilinear and linear parts of (6.22) which glue compatibly. By Proposition 6.4(3), on linear parts, it suffices to exhibit an  $A \otimes_R A^{\text{op}}$ -linear equivalence

$$\text{hom}_R(A, R^{\varphi C_2}) \simeq N_A \otimes_{R^{\varphi C_2}} N_{A^{\text{op}}}$$

L: todo: an  $R$ -linear enhancement of Proposition 3.1.16?

L: continue. discuss consequences of being Morita trivial (distinguished ‘point’ in  $N_A$ )? can we identify  $\overline{P}$ ?

L: In what other cases does a lift exist trivially? wrt Tate Poincaré structure, suffices to take a trivialization of  $A$ ?

L: rewrite for scheme with involution

and on bilinear parts, it suffices to exhibit an  $(A \otimes_R A^{\text{op}})^{\otimes_{R^2}}$ -linear equivalence

$$\text{hom}_{R \otimes R}(A \otimes_R A, R) \simeq M_A \otimes_R M_{A^{\text{op}}}$$

which glue compatibly. This follows from the definitions, concluding the proof.  $\square$

**Proposition 6.23.** *Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring. Let  $(\mathcal{C}, \mathcal{Y})$  be an invertible idempotent-complete  $R$ -linear Poincaré  $\infty$ -category. Suppose given a Poincaré object  $(P, q)$  of  $(\mathcal{C}, \mathcal{Y})$  so that  $P$  is a generator for  $\mathcal{C}$ . Then  $(\mathcal{C}, \mathcal{Y})$  is of the form  $(\text{Mod}_A^\omega, \mathcal{Y}_A)$  for some Azumaya algebra over  $R$  with genuine involution.*

L: later:  
add case of  
schemes with  
involution

**Corollary 6.24.** *Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring. Let  $(\mathcal{C}, \mathcal{Y})$  be an invertible idempotent-complete  $R$ -linear Poincaré  $\infty$ -category. Then  $(\mathcal{C}, \mathcal{Y})$  is of the form  $(\text{Mod}_A^\omega, \mathcal{Y}_A)$  for some Azumaya algebra over  $R$  with genuine involution.*

L: connectivity  
hypothesis on  
 $R$ ?

*Proof.* By Proposition 6.23, it suffices to exhibit a Poincaré object  $(P, q)$  whose underlying object is a generator of  $\mathcal{C}$ . By [AG14, Theorem 6.1],  $\mathcal{C}$  has a generator  $G$ . Now  $G \oplus D_{\mathcal{Y}}G$  promotes canonically to a Poincaré object of  $(\mathcal{C}, \mathcal{Y})$  by [Cal+20a, Proposition 2.2.5].  $\square$

*Proof of Proposition 6.23.* Similar to Example 6.18  $\square$

L: Need to  
show  $\mathcal{C}$  (or  
 $\text{Ind}(\mathcal{C})$ ) defines  
a  $R$ -linear  
category with  
descent?

## 6.2 Stacks associated to Poincaré- $\infty$ -categories

**Notation 6.25.** Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring. There is a functor

$$\begin{aligned} \mathbb{E}_\infty \text{Alg}_{R/-}^{BC_2} &\rightarrow \text{CAlg}_{R/-}^P \\ S &\mapsto (S, S \rightarrow R^{\varphi C_2} \otimes_R S \rightarrow S^{tC_2}) =: (R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2}) \otimes S, \end{aligned}$$

where the map  $R^{\varphi C_2} \otimes_R S \rightarrow R^{\varphi C_2} \otimes_{R^{tC_2}} S^{tC_2} \rightarrow R^{tC_2} \otimes_{R^{tC_2}} S^{tC_2} \simeq S^{tC_2}$  is given by base change along the Tate-valued norm composed with the structure map  $R^{\varphi C_2} \rightarrow R^{tC_2}$ . Composing the aforementioned functor with the functor that sends a Poincaré ring to its category of compact modules equipped with the canonical Poincaré structure defines a functor

$$\text{Mod}^P: \mathbb{E}_\infty \text{Alg}_{R/-}^{BC_2} \rightarrow \mathbb{E}_\infty \text{Alg} \left( \text{Cat}_{\infty, \text{idem}_R}^P \right).$$

**Notation 6.26.** Let  $X$  be a scheme with an involution  $\sigma$  and let  $\pi: X \rightarrow Y$  exhibit  $Y$  as a good quotient of  $X$ . If  $j: U \rightarrow Y$  is flat, let us write  $\pi^*U$  for the tuple  $(X \times_Y U, j^*(\sigma), U, j^*(\pi))$  of Remark 5.17. Then the assignment  $(j: U \rightarrow Y) \mapsto (\text{Mod}_{j^*X}^\omega, \mathcal{Y}_{j^*\sigma})$  defines a functor

$$\text{Mod}^P: \acute{\text{E}}t_Y^{\text{op}} \rightarrow \mathbb{E}_\infty \text{Alg} \left( \text{Mod}_{(\text{Mod}_X^\omega, \mathcal{Y}_\sigma)}(\text{Cat}_{\infty, \text{idem}}^P) \right).$$

**Proposition 6.27.** *Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring and assume that  $R^{\varphi C_2}$  and  $R$  are connective. Then the assignment of Notation 6.25 is a hypersheaf on the small étale site of  $R$ .*

**Corollary 6.28.** *Let  $(R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$  be a Poincaré ring and assume that  $R^{\varphi C_2}$  and  $R$  are connective. Write  $\text{PnBr}$  for the composite functor  $\acute{\text{E}}t_R \xrightarrow{\text{Mod}^P} \mathbb{E}_\infty \text{Alg} \left( \text{Cat}_{\infty, \text{idem}_R}^P \right) \xrightarrow{\text{PnBr}} \text{Sp}_{\geq 0}$ , where  $\text{Mod}^P$  is from Notation 6.25 and  $\text{PnBr}$  is Definition 6.1. Then  $\text{PnBr}$  is an étale sheaf.*

*Proof.* Follows from Proposition 6.27 and the fact that  $\text{PnBr}$  preserves all limits.  $\square$

**Proposition 6.29.** *Let  $(X, \lambda, Y, \pi)$  be a scheme with involution  $X$  and a good quotient  $Y$ . Then the assignment of Notation 6.26 is a hypersheaf on the small étale site of  $Y$ .*

*Proof of Proposition 6.27.* For now, use the notational shorthand  $R^P = (R, R \rightarrow R^{\varphi C_2} \rightarrow R^{tC_2})$ . Since limits in categories of algebras and modules are computed at the level of underlying objects, it suffices to show that the functor sends an étale hypercovering  $j_\bullet: S \rightarrow T^\bullet$  to a limit diagram in  $\text{Cat}_{\infty, \text{idem}}^P$ . By Proposition 6.1.4 of [Cal+20a], it suffices to show that the relevant diagram is a limit diagram in  $\text{Cat}_{\infty, \text{idem}}^h$ . The proof of Lemma 5.4 in [AG14] implies that the diagram defines a limit diagram on underlying  $\infty$ -categories. Thus by Remark 6.1.3 of [Cal+20a], it suffices to show that  $j_\bullet^*: \text{Mod}_{R^e \otimes_{R^{C_2}} S}^\omega \rightarrow \text{Mod}_{R^e \otimes_{R^{C_2}} T^\bullet}^\omega$  induces an equivalence  $\mathcal{Y}_{R^P \otimes S} \xrightarrow{\sim} \lim_{\Delta} \mathcal{Y}_{R^P \otimes T^\bullet} \circ (j_\bullet^*)^{\text{op}}$  of quadratic functors  $\text{Mod}_{R^e \otimes_{R^{C_2}} S}^{\omega, \text{op}} \rightarrow \text{Sp}$ . This follows from our assumption on  $S \rightarrow T^\bullet$  and [Cal+20a, Theorem 3.3.1].  $\square$

L: I think  
Propositions  
?? and 6.29  
have essen-  
tially the same  
proof—present  
them together?

L: recall  
étale hyper-  
sheaf/hypercov

## 7 Poincaré schemes

**Definition 7.1.** Let APS be the  $(\infty, 1)$ -category defined by the pullback

$$\begin{array}{ccc} \text{APS} & \longrightarrow & \text{Fun}(\Delta^2, \text{CAlg}(\text{Sp})) \\ \downarrow & & \downarrow d_1^* \\ \text{CAlg}(\text{Sp}^{BC_2}) & \xrightarrow{U(-) \rightarrow (-)^{tC_2}} & \text{Fun}(\Delta^1, \text{CAlg}(\text{Sp})) \end{array}$$

where  $U : \text{Sp}^{BC_2} \rightarrow \text{Sp}$  is the functor which forgets the  $C_2$ -action.

**Definition 7.2.** Define the category of affine Hermetian schemes, denoted AHS, to be the infinity category given by the Grothendieck construction applied to the functor

$$\text{CAlg}(\text{Sp})^{op} \rightarrow \text{Cat}_\infty$$

given by sending a ring  $R$  to the category  $\text{CAlg}(\text{Mod}_{NR})$  of  $\mathbb{E}_\infty$  algebras in modules with genuine involutions over  $R$ . Then define the category of affine Poincaré schemes, denoted by APS, to be the full subcategory of AHS spanned by the pairs  $(R, M)$  where  $M \in \text{CAlg}(\text{Mod}_{NR})$  is invertible.

We record here a few structural results about this category.

**Theorem 7.3.** *The following statements about APS hold:*

1. *The category APS is a cocomplete and symmetric monoidal infinite category;*
2. *the pullback diagram above is homotopy Cartesian;*
3. *the functor  $\text{APS} \rightarrow \text{CAlg}(\text{Sp}^{BC_2})$  is symmetric monoidal and (co)continuous;*
4. *the functor  $\text{APS} \rightarrow \text{CAlg}(\text{Sp})^{\Delta^2}$  is lax symmetric monoidal;*
5. *and the functor  $\text{APS} \rightarrow \text{CAlg}(\text{Sp})^{\Delta^2} \xrightarrow{\text{ev}_{[1]}} \text{CAlg}(\text{Sp})$  is symmetric monoidal.*

*Proof.* For (2) it is enough to show that  $d_1^*$  is an categorical fibration which follows from [Lur09, Corollary 2.3.2.5] and [Lur09, Corollary 2.4.6.5]. In fact  $d_1^*$  is a left fibration by [(taking  $p = id$ ,  $i = d_1 : \Delta^1 \rightarrow \Delta^2$ )]. There is a (pseudo-)functor

$$\begin{aligned} F : \text{Fun}(\Delta^1, \text{CAlg}(\text{Sp})) &\rightarrow \text{Cat}_\infty \\ (\varphi : A \rightarrow B) &\mapsto ((\text{CAlg}(\text{Sp})_{A/-/B})_{/\varphi}) \end{aligned}$$

which sends a square

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ \downarrow & & \downarrow \\ C & \xrightarrow{\psi} & D \end{array} \quad (7.4)$$

regarded as a morphism from  $\varphi$  to  $\psi$ , to the functor

$$\begin{aligned} (\text{CAlg}(\text{Sp})_{A/-/B})_{/\varphi} &\rightarrow (\text{CAlg}(\text{Sp})_{C/-/D})_{/\psi} \\ (A \rightarrow R \rightarrow B) &\mapsto C \simeq A \otimes_A C \xrightarrow{\varphi \otimes \text{id}_C} B \otimes_A C \rightarrow D \end{aligned} \quad (7.5)$$

where  $B \otimes_A C \rightarrow D$  is the canonical map induced by the commuting square (7.4). The functor  $F$  classifies the cocartesian fibration  $d_1^*$ .

N: I keep trying to make this work but the technical details are actively killing me. Better, I think, to use the following definition instead.

L: Is this reference correct? The conclusion asserts that some map of simplicial sets is a *categorical* fibration. The following argument is ‘sketchy’—depending on how precise we want to be about quascategories, we may want to argue with left/right anodyne maps instead.

N: To prove 2 we only need that  $d_1^*$  is a categorical fibration, although

For (3), let  $p : K \rightarrow \text{APS}$  be a map of simplicial sets,  $K$  a small simplicial set. Suppose the  $K^\triangleright \rightarrow \text{APS}$  be an extension such that  $K^\triangleright \rightarrow \text{APS} \rightarrow \text{CAlg}(\text{Sp}^{BC_2})$  is a colimit diagram. By [Lur09, Proposition 2.4.3.2] the diagram

$$\begin{array}{ccc} \text{APS}_{p/} & \longrightarrow & \text{CAlg}(\text{Sp})_{p/-}^{\Delta^2} \\ \downarrow & & \downarrow \\ \text{CAlg}(\text{Sp}^{BC_2})_{p/} & \longrightarrow & \text{CAlg}(\text{Sp})_{p/-}^{\Delta^1} \end{array}$$

is again homotopy cartesian. Then

$$\begin{aligned} \text{hom}_{\text{APS}}(p(\infty), -) &\simeq \text{hom}_{\text{CAlg}(\text{Sp}^{BC_2})}(p(\infty), -) \times_{\text{hom}_{\text{CAlg}(\text{Sp})}^{\Delta^1}(p(\infty), -)} \text{hom}_{\text{CAlg}(\text{Sp})}^{\Delta^2}(p(\infty)) \\ &\simeq \end{aligned}$$

□

We will denote elements of APS by  $\underline{A} = (A, s : A^{\Phi C_2} \rightarrow A^{tC_2})$ . Here  $s : A^{\Phi C_2} \rightarrow A^{tC_2}$  is the image of  $\underline{A}$  under the top horizontal map above. The use of the notation  $A^{\Phi C_2}$  is justified by the following.

**Lemma 7.6.** *Let  $\text{APS} \rightarrow \text{CAlg}(\text{Sp})$  be the composition of the functors*

$$\text{APS} \rightarrow \text{Fun}(\Delta^2, \text{CAlg}(\text{Sp})) \xrightarrow{ev_{[1]}} \text{CAlg}(\text{Sp}).$$

*Then this functor factors as a composition  $\text{APS} \rightarrow \text{CAlg}(\text{Sp}^{C_2}) \xrightarrow{(-)^{\Phi C_2}} \text{CAlg}(\text{Sp})$ .*

*Proof.* The commutativity of the diagram

$$\begin{array}{ccccc} & & \text{Fun}(\Delta^2, \text{CAlg}(\text{Sp})) & & \\ & & \downarrow d_1^* & \searrow d_0^* & \\ & & & & \text{Fun}(\Delta^1, \text{CAlg}(\text{Sp})) \\ & \text{CAlg}(\text{Sp}^{BC_2}) \xrightarrow{U(-) \rightarrow (-)^{tC_2}} & \text{Fun}(\Delta^1, \text{CAlg}(\text{Sp})) & \xrightarrow{ev_{[1]}} & \downarrow ev_{[1]} \\ & \searrow id & & \searrow (-)^{tC_2} & \downarrow \\ & & \text{CAlg}(\text{Sp}^{BC_2}) & \longrightarrow & \text{CAlg}(\text{Sp}) \end{array}$$

induces a functor on the pullback infinity categories  $\text{APS} \rightarrow \text{CAlg}(\text{Sp}^{C_2})$  which makes the corresponding cube commute. The functor  $ev_{[1]} : \text{Fun}(\Delta^2, \text{CAlg}(\text{Sp})) \rightarrow \text{CAlg}(\text{Sp})$  factors through  $d_0^*$  and so  $\text{APS} \rightarrow \text{Fun}(\Delta^2, \text{CAlg}(\text{Sp})) \rightarrow \text{CAlg}(\text{Sp})$  is equivalent to the composition

$$\text{APS} \rightarrow \text{CAlg}(\text{Sp}^{C_2}) \rightarrow \text{Fun}(\Delta^1, \text{CAlg}(\text{Sp})) \rightarrow \text{CAlg}(\text{Sp})$$

and the composition of the last two maps is the geometric fixed point functor as desired. □

The following Lemma gives the justification of the name Poincaré scheme.

**Construction 7.7.** We shall construct a functor

$$\text{Perf}^{\text{Pn}} : \text{APS} \rightarrow \text{Cat}_{\infty}^{\text{Pn}}$$

to the category of Poincaré infinity categories.

Recall that  $\text{Cat}_{\infty}^{\text{h}} \rightarrow (\text{Cat}_{\infty}^{\text{ex}})^{\text{op}}$  is a cocartesian fibration [Cal+20a, §1.4.] We will first construct a map of cocartesian fibrations

$$\begin{array}{ccc} \text{APS} & \xrightarrow{\quad \quad \quad} & \text{Cat}_{\infty}^{\text{h}} \\ \downarrow & & \downarrow \\ \text{CAlg}(\text{Sp}^{BC_2}) & \longrightarrow & (\text{Cat}_{\infty}^{\text{ex}})^{\text{op}} \end{array}, \quad (7.8)$$

L: For symmetric monoidal structure—maybe want to swap out  $\text{Mod}_{NA}$  for  $\text{CAlg}_{NA}$ ?

then show that the dotted arrow factors through the subcategory  $\text{Cat}_\infty^p \subseteq \text{Cat}_\infty^h$ . To construct a map of cartesian fibrations, it suffices to exhibit a natural transformation of classifying functors. Unraveling the definitions, by Theorem 3.2.13 of [Cal+20a] we must exhibit for each  $A \in \text{CAlg}(\text{Sp})^{BC_2}$ , a functor

$$(\text{CAlg}(\text{Sp})_{A/-/A^{tC_2}})_{/\varphi} \rightarrow \text{Mod}_{N^{C_2}(A^e)}(\text{Sp}^{C_2}) \quad (7.9)$$

(where  $\varphi: A \rightarrow A^{tC_2}$  is the Tate-valued Frobenius and  $N^{C_2}$  is the Hill–Hopkins–Ravenel norm) which is natural in  $A$ .

That the resulting functor factors through the subcategory  $\text{Cat}_\infty^p$  follows from Proposition 3.1.3 and Lemma 3.3.3 of *loc. cit.*

**Lemma 7.10.** *The functor of Construction 7.7 is symmetric monoidal and has essential image the subcategory spanned by objects  $(\text{Perf}(R), \mathfrak{P})$  which are  $\mathbb{E}_\infty$ -algebras.*

**Definition 7.11.** A map  $f: A \rightarrow B \in \text{APS}$  is faithfully flat if the underlying map  $f: A \rightarrow B$  is faithfully flat and the map  $f^{\Phi C_2}: A^{\Phi C_2} \rightarrow B^{\Phi C_2}$  is also faithfully flat.

**Lemma 7.12.** *The fpqc covers on APS form a Grothendieck site.*

## References

- [AG14] Benjamin Antieau and David Gepner. “Brauer groups and étale cohomology in derived algebraic geometry”. In: *Geom. Topol.* 18.2 (2014), pp. 1149–1244. ISSN: 1465-3060,1364-0380. DOI: [10.2140/gt.2014.18.1149](https://doi.org/10.2140/gt.2014.18.1149). URL: <https://doi.org/10.2140/gt.2014.18.1149>.
- [AGH21] Gabriel Angelini-Knoll, Teena Gerhardt, and Michael Hill. *Real topological Hochschild homology via the norm and Real Witt vectors*. 2021. arXiv: [2111.06970](https://arxiv.org/abs/2111.06970) [math.AT]. URL: <https://arxiv.org/abs/2111.06970>.
- [Alb61] A. Adrian Albert. *Structure of algebras*. Vol. Vol. XXIV. American Mathematical Society Colloquium Publications. Revised printing. American Mathematical Society, Providence, RI, 1961, pp. xi+210.
- [Bar+22] Clark Barwick, Saul Glasman, Akhil Mathew, and Thomas Nikolaus. *K-theory and polynomial functors*. 2022. arXiv: [2102.00936](https://arxiv.org/abs/2102.00936) [math.KT]. URL: <https://arxiv.org/abs/2102.00936>.
- [BRS12] Andrew Baker, Birgit Richter, and Markus Szymik. “Brauer groups for commutative  $S$ -algebras”. In: *J. Pure Appl. Algebra* 216.11 (2012), pp. 2361–2376. ISSN: 0022-4049,1873-1376. DOI: [10.1016/j.jpaa.2012.03.001](https://doi.org/10.1016/j.jpaa.2012.03.001). URL: <https://doi.org/10.1016/j.jpaa.2012.03.001>.
- [Cal+20a] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. *Hermitian K-theory for stable  $\infty$ -categories I: Foundations*. 2020. DOI: [10.48550/ARXIV.2009.07223](https://arxiv.org/abs/2009.07223). URL: <https://arxiv.org/abs/2009.07223>.
- [Cal+20b] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. *Hermitian K-theory for stable  $\infty$ -categories II: Cobordism categories and additivity*. 2020. DOI: [10.48550/ARXIV.2009.07224](https://arxiv.org/abs/2009.07224). URL: <https://arxiv.org/abs/2009.07224>.
- [Cal+21] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. *Hermitian K-theory for stable  $\infty$ -categories III: Grothendieck–Witt groups of rings*. 2021. arXiv: [2009.07225](https://arxiv.org/abs/2009.07225) [math.KT].
- [CHN24] Baptiste Calmès, Yonatan Harpaz, and Denis Nardin. *A motivic spectrum representing hermitian K-theory*. 2024. arXiv: [2402.15136](https://arxiv.org/abs/2402.15136) [math.KT]. URL: <https://arxiv.org/abs/2402.15136>.
- [Fau03] H. Fausk. “Picard groups of derived categories”. In: *J. Pure Appl. Algebra* 180.3 (2003), pp. 251–261. ISSN: 0022-4049,1873-1376. DOI: [10.1016/S0022-4049\(02\)00145-7](https://doi.org/10.1016/S0022-4049(02)00145-7). URL: [https://doi.org/10.1016/S0022-4049\(02\)00145-7](https://doi.org/10.1016/S0022-4049(02)00145-7).

- [FW20] Uriya A. First and Ben Williams. “Involutions of Azumaya Algebras”. en. In: *DOCUMENTA MATHEMATICA* Vol 25 (2020 2020), p. 527–633. DOI: [10.25537/DM.2020V25.527-633](https://doi.org/10.25537/DM.2020V25.527-633). URL: <https://www.elibm.org/article/10012038>.
- [Gab81] Ofer Gabber. “Some theorems on Azumaya algebras”. In: *The Brauer group (Sem., Les Plans-sur-Bex, 1980)*. Vol. 844. Lecture Notes in Math. Springer, Berlin, 1981, pp. 129–209. ISBN: 3-540-10562-X.
- [Lur09] Jacob Lurie. *Higher topos theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925. ISBN: 978-0-691-14049-0; 0-691-14049-9. DOI: [10.1515/9781400830558](https://doi.org/10.1515/9781400830558). URL: <https://doi.org/10.1515/9781400830558>.
- [Lur17] Jacob Lurie. *Higher algebra*. Manuscript available at <https://www.math.ias.edu/~lurie/papers/HA.pdf>. 2017.
- [Lur18] Jacob Lurie. “Elliptic cohomology II: orientations”. In: *available at https://www.math.ias.edu/~lurie/papers/Elliptic-II.pdf* (version dated 26 April 2018).
- [Lur24] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2024.
- [Nik23] Thomas Nikolaus. “Frobenius homomorphisms in higher algebra”. In: *ICM—International Congress of Mathematicians. Vol. 4. Sections 5–8*. EMS Press, Berlin, [2023] ©2023, pp. 2826–2854. ISBN: 978-3-98547-062-4; 978-3-98547-562-9; 978-3-98547-058-7.
- [NS18] Thomas Nikolaus and Peter Scholze. “On topological cyclic homology”. In: *Acta Math.* 221.2 (2018), pp. 203–409. ISSN: 0001-5962. DOI: [10.4310/ACTA.2018.v221.n2.a1](https://doi.org/10.4310/ACTA.2018.v221.n2.a1). URL: <https://doi-org.proxyiub.uits.iu.edu/10.4310/ACTA.2018.v221.n2.a1>.
- [PS92] R. Parimala and V. Srinivas. “Analogues of the Brauer group for algebras with involution”. In: *Duke Math. J.* 66.2 (1992), pp. 207–237. ISSN: 0012-7094,1547-7398. DOI: [10.1215/S0012-7094-92-06606-3](https://doi.org/10.1215/S0012-7094-92-06606-3). URL: <https://doi.org/10.1215/S0012-7094-92-06606-3>.
- [Sal78] David J. Saltman. “Azumaya algebras with involution”. In: *J. Algebra* 52.2 (1978), pp. 526–539. ISSN: 0021-8693. DOI: [10.1016/0021-8693\(78\)90253-3](https://doi.org/10.1016/0021-8693(78)90253-3). URL: [https://doi.org/10.1016/0021-8693\(78\)90253-3](https://doi.org/10.1016/0021-8693(78)90253-3).
- [Toë12] Bertrand Toën. “Derived Azumaya algebras and generators for twisted derived categories”. In: *Invent. Math.* 189.3 (2012), pp. 581–652. ISSN: 0020-9910,1432-1297. DOI: [10.1007/s00222-011-0372-1](https://doi.org/10.1007/s00222-011-0372-1). URL: <https://doi.org/10.1007/s00222-011-0372-1>.
- [Yan23] Lucy Yang. *On normed  $\mathbb{E}_\infty$ -rings in genuine equivariant  $C_p$ -spectra*. 2023. arXiv: [2308.16107](https://arxiv.org/abs/2308.16107) [[math.AT](https://arxiv.org/abs/2308.16107)].