

POTENZE A ESPONENTE INTERO
$$Z = \{..., -3, -2, -1, 0, 1, 2, 3, ...\}$$
 $A^{m} = \begin{cases} a^{m} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{a.m.} \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1}{a^{m}} & \text{se } m \ge 0 \\ 1 & \text{se } m \ge 0 \end{cases}$
 $A^{m} = \begin{cases} \frac{1$

V2 = 1,41	42135				
V = - 7					
A D O A C A M A D A A A A		-1,5	1 4000 (41145)	C.1 a a. 11 m 3.1	3
APPROSSIMAZIONI	SUCCESSIVE S	1 02	Arrossipazioni	Successive DI	7
1			31		
			11		
1,4			31,4		
			31,41		
1,41					
			34,414		
1,414			1 1112		
1,4142			3,4142		
			31,41421		
1,41421			3 '		
1 1 1			 		
			7		
De demostra	che quesa	succe	nione si av	vicine semp	e di
più (comer	se) o un.	unic s	ben determin	nots muners re	role
10 1		JZ			
the denotion	mo on 3				