Initialize functions


```
(*Ensure that variables are defined globally*)
SetOptions[EvaluationNotebook[], CellContext → "Global`"]
NotebookOpen[NotebookDirectory[] <> "GB-strain-scattering_init.nb"];
NotebookEvaluate[NotebookDirectory[] <> "GB-strain-scattering_init.nb"];
```


Define material values and phonon dispersion parameters

```
(*### Material values for Si ###*)
(*#### crystal properties ####*)
vs = 6084. (*[m/s] average speed of sound*);
V = 2 * 10^{-29} (*[m^3] \text{ volume of atom*});
n = 2(*# atoms per primitive unit cell (N in paper)*);
γ = 1(*Gruneissen parameter*);
v = 0.27(*Poisson's ratio*);
kmax = k0[V*n] // N(*[m^{-1}] edge of FBZ*);
\omega D = vs kmax(*[s^{-1}] Debye frequency*);
disp = "Debye";
\omega k[k] := vsk (*Debye dispersion relation*);
vg[k_] := vs (*phonon group velocity*);
(*#### phonon-phonon ####**)
C1 = 2.69 * 10^{-19} (*[s/K]*);
C2 = 167(*[K]*);
(*#### point defect ####*)
C3 = 1.81 * 10^{-45} (*[s^3]*);
(*#### microstructure ####*)
b = (V n)^{1/3} (*[m] Burger's vector*);
dGS = 350 * 10^{-9} (*[m] average grain size*);
n1D = 3 / dGS // N(*[m^{-1}]) number density of GBs *);
d = 3 * 10^{-9} (*[m] GB dislocation spacing (D in paper)*);
```

Diffraction peaks in $\Gamma_{qbs}(k)$

```
\texttt{rMakeList}[\theta\_, \phi\_] := \texttt{Table}\big[\big\{\omega\big/\omega\mathsf{D}, \, \texttt{rGBS}[\omega\,/\,\mathsf{vs},\,\theta,\,\phi] \, \star \, 10^{-9}\big\}, \, \big\{\omega,\,0,\,\omega\mathsf{D},\,10^{10}\big\}\big];
(*\theta=\pi/2 \phi=0 \text{ is phonon incident normal to the GB.*})
rlistGBS = rMakeList[\pi/2, 0];
ListPlot[{rlistGBS},
  PlotStyle → {Black},
  PlotRange \rightarrow \{\{0, 1\}, \{0, 40\}\},\
  Joined → True,
  FrameLabel \rightarrow \{ "k/k_{\text{max}}", "\Gamma_{\text{gbs}}(\mathbf{k} = k\hat{\mathbf{x}}) \text{ (ns}^{-1}) " \} ]
ListPlot[{rlistGBS},
  Joined → True,
  FrameStyle → Directive[FontFamily → "Times New Roman", FontSize → 20, Black],
  PlotStyle → Directive[Black, Thickness[0.01]],
  PlotRange \rightarrow \{\{0, 0.35\}, \{0, 7\}\}\}
```






Calculating $\tau_{\rm gbs}(\omega)^{-1}$

```
(*A convergence test was perfomed on \delta k by caluclating \kappa L at different \delta k. \delta k =
 2*10<sup>8</sup> was found to be fully converged. This
  calculation takes ~20 min on most personal computers*)
Clear[d]
τGBSList = {disp};
\delta k = 2. * 10^8;
For [d = 1.0 * 10^{-9}, d \le 8.0 * 10^{-9}, d = d * 2.0,
 Print[d];
 \tau GBSListd = Table[\{\omega / vs, \tau GBS\Theta \phi Integrate[\omega / vs, 90]\}, \{\omega, vs \delta k, \omega D, vs \delta k\}];
 AppendTo[\tauGBSList, {d, \thetacalc[b, d], \tauGBSListd}]
1. \times 10^{-9}
2. \times 10^{-9}
4. \times 10^{-9}
8. \times 10^{-9}
(*#### Exports τGBS results ###*)
Export[NotebookDirectory[] <> "../Results/tauSTGB_results.m", tGBSList]
/Users/rileyhanus/Desktop/Papers/Phonon_diffraction/Scripts/FOR_DISTRIBUTION/
  Phonon-scattering-and-transport-in-polycyrstals/core/../Results/tauSTGB_results.
(*#### Imports the last exported τGBS result ####*)
τGBSList = Import[NotebookDirectory[] <> "../Results/tauSTGB_results.m"];
Plotting \tau_{abs}(\omega) for a STGB
"STGB"
Show[ListLogPlot[{\tauGBExtract[\tauGBSList, 2],
   τGBExtract[τGBSList, 3], τGBExtract[τGBSList, 4], τGBExtract[τGBSList, 5]},
  PlotStyle -> {Directive[RGBColor[0., 0.5, 0.5], Thickness[0.0075]],
    Directive[RGBColor[0.7, 0., 0.5], Thickness[0.0075]],
    Directive[RGBColor[0., 0, 0.7], Thickness[0.0075]],
    Directive[Black, Thickness[0.0075]]},
  FrameLabel \rightarrow {Style["", 16], Style["\tau_{gbs}(\omega) (ns)", 16]},
  PlotRange \rightarrow \{\{0, 1\}, \{0.01, 5\}\},\
  Joined → True,
  PlotRange → All]]
Show
 ListPlot[{rGBExtract[rGBSList, 2],
```

```
rGBExtract[tGBSList, 3], rGBExtract[tGBSList, 4], rGBExtract[tGBSList, 5]},
  PlotStyle -> {Directive[RGBColor[0., 0.5, 0.5], Thickness[0.0075]],
     Directive[RGBColor[0.7, 0., 0.5], Thickness[0.0075]],
     Directive[RGBColor[0., 0, 0.7], Thickness[0.0075]],
     Directive[Black, Thickness[0.0075]]},
   FrameLabel \rightarrow {Style["\omega/\omega_D", 16], Style["\tau_{gbs}(\omega)^{-1} (ns<sup>-1</sup>)", 16]},
  PlotRange \rightarrow \{\{0, 1\}, \{0, 45\}\},\
   Joined → True,
  InterpolationOrder \rightarrow 3,
  PlotRange → All],
 Plot[\{\tau \text{invEmp}[\omega * \omega D, 1 * 10^{-9}, b, n1D] * 10^{-9}, 
    \tau inv Emp[\omega * \omega D, 2 * 10^{-9}, b, n1D] * 10^{-9}, \tau inv Emp[\omega * \omega D, 4 * 10^{-9}, b, n1D] * 10^{-9},
    \tau inv Emp[\omega * \omega D, 8 * 10^{-9}, b, n1D] * 10^{-9} \}, \{\omega, 0, 1\},
  PlotStyle → {Directive[Dashing[{.01, 0.03}],
       RGBColor[0, 0.2, 0.2], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], RGBColor[0.3, 0., 0.1], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], RGBColor[0.6, 0.6, 1], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], GrayLevel[0.7], Thickness[0.0075]]}]]
Show[
 ListPlot[{rGBExtract[rGBSList, 2], rGBExtract[rGBSList, 3],
    ΓGBExtract[τGBSList, 4], ΓGBExtract[τGBSList, 5]},
  PlotStyle -> {Directive[RGBColor[0., 0.5, 0.5], Thickness[0.01]],
     Directive[RGBColor[0.7, 0., 0.5], Thickness[0.01]],
     Directive[RGBColor[0., 0, 0.7], Thickness[0.01]],
     Directive[Black, Thickness[0.01]]},
   FrameLabel → {Style["", 16], Style["", 16]},
  PlotRange \rightarrow \{\{0, 0.4\}, \{0, 7\}\},\
  ImageSize → 250,
   Joined → True,
  InterpolationOrder \rightarrow 3,
  PlotRange → All],
 Plot[\{\tau inv Emp[\omega * \omega D, 1 * 10^{-9}, b, n1D] * 10^{-9}, 
    \tauinvEmp[\omega * \omegaD, 2 * 10<sup>-9</sup>, b, n1D] * 10<sup>-9</sup>, \tauinvEmp[\omega * \omegaD, 4 * 10<sup>-9</sup>, b, n1D] * 10<sup>-9</sup>,
    \tau invEmp[\omega * \omega D, 8 * 10^{-9}, b, n1D] * 10^{-9}\}, \{\omega, 0, 1\},
  PlotStyle → {Directive[Dashing[{.01, 0.03}],
       RGBColor[0, 0.2, 0.2], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], RGBColor[0.3, 0., 0.1], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], RGBColor[0.6, 0.6, 1], Thickness[0.0075]],
     Directive[Dashing[{.01, 0.03}], GrayLevel[0.7], Thickness[0.0075]]}]]
STGB
```



Convert $\tau(\omega)$ to $t(\omega)$ and plot

```
Show[ListPlot[{tGBExtract[rGBSList, 2], tGBExtract[rGBSList, 3],
    tGBExtract[\tauGBSList, 4], tGBExtract[\tauGBSList, 5]},
   PlotRange \rightarrow \{\{0, 1\}, \{0., 1\}\},\
   FrameLabel \rightarrow \{ "\omega/\omega_D", "Transmissivity, t(\omega)" \}, 
   Joined → True,
   PlotStyle -> {Directive[RGBColor[0., 0.5, 0.5], Thickness[0.0075]],
      Directive[RGBColor[0.7, 0., 0.5], Thickness[0.0075]],
      Directive[RGBColor[0., 0, 0.7], Thickness[0.0075]],
      Directive[Black, Thickness[0.0075]]}],
 Plot[{tEmp[\omega * \omega D, 1 * 10<sup>-9</sup>, b, n1D], tEmp[\omega * \omega D, 2 * 10<sup>-9</sup>, b, n1D],
    \mathsf{tEmp} \big[\omega * \omega \mathsf{D}, \, 4 * 10^{-9}, \, \mathsf{b}, \, \mathsf{n1D} \big], \, \mathsf{tEmp} \big[\omega * \omega \mathsf{D}, \, 8 * 10^{-9}, \, \mathsf{b}, \, \mathsf{n1D} \big] \big\}, \, \{\omega, \, 0, \, 1\},
   PlotStyle → {Directive[Dashing[{.01, 0.03}],
       RGBColor[0, 0.2, 0.2], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], RGBColor[0.3, 0., 0.1], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], RGBColor[0.6, 0.6, 1], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], GrayLevel[0.7], Thickness[0.0075]]}]]
Fransmissivity, t(\omega)
    0.8
    0.6
    0.4
    0.0
                 0.2
                            0.4
                                       0.6
                                                  0.8
                                                             1.0
                                \omega/\omega_D
```

Calculating κ_L

```
Timing[
 κTList = {disp};
 For[i = 2, i \le 5, i++,
  Print[tGBSList[[i, 1]] // N];
  \kappa T = \{\};
  For [T = 5, T \le 500, T = T * 1.1,
   AppendTo[xT, {T, xLCalc[rTotCalc[rGBSList[[i, 3]], T], T]}]];
  AppendTo[\kappaTList, {\tauGBSList[[i, 1]] // N, \tauGBSList[[i, 2]], \kappaT}]]]
```

```
1. \times 10^{-9}
2. \times 10^{-9}
\textbf{4.}\times\textbf{10}^{-9}
8. \times 10^{-9}
{0.28003, Null}
Show
 ListLogLogPlot[{xTList[[2, 3]], xTList[[3, 3]], xTList[[4, 3]], xTList[[5, 3]]},
   FrameTicks → Automatic,
   Joined → True,
   PlotStyle →
     {RGBColor[0., 0.5, 0.5], RGBColor[0.7, 0., 0.5], RGBColor[0., 0, 0.7], Black},
   PlotRange \rightarrow \{\{5, 500\}, \{0.1, 500\}\},\
   ImageSize \rightarrow 350],
  LogLogPlot[
   \{\kappa L[T, (\tau invEmp[\omega, 1*10^{-9}, b, n1D] + \tau ppModel[\omega, T, C1, C2]^{-1} + \tau PD[\omega, C3]^{-1})^{-1},
      vs, \omega / vs, \{\omega, \Theta, \omega D\}],
    \times L[T, (\tau invEmp[\omega, 2*10^{-9}, b, n1D] + \tau ppModel[\omega, T, C1, C2]^{-1} + \tau PD[\omega, C3]^{-1})^{-1},
      vs, \omega / vs, \{\omega, \Theta, \omega D\},
    \times L[T, (\tau invEmp[\omega, 4*10^{-9}, b, n1D] + \tau ppModel[\omega, T, C1, C2]^{-1} + \tau PD[\omega, C3]^{-1})^{-1},
      vs, \omega / vs, \{\omega, \Theta, \omega D\}],
    \kappa L[T, (\tau invEmp[\omega, 8*10^{-9}, b, n1D] + \tau ppModel[\omega, T, C1, C2]^{-1} + \tau PD[\omega, C3]^{-1})^{-1},
      vs, \omega / vs, \{\omega, 0, \omega D\}\}, \{T, 5, 500\},
   PlotStyle → {Directive[Dashing[{.01, 0.03}],
       RGBColor[0, 0.2, 0.2], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], RGBColor[0.3, 0., 0.1], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], RGBColor[0.6, 0.6, 1], Thickness[0.0075]],
      Directive[Dashing[{.01, 0.03}], GrayLevel[0.7], Thickness[0.0075]]]]
 ListLogLogPlot[{T2[18, {Tref, 8, 25}], T3[0.15, {Tref, 10, 35}]}, Joined → True]
     500
      50
\kappa_{\rm L} \; ({\rm Wm}^{-1}{\rm K}^{-1})
       5
     0.5
               10
                                 50
                                       100
                                                         500
```


T(K)


```
LineLegend[{Directive[Black, Thickness[0.0075]],
  Directive[RGBColor[0., 0, 0.7], Thickness[0.0075]],
  Directive[RGBColor[0.7, 0., 0.5], Thickness[0.0075]],
  Directive[RGBColor[0., 0.5, 0.5], Thickness[0.0075]]},
 {Style["
                        2", FontFamily → "Times New Roman", 18],
  Style["
                        5", FontFamily → "Times New Roman", 18],
  Style["
               2
                        10", FontFamily → "Times New Roman", 18],
  Style["
                        19", FontFamily → "Times New Roman", 18]}, Spacings → 0.3]
             2
5
10
```

Supplemental: Pole plots of $\Gamma^{-1}(k)$

```
d = 3 * 10^{-9};
For [i = 1, i \le 4, i++,
 Print["k=" <> ToString[i] <> "\frac{\pi}{d}"];
 Print[\tau PolarPlot[i\pi/d]]]
k=1\frac{\pi}{d}
        1.0
        0.5
\tau_{\rm gbs} \sin(\phi) (ns)
        0.0
     -0.5
             -1.5 -1.0 -0.5
                                          0.0
                                                    0.5
                                                             1.0
                                                                      1.5
                                   \tau_{\rm gbs} \cos(\phi) (ns)
```

 $k=2\frac{\pi}{d}$





 $k=4\frac{\pi}{d}$

